mal from the typical action of the extracts of exophthalmic goiter.

6. The tolerance of the serum of exophthalmic goiter apparently protects the animal against the action of the extracts of exophthalmic goiter.

CONCLUSION

There is a powerful depressor substance in saline extracts of exophthalmic goiters, and apparently the same substance is present in the blood of individuals suffering acutely from this disease.

ABSTRACT OF DISCUSSION

ON PAPERS OF DRS. WILSON, SMITH, SANFORD AND BLACKFORD

DR. E. T. BELL, Minneapolis: I should like to ask Dr. Wilson to explain his grouping of the toxic cases.

DR. A. W. HEWLETT, Ann Arbor, Mich.: Was the fall in blood-pressure from exophthalmic serum prevented by a previous injection of exophthalmic extract?

DR. A. H. SANFORD, Rochester, Minn.: Apparently tolerance to the depressor substance in an extract of exophthalmic goiter would prevent the fall in blood-pressure from a potent serum of an exophthalmic goiter patient. I use the word "potent" because only in the acute stage of intoxication is a depressor substance obtained. We have used altogether about twenty-six serums and about forty-nine extracts. Some of these serums were from patients with simple goiter. These serums contained no potent depressor substance. We also used serums from normal persons and from the horse. We found no drops in normal serums.

DR. A. W. HEWLETT, Ann Arbor, Mich.: We have injected the serums from patients with high blood-pressure into dogs and rabbits and in these cases there was a striking fall in blood-pressure. We were under the impression that this fall might occur after injections of normal serums.

DR. L. B. WILSON, Rochester, Minn.: The grouping into toxic non-exophthalmic and into toxic exophthalmic cases is purely clinical. It is found that the exophthalmic toxic cases invariably show hypertrophy and hyperplasia, while the non-exophthalmic cases do not show these symptoms.

DR. A. H. SANFORD, Rochester, Minn.: There was a fall in blood-pressure even after both vagi were cut. Furthermore, Gley and Cleret in their series stated that there is inhibition of vagus action after the injection of the serums and goiter extracts. We could not demonstrate this. By the use of the intestinal plethysmograph we were able to demonstrate that a marked splanchnic dilatation occurred coincident with this fall in blood-pressure. We injected these serums as we obtained them from patients without noting their histories. Sometimes we obtained marked fall in blood-pressure and at other times nothing at all. We did not know beforehand what our results would be. In reviewing the case histories it was found that the serums in which we had a marked fall in blood-pressure were from those patients who had acute intoxication from exophthalmic goiter and that those from which no effect was obtained were from the chronic cases. It appeared to us quite significant as to the amount of depressor substance present in the blood of these patients at that particular time.

Rural Sanitation.—In many states the large cities are the only points which have any real sanitary protection. Our rural population of 49,000,000 people, including the 30,000,000 who live on farms, with the exception of those fortunate rural dwellers in one or two exceptional states which regard the health of the country people as also important, receive very little state health protection and maintain no local protective system of their own against contaminated local milk, meat, shell-fish, fish or vegetables. Moreover, only a small number of these rural inhabitants are safeguarded, by competent inspection, against polluted water-supply or sewage-disposal methods dangerous to health and life.—Carl L. Alsberg, *Am. Jour. Pub. Health*.

FIELD-WORK IN TUBERCULOSIS*

MARY E. LAPHAM, M.D.

HIGHLANDS, N. C.

It is small comfort to a man dying of tuberculosis to be told that, if the disease had been discovered in time and properly treated, his life might have been saved. It is more than doubtful if the frequency of this failure to discover the disease in time is adequately appreciated outside a sanatorium. There we learn with appalling emphasis the lesson taught by experience that death in tuberculosis is not entirely due to an infection by tubercle bacilli, but equally, and possibly more than equally, due to failure to detect the beginnings of pathologic processes. The fault is not altogether in the inability or the neglect of physicians, because only too often the patient is all unconscious of the danger and does not seek help. It is to emphasize the importance of educating the people to a sense of this danger that this paper is written.

These cases of delayed diagnosis in which the patients do not seek a doctor until it is too late are due to lack of manifestations, so that the patient is not aroused in any way to the fact that there is any trouble in his lungs. He does not cough, has no fever, feels well, looks well and does his day's work with ease. We have at the Highlands Camp Sanatorium under our care patients who, up to the very time that they finally felt ill enough to call in a doctor, were feeling as well and working as well as they ever had. How shall we teach our people to realize that apparently good health is no guarantee against tuberculosis and that good working capacity is perfectly compatible with bad lungs? How can we make them appreciate the dangers of tuberculosis among apparently well people and the desirability of guarding against its spread and development by periodic examinations? How can we explain the long years of preparation and teach them that the victims discovered to-day have been years in the making, and that, therefore, the time to attack tuberculosis is not at the end, when it is perforce discovered, but during those long years of slow development when the crops were growing?

The recent work done in the South against the hookworm with such splendid results suggests the desirability of employing just such tactics against tuberculosis. In the South our people did not know why they could not work or that they had hookworms or that they could be cured. They had no hope of help and rarely sought it. By going out into the field and searching out these cases and teaching the people and the physicians on the spot an incredible amount of working capacity has been acquired for the South and this saving will steadily increase while the expenses of the campaign will lessen and cease. In the ordinary course of events how long would it have taken for this knowledge to filter down through to physicians and patients?

If another Rockefeller would establish a fund for field-work in tuberculosis so that we could plan the campaign along the same lines as that against the hookworm and put a tuberculosis expert in charge of each district, what should we hope to accomplish? First, we should want to educate the people so that they would understand the problems connected with the prevention of the disease; and, second, we should make a thorough canvass in order to learn how many cases of unsuspected latent tuberculosis were insidiously preparing to become manifest.

* Read in the Section on Preventive Medicine and Public Health of the American Medical Association, at the Sixty-Fourth Annual Session held at Minneapolis, June, 1913.

The district physicians will make a thorough search for suspicious cases and bring them for examination on the appointed day by the expert. This will teach the profession the most reliable and newest methods of examination and enable them to learn to detect the presence of a tuberculous process long before there are any manifestations. The expert will insist on the examination of apparently well children because he knows that it is precisely among these seemingly well children that concealed tuberculosis is found, and that if these cases are put under proper treatment many an adult will be saved. The expert will teach his physicians the necessity for distinguishing between an infection by tubercle bacilli and tuberculosis, because whereas over 90 per cent. of our children are infected by tubercle bacilli by the time they reach maturity, only a very small percentage manifest the disease and it is therefore evident that not all these infections develop into a tuberculous process. We can detect the unsuspected presence of tubercle bacilli by a tuberculin test, but this may be simply a biologic and not a pathologic reaction. The ability to react to tuberculin increases with each year of life from birth to maturity, so that we as constantly find an increase in the number of infections by tubercle bacilli, but this does not mean that the manifest cases of tuberculosis increase in the same proportion.

From birth to maturity there is an ever-increasing disproportion between the percentage of tuberculin reactions, the manifested cases of tuberculosis and the death-rate. We cannot tell by a tuberculin test whether or not a new-born babe has a tuberculosis infection because the ability to respond to tuberculin is still undeveloped.

With every year of life we find an increase in the number of children responding to tuberculin and we have argued from this increase a corresponding increase in the number of infections. This inference may be wrong in some cases because instead of a new infection we may have simply a newly developed ability to respond. Every day the study of the newly born and of the prematurely born is showing with ever-increasing frequency the presence of acid-fast bacilli or their prebacillary forms known as Much's granules in the blood and tissues when there was absolutely nothing to suggest their presence. The latency of the infections is so great that we are beginning to believe that they are as universal and as harmless as those by colon bacilli.

It is possible that the story of tuberculosis begins at birth and that the growing organism has to decide at the very start the question of its relations with this infection. If it is able to assert a phylogenetic ability to retain acid-fast infections in a state of harmless residence, as we all do colon bacilli, then tubercle bacilli are not dangerous to that individual. If there is a flaw in this mechanism, if the adaptation is not complete, the manifestation of this flaw will depend on an infinitude of factors. If the failure is conspicuous, the newly born will manifest it and die. One per cent. of all our children manifest this failure during the first year of life and die, not because they have been exposed, but because they have failed to adapt themselves. These failures inevitably die because they have never established the ability to live and the proportion between the number of manifest cases and the deaths is 100 per cent. The proportion of the death-rate to the total number of children is 1 per cent., and the percentage of infections is the same. Even in the second year of life discrepancy appears between the percentage of tuberculin reactions, the manifest cases and the death-rate, because no distinction is made between the tuberculin reactions and the manifest cases.

We argue from the increased number of reactions an increase in the number of infections and because the death-rate does not correspondingly increase it falls in proportion. If we should carefully distinguish between these tuberculin reactions and the evident manifest cases we might find that the death-rate was the same for these as in the first year of life and that the discrepancy was solely due to an increased ability to respond to tuberculin. That this is true is supported by the fact that acid-fast bacilli and Much's granules are found in infants that do not respond to tuberculin until later. With each year this discrepancy becomes greater until at maturity we find over 90 per cent. of tuberculin reactions and a death-rate of only 2 per cent. These figures will not reveal their true significance until a careful estimate is made of the distinction between manifest cases and biologic reactions. It may be that we shall find the actual death-rate higher at maturity than at any other period of childhood because of the ever-developing slowly manifested inability to live in safety with tubercle bacilli. It may be that the child that dies at 14 does not die because of a new infection but because the disparity between the danger from the infection and the ability to resist it increases. If this proves true, the tuberculosis work in the future must be directed toward detecting the children threatened with this failure and endeavoring to correct it. Prophylaxis will consist in preventing the development of this flaw and never allowing it to go on to the tuberculosis of adults.

To be able to distinguish between pathologic and biologic reactions our experts will need a well-equipped laboratory and Roentgen apparatus. This will mean great diagnostic skill and a great deal of work with consequent need of assistance. This will give our recent graduates a splendid opportunity to learn the most difficult task of distinguishing between a simple presence of tubercle bacilli and their assumption of pathogenic qualities. The keeping of these records and their comparison with the subsequent histories will teach us more concerning the relationship of these childhood infections with the tuberculosis of adults than we could possibly learn in any other way. We know that primary infection of the adult is impossible, but we do not know the history of these years between the primary infection and its final manifestation. By periodic examinations of our patients we can reveal these connections and study the effect of saving the child on the future death-rate from tuberculosis. This great effort cannot be forced on the people by the profession and they must therefore be taught to demand it, and to see that their future safety depends on it. The expense seems prohibitive, but we may come to realize that the money has to be spent anyway, and that it is only a question of when and how—whether we shall build and maintain sanatoriums for manifest cases and be utterly unequal to the problem of caring for them all, or whether preventoriums will not save years of wasted educational and living expenses lost at the very time the expense was to be justified.

The campaign of education will be greatly furthered by the stimulus to the comprehension of the people from the practical work carried on among them. Explanatory of these results, they will understand pathologic sections of lungs made during all stages of the disease, from the tissues swarming with the bacilli and yet not in the least injured, to the rotten lungs reeking with corruption.

Lectures illustrating the slow development of these processes in apparently well people and their compatibility with good health and working capacity will stimulate a desire for protection against this danger so subtle that

it cannot possibly be appreciated. Above all else, the length of time should be emphasized, because there is a strong belief that an infection by tubercle bacilli resembles other infections, such as typhoid, and that recovery is, therefore, comparable to that from typhoid. If the people could be made to understand the nature of recovery in tuberculosis there would be fewer deaths from idiotic impatience. In our schools the teachers and pupils can be easily taught the whole story of tuberculosis as soon as the knowledge is sufficiently diffused to furnish us with teachers. How else can this be done as quickly as by field-work done just as it was against the hookworm?

We guard our schoolchildren from infection in every way we can, but there are two sources of infection that do not sufficiently attract our attention. One source is in the unsuspected carriers of tubercle bacilli that during a lifetime cough a little, or have "winter colds" or "chronic bronchitis," but whose general health and appearance do not suggest tuberculosis. One man complacently told me when he brought his wife that he had lost two wives from tuberculosis and probably he would lose this one, which he did. This man was the unsuspected carrier of tubercle bacilli. A few months ago the mother of one of my patients died at 68 and tubercle bacilli were found just shortly before her death, but not until after two of her children had died from tuberculosis. We distinguish between the danger from cases with tubercle bacilli in the sputum and those without. In many cases the prebacillary forms of Much's granules are present when the adult bacilli are not and their virulence is equally great, if not greater, so that the distinction between an open and a closed case must be changed to sputum containing no Much's granules and not capable of causing tuberculosis in a guinea-pig by inoculation.

Prophylaxis in the future must consist in preventing the development of a universal infection common to all mankind, which can in no way be avoided and only restrained. The direct transmission from case to case is entirely inadequate to account for this universal infection and we must seek some other explanation. We know that throughout the grasses and among trees and with the lower cold-blooded animals such as snakes, fish, turtles and lizards, and with the higher animals, such as birds, poultry, cats, dogs, sheep, pigs and cows, each class has its own peculiar kind of acid-fast bacilli capable of producing tubercles and manifesting their genetic relationship to each other by tuberculin reactions. That these tubercle-producing acid-fast bacilli are universally present throughout Nature and that they all belong to the same family is unquestioned, and that changes in type can be induced by changes in the host is indisputably proved.

Starting with these undoubted facts, it may be eventually proved that the universal infection of the human race by acid-fast bacilli is comparable with its universal infection by colon bacilli and that the danger essentially consists in the assumption of pathogenic qualities. As a rule the human race has acquired the ability to restrain this assumption, but occasionally in a very small percentage of cases this ability cannot be established and tuberculosis results. The detection of this change at the earliest possible moment and its conversion into safety is the fundamental basis of our future efforts against the spread of tuberculosis. Periodic examinations of our schoolchildren would do much to reduce the tuberculosis of adults.

Highlands Camp Sanatorium.

ABSTRACT OF DISCUSSION

DR. ROSINA WISTEIN, Chicago: I do not agree with Dr. Lapham that the education of the public must come from the laity, and not from the medical profession. The laity are ignorant of the dangers that are confronting them; and I believe that it is the duty of the medical profession to teach others, and only in proportion as we are interested can we hope to interest others. I find the greatest indifference and carelessness not on the part of the laymen, but on the part of the medical practitioner.

SEVERANCE BURRAGE, PH.D., Indianapolis: I have met all kinds of people in this antituberculosis work, and I wish to corroborate what has been said in regard to the inactivity and careless attitude of physicians in general. In Indiana we are organizing the whole state in an antituberculosis campaign. We are organizing a local society in each county to carry on a campaign of education and disease prevention in that county. We find that the women are the best workers in most of these organizations, and that the physicians are really the hardest ones to induce to do active work. Some of them seem to feel that if they take an active part in a campaign for a local hospital—tuberculosis hospital—it will give the appearance of local advertisements or create certain jealousies among the physicians that will be undesirable, and will be looked on, possibly, as unethical or outside the pale of the physician's etiquette. For that reason we have found it difficult in many cases to enlist the active interest of physicians in the antituberculosis work. Some of the best workers are schoolteachers. They have a great deal of teaching to do, anyway; and just as schoolteachers are oftentimes enlisted in the Sunday-school work and in other kinds of teaching, they are also enlisted in this work. It does not seem right to me; I think that in a community in which facts in regard to disease—causes of disease and disease prevention—are concerned that the physicians should be the leaders in this educational campaign. They are the ones to whom the public should look for accurate information on these subjects; and yet, as I have stated, we find in the majority of cases that active workers among the physicians are hard to find.

DR. C. C. BROWNING, Los Angeles: I believe that the plan of fieldwork which the doctor has mapped out is worthy of more than passing notice. On account of my connection with the state tuberculosis commission of California, I have had occasion within the last two years to consider this subject somewhat carefully. We have adopted in our report a scheme founded on the campaign against the hookworm in the South. Our recommendation is the division of the state into sanitary districts, each of which is to be presided over by a dispensary physician, aided by a corps of nurses, for the express purpose of the detection of tuberculous patients. I believe that this plan promises the greatest amount of good in the future for the amount of money expended.

DR. W. A. EVANS, Chicago: There is, I think, a general consensus of opinion that tuberculosis in the cities is on the decline. Among the older cities—cities along the Atlantic seaboard—and the cities of continental Europe, the decline is about 50 per cent. in round figures. In our section of the country—in the Middle West—the decline is about 25 per cent. Our campaigns were not inaugurated so early as were the campaigns to the east of us, and probably they have not been so effectively conducted. It seems to me that the teaching of these experiences is quite conclusive that in communities that have undertaken this work thoroughly results follow, and that we are justified in concluding that tuberculosis will respond to efforts at its control. There are probably some states in which there is a decline in tuberculosis outside of the cities; but in the main, I think, the statement is true that there is—aside from the increasing amount of discovered tuberculosis—an actually increasing amount of tuberculosis in the rural districts.

The work for the control of tuberculosis in the rural districts is not so well developed as in the cities. I know of the plans that are being pursued in various cities in the vicinity of Chicago. It has seemed to me that the best of those plans was that in operation in Wisconsin. In Illinois the women's clubs are conducting a tuberculosis survey—a survey the

results of which will be inaccurate, but which, nevertheless, will serve to concentrate the attention of the women on the tuberculosis problem in their respective communities. We are employing visiting nurses, under the auspices of the state tuberculosis society. A nurse investigates conditions in one county, and the idea is that when she leaves and moves over to another county there will be left behind in that county one or more local county nurses maintained by philanthropy. So far as I know, this idea has been better worked out in Wisconsin than in any other state in the Union, certainly better than in any other state in our part of the country.

DR. GERTRUDE FELKER, Dayton, Ohio: Four years ago the House of Delegates gave to the women physicians of the American Medical Association the task of forwarding the education of the people. This committee is now known as the Committee for Public Health Education Among Women. The effort has been to find some physician in every state who would be interested in education; and through that physician as state officer to find some one in each county who would talk public health education and have in each county society a public health education committee appointed. We have asked that a woman be put on that committee in counties in which creditable work is being done by women physicians. After four years' work almost five hundred counties have appointed committees.

DR. MARY E. LAPHAM, Highland, N. C.: The purpose of my paper was to urge the appointing of an expert from the medical profession to teach the people the earliest suggestions of tuberculosis. The education of the people should come from the profession. The earliest recognition of tuberculosis must come from the people in order that the profession may have the earliest opportunity to confirm it. It is because the people do not understand the situation that the general practitioner does not have the opportunity to make a sufficiently early diagnosis.

THE HISTOPATHOLOGIC LOAN COLLECTION AND RELIEVING THE CURRICULUM *

H. S. STEENSLAND, M.D.

AND

H. G. WEISKOTTEN, M.D.
SYRACUSE, N. Y.

It was pointed out by Bardeen¹ that it is essential to have *good* histologic preparations if the student of normal histology is to see the details of structure as they exist in the tissues. Otherwise he is likely to make mistakes and to draw wrong conclusions from what he sees, and perhaps eventually to fabricate from grosser features of a histologic preparation the finer details that it does not present. This is just as true in the case of pathologic histology. A common method of handling routine teaching material is to give the student sections which he is expected to stain and mount for himself. Another practice is to distribute stained sections for the student to mount. Frequently it results that the preparations are not satisfactory and that much of the time and energy of the student is wasted in puzzling over them. For this reason it would appear to be wise to make extensive use of loan preparations.

The amount of time that can be given each year by members of the teaching staff to the preparation of histologic teaching material is limited. If this time is given to the making of a limited number of permanent loan preparations of high quality covering a limited field, there will accumulate eventually a loan collection of high

grade covering a large field. Then there will be histopathologic loan collections comparable to the Minot embryologic collection at Harvard and the Mall embryologic collection at Johns Hopkins. Also the student will have for study carefully selected material, well fixed, cut into thin sections, and carefully stained and mounted. The time that the student formerly gave to the routine of staining and mounting sections he can give to actual study of the preparations.

It may be objected that the student should have the technical experience involved in staining and mounting; but with the gain in time he can be given a short but more varied training in technic in place of the oft-repeated routine. The ownership by the student of a collection of sections is a doubtful advantage as compared with the length of time that he spends in staining and mounting. The sections probably are rarely referred to after the course is completed. The mere collecting of the specimens is apt to be uppermost in the student's mind in place of the more important careful study of the material. It is not necessary to have one preparation of each kind for each student. One preparation for each pair or larger group of students is sufficient. The selection of material for permanent preparations for teaching purposes and the preparation of this material up to the point of orienting the blocks of tissue should have expended on it the greatest skill that the staff affords.

A modern course in pathology requires the use of a larger amount of experimental histologic material than was used ten or fifteen years ago. It is necessary to use to a larger extent tissues affected by means of known injurious agents under accurately known conditions. The amount of time required for the preparation of such material is very large, and the amount of material yielded by each experiment is frequently small. It is therefore obviously impossible to duplicate such material as frequently as would be required for giving each student a specimen as his personal property.

At Syracuse a loan collection as above set forth was begun by one of us (Steenland) ten years ago. A large gift from Dr. Mallory of imbedded specimens selected from his Harvard teaching material formed the nucleus of the collection and constituted a model and ideal of quality toward which to strive. As a result of the use of special care in the technic to prevent fading of the stains the oldest material is to-day apparently as good as when first prepared. Additions can be made to the collection as advances in the knowledge of diseased tissues require, and antiquated material can be replaced by new material. Each year special selection from the collection may be made for use by the students according to the trend in special directions which is given to the course in that particular year.

Pearce² and Gay³ have pointed out the great advantage of correlating the work in experimental pathology and pathologic physiology with the work in pathologic histology. With a given number of scheduled hours the possibility of gain in time resulting from the use of a loan collection has an important bearing here. This gain in time for the student and the teaching staff will permit of the introduction of more work in experimental pathology and pathologic physiology into the usual course in pathology. The amount of such work that can be done by students in any particular department will depend on

* From the Department of Pathology, Syracuse University, College of Medicine.

1. Bardeen, C. R.: Report of the Subcommittee on Anatomy to the Council on Medical Education of the American Medical Association, April, 1909, *Anat. Rec.*, 1909, iii, 415.

2. Pearce, Richard M.: The Teaching of Experimental Pathology and Pathological Physiology to Large Classes, *Bull. Johns Hopkins Hosp.*, 1911, xxii, 404.

3. Gay, Frederick P.: A Method of Correlated Teaching of Pathology and Bacteriology in the Second Year of Medical Instruction, *Bull. Johns Hopkins Hosp.*, 1912, xxiii, 170.