I saw both whales' bones and drift-wood *at least thirty feet*, and I think as much as forty feet, above high-water mark. Geologists can form their own deductions from this; but I may mention that the seal-fishers all believe the land to be rising, or, as they generally express it, "the sea is going back."

Whales used to be formerly very abundant around the shores and bays of Spitzbergen, but of late years the true whale (*Mysticetus*) is unknown within many miles of Spitzbergen. Macculloch and other commercial writers attribute this to the persecution they have undergone; but the seal- and walrus-fishers all say that "the seas have got too shallow for them." This may also arise partly from the enormous quantities of mud and débris brought down by the numerous rivulets running from the steep slopes of the mountains. Some of these rivulets have immense deltas of semifluid mud at their mouths.

The Thousand Islands are an immense cluster of low rocky islands lying off the south-east corner of Spitzbergen, and consist entirely (at least, as far as I could observe) of a coarse granular trap. They are mostly very low, and with very little soil on their surfaces. I observed the skeletons of whales upon some of them at such a height that it was quite impossible they could have been washed there with the rocks at their existing level; and this seems to me to afford conclusive evidence that Spitzbergen and the adjacent islands are emerging from the deep at a rate even more rapid than that at which the coasts of some parts of Norway have been clearly demonstrated by distinguished geological writers to be rising.

2. On the FORMATION of GYPSUMS and DOLOMITES.

By T. STERRY HUNT, Esq., of the Geological Survey of Canada.

(Communicated by Professor Ramsay, F.R.S., F.G.S.)

THE deposits of gypsum in nature may be divided into two classes. One of these comprises the gypsums formed by the alteration of beds of limestone. The vapours of solfataras, the sulphuric acid produced by the slow oxidation of moist sulphuretted hydrogen gas (as shown by Dumas), and springs containing free sulphuric acid are the agents which have changed and are still changing carbonate into sulphate of lime. Such springs, containing from $\frac{2}{1000}$ to $\frac{4}{1000}$ ths of free sulphuric acid (evolved probably from the reaction between sulphate of lime and siliceous matters under the influence of heat at considerable depths), are frequent in Western Canada; and their effects in giving rise to masses of gypsum in the quaternary clays of that region I have long since pointed out (Rep. Geol. Survey of Canada, 1848, p. 155).

The gypsums of the second class, which are the more frequent, are met with interstratified with marls, dolomites, and rock-salt, with which they are evidently contemporaneous. Reserving for

another occasion the discussion of the theories which have been proposed to account for the origin and association of these substances, I propose to mention some recent observations of mine which serve to throw light upon the question.

When a solution of bicarbonate of lime in carbonic-acid water is mingled with one of sulphate of magnesia, there are formed sulphate of lime and bicarbonate of magnesia; so that in presence of an excess of bicarbonate of lime there is obtained a saturated solution of gypsum, from which this salt may be thrown down by the addition of alcohol.

In the same way I have found that a solution of sulphate of soda is decomposed by bicarbonate of lime, whose solubility in water is much augmented by the presence of the alkaline sulphate. From such a solution the addition of alcohol precipitates gypsum, leaving bicarbonate of soda in solution.

When a mixed solution of sulphate of magnesia and bicarbonate of lime is evaporated at a gentle heat (from 80° to 120° F.), the gypsum, being much less soluble than the bicarbonate of magnesia, is first deposited in a crystalline form, and at a later period the magnesia separates as a hydrous carbonate. The presence of sea-salt does not prevent this reaction, nor the precipitation of gypsum by evaporation; so that by continued additions of bicarbonate of lime to a basin of sea-water, its sulphate of magnesia might at length, under favourable conditions, be entirely converted into gypsum.

The hydrated carbonate of magnesia deposited from solutions of the bicarbonate is very soluble in dilute acids; but when heated to 300°–400° F. under pressure to prevent the escape of carbonic acid, it is converted into a crystalline, anhydrous, and difficultly soluble carbonate, already obtained by De Senarmont, and apparently identical with magnesite. When the hydrous carbonate is thus heated in presence of carbonate of lime, the two combine to form a double carbonate, which, by taking advantage of its sparing solubility in cold dilute acetic acid, may be separated from any admixture of free carbonate of lime. This double carbonate of lime and magnesia is dolomite. As prepared under various conditions, I found it to contain from 35 to 45 per cent. of carbonate of lime, the excess of magnesia being due apparently to a variable admixture of magnesite. Its formation by the direct union of the carbonates of lime and magnesia takes place alike in the presence of solutions of earthy chlorides and alkaline carbonates. In Von Morlot's experiment, however, which consists in heating a mixture of two equivalents of carbonate of lime and one of sulphate of magnesia in sealed tubes to 200° C., the sulphate of lime which is formed seems to prevent the union of the two carbonates, and the product I have found to be chiefly magnesite mixed with carbonate of lime. In Marignac's experiment, where chloride of magnesium is substituted for the sulphate, the newly formed carbonate of magnesia unites with the carbonate of lime, and a portion of dolomite is formed.

But the solutions of bicarbonate of magnesia which give rise to deposits of hydrous carbonate of magnesia, that, according as they

are pure or mingled with carbonate of lime, are subsequently converted into magnesite or magnesian limestone, may also be formed by another reaction, which is independent of the formation of sulphate of lime. I have already alluded to it, in my paper "On some points in Chemical Geology," read before this Society on the 5th of January last. Waters holding in solution bicarbonate of soda (a constant product of the decomposition of feldspathic minerals), when mingled with sea-water, decompose the lime-salts first, the carbonate of lime separating in a nearly pure state; so that in lakes and limited sea-basins there must result from this reaction waters that, like the bitters from which all the lime has been removed by evaporation as sulphate (another source of gypsum), contain only salts of soda and magnesia. The further action of solutions of bicarbonate of soda upon such magnesian waters must give rise to bicarbonate of magnesia, and, aided by evaporation, to precipitates of magnesian carbonate—which, mingled with the carbonate of lime generally accompanying the bicarbonate of soda in alkaline waters, would afford the material for those great beds of magnesian limestone which are independent of gypsum. (See Proc. Royal Soc. for May 18, 1858; Phil. Mag. vol. xvi. p. 379.)

The points to which I wish to call attention in the present note are, first, the formation of sulphate of lime and bicarbonate of magnesia by the action of bicarbonate of lime upon a solution of sulphate of magnesia, and their successive deposition in the forms of gypsum and hydrous carbonate of magnesia during the process of evaporation; and secondly, the direct union, under certain conditions, of this carbonate of magnesia with carbonate of lime to form a double carbonate, which is dolomite.

The application of these reactions to explain the formation of dolomites and a great portion of the gypsums, both in marine and freshwater deposits, will readily be made. The details of my experiments will be found in part in the Report of the Geological Survey of Canada for 1857*, and completed in the forthcoming Report for 1858.

-
3. *On the TERTIARY DEPOSITS, associated with TRAP-ROCK, in the EAST INDIES.* By the REV. STEPHEN HISLOP. *With DESCRIPTIONS of the FOSSIL SHELLS,* by the REV. S. HISLOP; *and of the FOSSIL INSECTS,* by ANDREW MURRAY, Esq., F.R.S.E.; *and a Note on the FOSSIL CYPRIDÆ,* by T. RUPERT JONES, Esq., F.G.S.

[Communicated by the President.]

(PLATES V.—X.)

In introducing the following descriptions of some fossils from Peninsular India, I shall confine my remarks to the rocks, volcanic

* See also Am. Journ. Science, 2nd ser. vol. xxvi. p. 109.

TERTIARY SHELLS: NAGPUR.

TERTIARY SHELLS: NÁCPUR.

26

28

27

TERTIARY SHELLS : NÁCPUR.

TERTIARY SHELLS. RAJAMANDRI.

TERTIARY SHELLS RÁJÁMANDRI

TERTIARY SHELLS, INSECTS, & CYPRIDES RÁJÁMANDRI, NARBADDÁ, & NÁGPUR.