

Theory and simulation of the ultrafast double-bond isomerization of biological chromophores

Samer Gozem[†], Hoi Ling Luk^{‡,††}, Igor Schapiro[¶] and Massimo Olivucci^{‡,#}

[†] Department of Chemistry, Georgia State University, 50 Decatur St. SE, 262 Natural Science Center Atlanta, Georgia 30302, United States; [‡] Chemistry Department, Bowling Green State University, Overman Hall, Bowling Green, Ohio, 43403, United States; [¶] Fritz Haber Center for Molecular Dynamics, Institute of Chemistry, The Hebrew University of Jerusalem, Jerusalem 91904, Israel; [#] Dipartimento di Biotecnologie, Chimica e Farmacia, Università di Siena, via A. Moro 2, 53100 Siena, Italy.

KEYWORDS: Photoresponsive Proteins, Rhodopsins, Xanthopins, Phytochromes, Photoreceptors, Ultrafast Photoisomerization.

ABSTRACT: Ultrafast processes in light-absorbing proteins have been implicated in the primary step in the light-to-energy conversion and the initialization of photoresponsive biological functions. Theory and computations have played an instrumental role in understanding the molecular mechanism of such processes, as they provide a molecular-level insight of structural and electronic changes at ultrafast time scales that often are very difficult or impossible to obtain from experiments alone. Among theoretical strategies, the application of hybrid quantum mechanics and molecular mechanics (QM/MM) models is an important approach that has reached an evident degree of maturity, resulting in several important contributions to the field. This review presents an overview of state-of-the-art computational studies on sub-nanosecond events in rhodopsins, photoactive yellow proteins, phytochromes, and some other photoresponsive proteins where photoinduced double-bond isomerization occurs. The review also discusses current limitations that need to be solved in future developments.

CONTENT

1. Introduction
2. A case study: the primary event in vision
3. Theory of double-bond isomerization
4. Rhodopsins
5. Photoactive yellow proteins
6. Phytochromes
7. Other photoresponsive proteins
8. Conclusions and perspectives

1. INTRODUCTION

Nature has developed light-responsive proteins capable of exploiting the energy of a photon to accomplish specific electronic and geometrical changes at the molecular level. Through such changes, the photon energy is stored and then employed to initiate a series of conformational transformations

defining the protein photocycle. Such photocycles are implicated in a variety of biological functions such as daylight and nocturnal visual perception in vertebrates and invertebrates¹, light-oriented growth in plants², phototaxis³, light sensing⁴ and ion pumping⁵ in eubacteria and Archaea, and ion gating in unicellular algae⁶, to cite a few.




Figure 1.1. Light-responsive proteins. A. Families of light responsive proteins in plants and fungi. Ultraviolet-B receptor (UV-B), light-oxygen-

voltage-sensing domains (LOV), cryptochromes (CRY), rhodopsins and phytochromes (PHY). The structure of the prosthetic groups is also given. Rhodopsins and phytochromes exploit a double-bond photoisomerization to initiate the corresponding photocycles. Adapted with permission from ref. ⁷. Copyright 2012 John Wiley & Sons, Inc. B. Structure of bovine rhodopsin. C. Details of the tight chromophore cavity of bovine rhodopsin and of the structure of its 11-*cis* retinal chromophore. The chromophore isomerizing double bond (C11=C12) is marked with a curly arrow. Adapted with permissions from ref. ⁸. Copyright 2001 American Physiological Society.

In their simplest form, light-responsive proteins consist of a cytoplasmic or membrane protein hosting a prosthetic group, the chromophore (see Fig. 1.1A), responsible for light absorption, photon energy storage and photocycle initiation. These relatively basic systems are generally classified in families which include rhodopsins, phytochromes, xanthopsins, cryptochromes, phototropins, and BLUF proteins.⁹ The latter three families have a flavin chromophore that initiates various intermolecular photophysical and/or photochemical processes (e.g. LOV and CRY in Fig. 1.1A belong to these three families). In contrast, the former three families feature diverse chromophores which, upon illumination, undergo the same "elementary" intramolecular chemical process: a sub-picosecond isomerization of one of the double bonds of their conjugated frameworks (e.g. RHODOPSINS and PHY in Fig. 1.1A). In the present review we mainly focus on these "photoisomerizing" protein families.

The rhodopsin, phytochrome and xanthopsins families have served as a virtually endless source of interest, as well as inspiration, for scientists working in different fields such as photochemistry, photobiology and biophysics. As a consequence, there have been numerous theoretical studies aimed at understanding their spectroscopy, color tuning and photochemistry, not to mention the light energy transduction mechanism characterizing their functions. While here we will be concerned exclusively with double-bond isomerization, we need to warn the reader that the present review is not exhaustive even within this ostensibly narrow topic. Rather, this review reflects the interest of the authors, whose aims are a complete molecular-level understanding of this process and the design of efficient computational tools/strategies providing access to it.

In order to illustrate the impact that the theoretical investigation of double-bond isomerization has on our comprehension of light-responsive proteins, we start with a section (Section 2) reviewing the results of complementary spectroscopic and computational studies on the photoreceptor mediating scotopic (i.e. dim light) vision in vertebrates: rod rhodopsin. We start with this example because it is one of the most studied processes in photobiology, both experimentally and theoretically, and therefore serves as a prototypical system. The review will then continue by looking at the current understanding of specific protein families. To do so, it is necessary to start with a section (Section 3) revising our present understanding of double-bond isomerization in conjugated organic molecules and the computational methods available to construct and study light responsive protein models. Three more specific sections will then follow focusing on rhodopsins (Section 4), photoactive yellow proteins (Section 5) and phytochromes (Section 6). The involvement of double-bond isomerization in other light-responsive proteins will be finally discussed in an additional

section (Section 7). Most of these sections are organized in a way to highlight studies that investigate the structure of CIs first, then the mapping of excited and ground-state reaction paths, and trajectory studies last. The review is closed by a conclusion section also examining the perspectives of this research field (Section 8). We have also included an appendix (Appendix I) that reviews some of the computational methodologies available for studying the thermal and light-induced isomerization of biological chromophores. To get a better appreciation of the work reviewed here, we recommend the reader to read Appendix I before proceeding to Sections 4-7. This will not only serve to get familiar with terms such as "CASPT2", "CASSCF", and "QM/MM", but also to equip the reader with knowledge of the advantages and drawbacks of different approaches thus allowing them to better understand the results and discussions.

2. A CASE STUDY: THE PRIMARY EVENT IN VISION

2.1 Spectroscopic studies: The photoisomerization of rod rhodopsins is vibrationally coherent.

Dim-light vision is initiated by the light-induced isomerization of one of the bonds of the 11-*cis*-retinal protonated Schiff base (rPSB11). This is the chromophore of rhodopsin: one of the light-responsive proteins found in the retina of vertebrates. The crystallographic structure of bovine rhodopsin is shown in

Fig. 1.1B and features seven transmembrane α -helices forming a tight cavity where a lysine residue is covalently linked to the chromophore through a Schiff base group (see Fig. 1.1C). Upon illumination, rPSB11 isomerizes inside the tight protein cavity about the C11=C12 double bond producing the corresponding all-*trans* isomer (rPSBAT) on a sub-200 fs timescale.¹⁰ This event corresponds to the first step of the rhodopsin photocycle (see Fig. 2.1) and leads to the formation of the photorhodopsin intermediate that, after vibrational relaxation, yields bathorhodopsin. The photon energy stored in the bathorhodopsin intermediate is then used to drive the photocycle and, ultimately, the protein function through the production of the meta II activated state.




Figure 2.1. The rhodopsin photocycle involves several intermediates defined by the characteristic wavelength of their light absorption maximum (λ_{max}). The first spectroscopically detectable intermediate after illumination is photorhodopsin, which is interpreted as a vibrationally hot

form of the first isolable intermediate bathorhodopsin. Subsequent intermediates emerge through further thermal relaxation. Bathorhodopsin may be regarded as the intermediate storing the photon energy. This energy is then used for the production of meta II: a key event within the photocycle constituting the signal-transducing state of rhodopsin (i.e. its biological function). Since the C11=C12 isomerization is fully accomplished at the level of the photorhodopsin/bathorhodopsin intermediates, the present contribution reviews the mechanistic studies focusing on the framed part of the photocycle. Adapted with permission from ref. ¹¹. Copyright 2014 John Wiley & Sons, Inc.




Figure 2.2. Mechanism of the coherent isomerization of bovine rhodopsin. A. Conical intersection mechanism for the ultrafast isomerization of bovine rhodopsin occurring via population transfer from the excited-state to the ground-state potential energy surfaces (adapted with permission from ref. ¹². Copyright 2015 Nature Publishing Group). B. Avoided crossing mechanism proposed for the same ultrafast isomerization (adapted with permission from ref. ¹⁰. Copyright 1991 The American Association for the Advancement of Science and ¹³ Copyright 1994 The American Association for the Advancement of Science). S_0 and S_1 are labels indicating that the initial light absorption promotes the system from the ground singlet state to the first singlet excited state, respectively.

A recent spectroscopic study by Miller and collaborators¹⁴ has indicated that the transition from the excited state of rPSB11 to the ground state of rPSBAT is not only ultrafast, but vibrationally coherent. In a contribution¹² on the significance of such a finding, Mathies retraces the progress made through the spectroscopic investigation of double-bond isomerization in bovine rhodopsin (Rh), highlighting Miller's work as the conclusion of a research line started in the early '90s. In their con-

tributions, both Miller and Mathies provide a mechanistic interpretation of the observed spectroscopic signals in terms of first singlet excited state (S_1) decay at a *conical intersection* (CI) progression along S_1 and ground state (S_0) *reaction coordinates (and/or paths) intercepting the CI*, and *trajectories* that describe the evolution of excited molecules with time and their transfer from the S_1 to the S_0 potential energy surfaces in the CI region. Such terminology is rooted in theoretical research efforts that have advanced the mechanistic understanding and computer simulation of organic photochemical reactions during the last few decades. Such advancements started to consolidate and became routinely available for computations in the '90s, therefore roughly at the same time of the early spectroscopic measurements on Rh.

CIs, reaction coordinates and trajectories constitute key elements for the mechanistic description of photoinduced double-bond isomerizations, as well as for any other photochemical reactions not involving a change of spin symmetry (e.g. singlet photochemical reactions).¹⁵ Indeed, in the Miller¹⁴ and Mathies¹² contributions, the observed coherences in the spectroscopic signals are effectively interpreted as collective nuclear (vibrational) motions of the molecular population towards or away from specific CI structures. Such motion not only ensures that the population moves, as a whole, out of the vertical excitation region of the excited state potential energy surface (see Fig. 2.2A), but also that, in each molecule, the distinct vibrational modes describing such progression maintain a certain phase relationship. More specifically, it is proposed that after light absorption, the population of protein-embedded rPSB11 chromophores initiate a vibrationally phased motion along critical modes such as the bond-length-alternation (BLA) stretching, the hydrogen-out-of-plane (HOOP) wagging, and the torsional deformation of the reactive double bond. These modes would drive the population towards a CI between the S_1 and S_0 potential energy surfaces (in green and red respectively in Fig. 2.2A) thus transferring it to S_0 . Such a mechanistic description is consistent with the results of a full body of multiconfigurational quantum chemical computations¹⁶⁻¹⁸ that were initiated at the end of the '90s. Such computations revealed: (i) the existence of low-lying CIs in rPSB11, (ii) the relationship between electronic origin and structure of the isomerization coordinate of rPSB11 and (iii) the molecular modes dominating the rPSB11 excited-state progression. In the present review article, we will highlight computational work focusing on the structure of CIs, the mapping of reaction paths, and trajectory simulations. In fact, each section of this review is organized such that it mirrors this section.

2.2 Computational studies: Rh isomerizes via a two-state, three-mode mechanism.

It is instructive to discuss at this point how computational studies connect with the experimental results i-iii introduced above and, at the same time, provide a motivation for the notions that will be examined in the following six sections. On the other hand, the reader may also skip the present subsection without loss of continuity, and revisit it at a later stage after reading Sections 3 and 4 if they feel they would prefer a more

general introduction to the theoretical terms and concepts used in this subsection.

The impact of the the existence of low-lying CIs in rPSB11 is apparent when comparing Fig. 2.2A and Fig. 2.2B illustrating the mechanism proposed in Miller and Mathies' contributions^{12,14} with that given in an early report by Schoenlein *et al.*,¹⁰ respectively. In contrast to the early view proposing decay at an avoided crossing (AC), the population transfer from S_1 to S_0 is presently assigned to the efficient internal conversion typical of CIs. Building on the seminal work of Michl and Bonačić Koutecký,¹⁹ a CI mediating the isomerization of a reduced rPSB11 model has been computed in 1997²⁰ shortly after CIs were shown to be implicated in common organic photochemical reactions. In the same report, it was shown that a barrierless coordinate connects the vertical excitation region to the CI located at the bottom of the excited-state potential energy surface.¹⁵ Also, the electronic structure of the excited state driving isomerization was assigned to a Bu-like (charge-transfer) rather than Ag-like (diradical) wave function symmetry (see also Section 3.1): a result that has been used by Miller and collaborators¹⁴ in their analysis of the coherences assigned to the excited-state HOOP mode describing the rPSB11 relaxation.

The second result, the relationship between electronic origin and structure of the isomerization coordinate of rPSB11, is a consequence of the Bu-like S_1 electronic structure. Promotion to a Bu-like state leads to an instantaneous charge-transfer process effectively unlocking the double bonds and locking the single bonds of rPSB11. This induces a stretching relaxation along C11=C12 and other conjugated bonds contributing to a BLA mode (this is quantified by the difference between the average of the double-bond lengths and the average of the single-bond lengths of the conjugated chain). The resulting =C9-C10=C11-C12=C13-C14= pattern (compare the ground-state and excited-state double-bond pattern in Fig. 2.3A) allows for a facile torsion about the reactive C11=C12 bond. In Rh, this torsion is accompanied by torsional motion along the adjacent C9=C10 bond.^{20,21} Such a mechanism, termed the bicycle-pedal coordinate, was first proposed by Warshel²² and confirmed for rhodopsins from different organisms^{23,24} after the development of CASSCF-based quantum mechanics/molecular mechanics models (*i.e.* full protein models).²⁵ While such a confined coordinate facilitates isomerization in the tight protein cavity, it has been shown that isolated rPSB11²⁶ and quantum mechanical/molecular mechanical (QM/MM) models²¹ feature essentially the same bicycle-pedal coordinate that, however, is aborted after decay to the ground state (*i.e.* the C9=C10 twisting reverts to the original configuration yielding isomerization at C11=C12 exclusively). A more detailed description of these results will be given in Section 4.3.




Figure 2.3. Reaction coordinate driving the isomerization of bovine rhodopsin. A. Change in the conjugation pattern upon photoexcitation of rPSB11 to a Bu-like state. B. QM/MM excited-state trajectory (in red) simulating the photoisomerization of Rh (data from ref. ²¹ and ²⁷) and representing the center of a coherent population (*i.e.* the vibrational wavepacket). The trajectory is dominated by progression along three modes: BLA (difference between average single and double-bond lengths of the rPSB11 backbone in Å), HOOP (in degrees) and the bicycle-pedal mode incorporating the C11=C12 torsion (in degrees). The initial 5 fs time segment is dominated by BLA but the HOOP progression is active on a 15 fs timescale. The dashed curve provides a qualitative representation of the projection of the intersection space (indicated with the IS label) of the torsion-HOOP sub-space. As we will mention in Section 3.2 and discuss in the appendices, the element of the intersection space are geometrically and energetically distinct CI points.

The third finding, the modes dominating the rPSB11 excited-state progression, is related to the second and refers to an early report by Gonzalez-Luque *et al.*¹⁶ where rPSB11 models were used to investigate the isomerization coordinate. It was shown that the aforementioned relaxation along the BLA mode (incorporating the reactive C11=C12 stretch of Fig. 2.3A) precedes the reactive C11=C12 torsion. The result pointed to a coherent two-state two-mode motion of the Rh population. Computational evidence for the participation of the HOOP mode in the coherent excited-state relaxation came later when semi-classical trajectory computations became possible.^{27,28} The development of QM/MM models of Rh^{21,25,29-31} in the first decade of 2000 allowed for the simulation of the full 200 fs dynamics probed via femtosecond-transient-absorption spectroscopy showing that three modes are involved in the excited-state motion. The results^{21,27,29,32-34} reported in Fig. 2.3B display a progression dominated by the BLA, HOOP and C11=C12 torsion modes held responsible for the observed coherent features.


Quantum chemical calculations have, as discussed above, provided a framework for the mechanistic/theoretical understanding of the photochemistry of dim-light vision. However, the

reliability of computational studies depends critically on the quality of the computer model used to represent the protein and the light-induced processes in the chromophore. Such models are complex and, currently, not univocally defined. For this reason they require a validation based on the simulation of observed quantities (e.g. absorption maxima and excited-state lifetimes) or the prediction of quantities that could be experimentally verified. For instance, one of the Rh models²⁷ introduced above predicted a HOOP mode with a ~ 40 fs period (~ 830 cm^{-1}) and a BLA mode with a ~ 20 fs period (~ 1670 cm^{-1}) and thus qualitatively consistent with the values recently reported by Miller and collaborators for the HOOP (746 cm^{-1}) and C11=C12 stretch (1679 cm^{-1}). Such comparisons support the initial relaxation mechanism provided by the model which points to a C11=C12 torsional progression reaching a 40° value at 50 fs.

Validated models are also needed to open new alleys for the comprehension of key properties of double-bond photoisomerization such as excited-state lifetime and quantum yield. For instance, an effect still under computational investigation is associated with the intersection space structure of rPSB11³⁵ (IS, see Fig. 2.3B and associated legend). The IS is a $(N-2)$ -dimensional nuclear coordinate subspace (N being the vibrational degrees of freedom of the system) formed by CI points. The CI of Fig. 2.2A and 2.3B is therefore only one point along the IS. Multiconfigurational quantum chemical calculations show that the IS may be approached not only via large, ca. 90° C11=C12 torsions, but also via progression along the BLA and HOOP modes.²⁷ Thus it could be possible to achieve a CI region at smaller value of the C11=C12 torsion when the progression along the HOOP mode approaches the IS (this is schematically represented by the dashed curve in Fig. 2.3B). See Section 3.4 for further details.

3. THEORY OF DOUBLE-BOND ISOMERIZATION

The Rh case study reviewed above shows that a theoretical framework is needed for interpreting or planning experiments designed to reveal the significance of double-bond isomerization in light-responsive proteins. Accordingly, in the present section we review the basic *theoretical and mechanistic concepts* for double-bond isomerization that have emerged from quantum chemical studies. Note that a discussion of the *methodologies* that may be employed in these studies is available in Appendix I. Here, we continue to focus primarily on protonated Schiff bases as paradigmatic examples of isomerizing chromophores.



Scheme 3.1. Schematic representation of photochemical (full arrows) and thermal (dotted arrows) isomerization paths. The two paths develop along different reaction coordinates. In fact, the CI is located energetically above the transition state controlling the thermal isomerization (the thermal transition state is indicated with the label TS and corresponds to a first order saddle point located along the S_0 potential energy surface), features a different geometrical structure, and drives a far-from-equilibrium process. $\Delta E(S_1-S_0)$, $\tau_{cis \rightarrow trans}$, and E_a^T (in red) are computable quantities. As shown at the top-right, quantum yields (QYs) can also be computed as the ratio of non-adiabatic trajectories reaching the isomerized chromophore (N_p) over the total number of computed trajectories launched (N_t). The competing photoproduct photoisomerization (dashed arrows) yielding the initial reactant may also be computationally investigated as will be discussed in Section 4.2.1.

To understand how Nature has utilized the simple double-bond isomerization motif to drive, very efficiently and effectively, different photobiological processes, two important questions come to mind. First, why has Nature selected certain organic chromophores in photobiological systems rather than others? While the answer to this first question depends on different factors, including the availability of biosynthetic pathways leading to specific conjugated frameworks, in this section we provide a partial answer by focusing on the factors that determine light sensitivity at the molecular level. Second, how does Nature optimize the properties of the chromophore (e.g., to maximize sensitivity to light) by tuning its environment? Indeed, the chromophore environment (i.e. the amino acid composition of the protein cavity) can modulate its maximum absorption wavelength (λ_{max}), its excited-state lifetime ($\tau_{cis \rightarrow trans}$), its photoisomerization “quantum yield” (QY), and the barrier (E_a^T) controlling the rate of the competing thermal isomerization.

The events included in the framed region of the photocycle of Fig. 2.1 are associated with specific features of the S_0 and S_1 potential energy surfaces of the chromophore. In the simplified Scheme 3.1, these S_0 and S_1 N -dimensional potential energy surfaces (with N being the number of independent nuclear degrees of freedom or independent vibrational modes) are reduced to one-dimensional curves along a reaction coordinate. This scheme can be used to assign computable quantities and their relation to experimental observables. Accordingly, light absorption is characterized by the vertical excitation energy $\Delta E(S_1-S_0)$ approximating the photon absorption wavelength λ_{max} of Fig. 2.2B. The absorbed photon promotes the chromophore from its S_0 equilibrium structure to the Franck-Condon (FC) point of its S_1 potential energy surface. At this

stage, to enhance light sensitivity of the system, it is important to increase the efficiency of the photochemical process. For example, the isomerization QY must be maximized in order to increase the ratio of reactive trajectories relative to non-reactive ones that do not trigger the photocycle. At the same time, any competing processes must be minimized. This includes thermal isomerization, which could compete with the photoisomerization process or, in the case of visual rhodopsins, creates the “background noise” in the signal and therefore lowers visual sensitivity. Therefore, an understanding of how to improve light sensitivity requires both calculation of QY, and rate of competing thermal isomerizations. To compute QY, one may start by considering the correlation proposed by Weiss and Warshel³⁶ and Mathies and coworkers.³⁷ The idea is that, consistently with the Landau-Zener model,³⁸ a larger velocity of the S_1 population moving toward the CI results in a higher reaction QY. This indicates that variation in QY correlates with the change in S_1 lifetime estimated, for instance, by computing the $\tau_{cis \rightarrow trans}$ time required to reach the CI or, more generally, the point of hop from S_1 to S_0 (i.e., the point when the trajectory algorithm decides to switch or “hop” from one state to another). This simplifies QY calculations considerably because one can get a sense of differences in QY in different systems by simulating the excited state lifetime of a single trajectory. Instead, as illustrated in the inset of Scheme 3.1, a QY value can be obtained in a more brute-force way by exploiting the S_0 and S_1 potential energy gradients and non-adiabatic couplings to compute a set (sometimes called ensemble or swarm) of non-adiabatic trajectories. Such calculations start with a simulation of the S_0 conformational population of the protein in the dark (i.e. ideally a representation of the Boltzmann distribution). This provides a finite number (N_i) of S_1 initial structures and velocities whose time evolution is simulated. The QY value can then be computed as the ratio between the number of trajectories successfully producing the isomerized chromophore (N_p) and N_i . Such explicit QY computations have been reported^{32,39,40} for Rh. Finally, the variation in thermal isomerization rate can be computed from the potential energy barrier E_a^T corresponding to the energy difference between the transition state (TS) and the protein equilibrium structure. In conclusion, the computational investigation of the photochemical and thermal isomerization of a double bond requires, at the very least, the calculation of the S_0 and S_1 state potential energy surfaces of the chromophore. As discussed below, the S_2 potential energy surface may also be implicated, so also needs to be computed for some cases.

3.1 Vertical excitation energies and conical intersections.

The selection of a suitable conjugated system hosting the reactive double bond may be discussed on the basis of the comparison of linear conjugate hydrocarbons and protonated Schiff bases. Here we focus on systems with five conjugated double bonds. Accordingly, we compare deca-1,3,5,7,9-pentaene to the nona-2,4,6,8-tetraeniminium cation, which are referred to as decapentaene and PSB5 respectively. Note that PSB5 serves as a model of rPSB11 (due to the large ca. 50° twisting about the C6-C7 bond, the C5=C6 double bond of rPSB11 does not efficiently conjugate with the rest of the rest of the

chain,^{41,42} making it effectively similar to a five double-bond conjugated protonated Schiff base).




Figure 3.1. Comparison of relative energies of FC and CI energies in the isoelectronic decapentaene and PSB5. The energies computed at the 3-root-SA-CASPT2(IPEA=0)//2-root-SA-CASSCF/6-31G* level of theory (the method acronyms are introduced in Section A.1.1). Note that the double slash notation indicates that energies were computed at the level of theory at the left, here 3-root-SA-CASPT2, and geometries at the second level of theory at the right, here 2-root-SA-CASSCF). A. Relative stability of the FC states and low-lying CIs of decapentaene. The schematic representation of the geometrical and mixed S_1/S_0 electronic structure of the CIs is also given. The dashed lines connect FC and CI structures for visibility. The arrows represent relaxation paths from the FC points. The framed structure represents the CI driving decay from an intermediate (P-state) in stilbene and is shown for comparison. The dashed segment connecting the two radical centers indicates that these electrons are singlet coupled. B. Relative stability of the FC states and low-lying CIs of PSB5. Even in this case the dashed segments connecting two radical centers indicate that these are coupled (i.e. they appear uncoupled only because they reside in two nearly orthogonal π -systems). In all cases the Lewis formula used for representing the electronic structure of the CI points represents mixed electronic structures. This is illustrated in Scheme 3.2 for the case of PSB5. Data from ref. 43.

The first three electronic singlet states (S_0 , S_1 and S_2) of the reference hydrocarbon decapentaene in its S_0 equilibrium C_{2h} structure can be classified as 1Ag, 1Bu and 2Ag. The 1Ag character corresponds to S_0 and has a closed-shell covalent electronic structure. The 1Bu state corresponds to S_1 and it is dominated by a singly-excited electronic configuration leading to a charge-transfer character, while the 2Ag state, corresponding to S_2 , is dominated by double excitations and has a diradi-

cal character. The selection rules for electronic transitions impose only the 1Ag to 1Bu and 1Bu to 2Ag transitions have non-zero oscillator strength.


In the discussion below the electronic character of the singlet electronic states of our model chromophores will still be classified using the C_{2h} symmetry labels 1Ag, 1Bu and 2Ag (i.e. related to a polyene hydrocarbon such as *tEt*-hexa-1,3,5-triene in its planar S_0 equilibrium geometry (italics stereochemical labels indicate the configuration of single (trans or cis) and double bonds (E or Z) respectively), even for structures with C_s symmetry like PSB5 or no group symmetry like any structure located along the PSB5 isomerization reaction coordinate.

Alternatively, the same states will be labeled $\Phi_{\text{COV/DIR}}$, Φ_{CT} , and Φ_{XDIR} , respectively, to explicitly indicate the character of the dominating electronic configuration (or diabatic state). Notice that along an isomerization path or a trajectory the character of the potential energy surfaces associated with S_0 , S_1 and S_2 (i.e. classified according their energy order which will always be, by definition, $S_0 < S_1 < S_2$) may change. Accordingly, one may have an S_0 potential energy surface with a dominating $\Phi_{\text{COV/DIR}}$ (i.e. here associated with a 1Ag) character in certain region and a Φ_{CT} (i.e. 1Bu) character in some other region, or a S_1 potential energy surface with a dominating Φ_{CT} character in a region and a $\Phi_{\text{COV/DIR}}$ or Φ_{XDIR} (i.e. 2Ag) character in a different one.

Note that later on, in Section 3.2, we will further distinguish $\Phi_{\text{COV/DIR}}$ as being either Φ_{COV} or Φ_{DIR} , depending on the geometry, but it is important to recognize that these two labels both describe the same electronic state. To elaborate, $\Phi_{\text{COV/DIR}}$ indicates that along a double-bond twisting path, the Φ_{COV} closed-shell electronic configuration dominating the S_0 state of the reactant (or product), smoothly changes into a Φ_{DIR} open-shell configurations representing an orthogonal diradical. This is equivalent to adopting a valence-bond view of a homolytic bond-breaking process where two spin-paired electrons become localized on distinct fragment π -orbitals and the bond breaking is described by a change in overlap between these singly-occupied orbitals. Therefore, Φ_{COV} and Φ_{DIR} are seen as different components of the S_0 electronic structure. Notice that Φ_{XDIR} is distinct from Φ_{DIR} . In fact, Φ_{XDIR} represent a diradical state obtained via electronic excitation at the untwisted reactant geometry and, as we will see below, may assume a tetraradical character upon double-bond twisting. When using the same valence bond view, Φ_{CT} corresponds to an electron transfer from one fragment π -orbital to the other.

The energy of lowest-lying decapentaene CIs are shown in Fig. 3.1A. The diagram indicates that excitation in decapentaene occurs to the S_2 state, after which a 2Ag/1Bu CI between the S_2 and S_1 potential energy surfaces may be readily accessed from the FC point after photon absorption. After decay to S_1 , the evolution along the S_1 potential energy surface may

then lead to a 2Ag/1Ag CI where the decay to S_0 would occur. A qualitative analysis of the geometrical and electronic structure of these CIs indicates that: (i) the S_0 photoproduct may be reached with UV rather than visible light excitation, (ii) the E to Z isomerization at C5=C6 may occur via the 2Ag/1Bu CI, potentially generating an isomerized photoproduct on the S_1 potential energy surface, and (iii) the decay to S_0 occurs via a 2Ag/1Ag CI featuring a $-(\text{CH})_3-$ kink with a tetraradical character.⁴⁴ The "kink" geometrical and electronic structures support the possibility to simultaneously generate two different primary photoproducts (a cyclopropyl diradical intermediate and a Z-isomer) in addition to the possibility of regenerating the initial *E*-isomer. Points i-iii make it clear why linear polyenes would not be favorable chromophores for absorbing photons of visible light selectively generating the isomerization product without going to secondary (e.g. cyclization) photoproducts. In other words, a protein cavity hosting a decapentaene chromophore would have to severely modify, through environmental effects, the three potential energy surfaces to achieve favorable spectroscopic and reactivity properties. Furthermore, in a short polyene hydrocarbon, the charge separation occurring upon vertical excitation (i.e. via the 1Ag to 1Bu transition) is spread along the chain and does not have a direction, making it more difficult to control the chromophore properties via local electrostatic effects (i.e. via interaction with specific polar residues located in the protein chromophore cavity).



Scheme 3.2. Lewis resonance formulas representing the electronic configurations degenerate at a 1Ag/1Bu CI in PSB5 featuring a ca. 90° twisted C5=C6 double bond. At the FC point the S_0 and S_1 adiabatic electronic states are dominated by the 1Ag and 1Bu configurations respectively. Note that 1Ag corresponds to Φ_{COV} at the FC point but to Φ_{DIR} at the twisted CI geometry. However, in the CI vicinity, the S_0 and S_1 states display a mixed and interchanging electronic character. The crossing of 1Ag and 1Bu electronic characters at the CI is represented by the resonance formula at the right.

The results discussed above indicate that, in decapentaene, there is no direct access to a low-lying 1Bu/1Ag CI where decay to S_0 may occur directly. However, an accessible 1Bu/1Ag CI has been shown to exist in arylethenes such as stilbene (see inset in Fig. 3.1A). With respect to the FC point, such a CI is found⁴⁵ lower than the decapentane 1Bu/2Ag CI and is related to an observed S_1 intermediate called the "P-state". This intermediate features a highly twisted double bond with a pyramidalized sp^3 carbon hosting an anionic center and a positive charge partially delocalized on a benzyl moiety. Thus, *E*-stilbene decays to S_0 from the initially populated S_1 state leading to its *Z*-isomer after having accessed the P-state and the nearby 1Bu/1Ag CI. However, this decay mechanism does not occur for decapentaene.

We will now compare the energy of lowest-lying PSB5 CIs to the one described above for decapentaene. Fig. 3.1B shows a

1Bu/1Ag CI that could be readily access from the FC point. In other words, in the absence of an S_1 energy barrier, the events initiated by the photon absorption leads directly to S_0 without involving a 2Ag/1Bu CI, which is now located at a significantly higher energy. Most importantly, due to the positive charge residing on the chromophore, the 1Bu/1Ag CI features a single 90° twisted $C5=C6$ double bond with no other major skeletal distortions. This CI is thus geometrically different from the stilbene 1Bu/1Ag CI characterized by an S_1 state with charge separation rather than by a charge transfer. In fact, the S_1 state of near the PSB5 CI may only feature a neutral lone pair localized on the nitrogen atom of the Schiff base functional group with a positive charge far away from it. A qualitative analysis of the geometrical and electronic structure of the PSB5 CI indicates that: (i) the CI may be reached with visible light excitation, (ii) the isomerization of the central $C5=C6$ double bond occurs via a fully (90°) 1Bu/1Ag CI potentially generating exclusively an S_0 isomerized photoproduct. Point i and ii appear to be favorable features for a biological chromophore.




Figure 3.2. $\Delta E(S_1-S_0)$ and $\Delta E(S_2-S_1)$ vertical excitation energies variations of protonated Schiff base chromophores. A. Environmental effect on the $\Delta E(S_1-S_0)$ value of bovine rhodopsin (ca. PSB5). The balloons represent the charge distribution (blue positive and red negative) computed in the S_0 and S_1 states. The level diagram illustrates, schematically, the $\Delta E(S_1-S_0)$ and $\Delta E(S_2-S_1)$ variations starting (left to right) from the chromophore *in vacuo* (λ_{\max} ca. 610 nm),⁴⁶ moving to the bovine rhodopsin cavity (λ_{\max} ca. 500 nm)⁴⁷ and to the chromophore in methanol solution (λ_{\max} ca. 440 nm)^{48,49} where a negative “counterion” strongly stabilizes S_0 with respect to S_1 . More specifically, the counterion stabilizes preferentially the 1Ag and 2Ag states (i.e. 2Ag has diradical character and its charge distribution is similar to the 1Ag charge distribution) with respect to the charge-transfer (1Bu) state. Notice that due to the stabilization of the 2Ag state, the energy difference between the 2Ag and 1Bu states is much reduced in the solution environment. B. CASPT2(IPEA=0)//2-root-SA-CASSCF/6-31G* computed vertical excitation energies for protonated Schiff bases of different length (from PSB6 with six conjugated double bonds to PSB4 with four conjugated double bonds) and equilibrium geometries. Data from ref.²³.

A protein cavity hosting a “PSB5-like” chromophore such as rPSB11, would easily modulate the $\Delta E(S_1-S_0)$ and therefore the λ_{\max} . In fact, in contrast with decapentaene, the charge

transfer occurring upon S_0 to S_1 vertical excitation would result in a positive charge translocation (delocalization) from the $N=C2$ bond to the β -ionone ring. Such a charge translocation is a general feature of protonated and alkylated Schiff bases and is documented in Fig. 3.2A for rPSB11. The molecular environment (i.e. the protein and solvent cavities) interacts with the charge distributions of the S_0 and S_1 states differently and, ultimately, modulates the $\Delta E(S_1-S_0)$ value and thus the corresponding λ_{\max} . The mechanism through which proteins influence the absorption wavelength λ_{\max} of the chromophore is known as color tuning or spectral tuning, a topic that has been widely studied computationally⁵⁰⁻⁵³ and experimentally.^{54,55}

Other $\Delta E(S_1-S_0)$ and $\Delta E(S_2-S_1)$ modulating effects are achieved by changing the effective length of the conjugated chain via single bond twisting (e.g. the $C6-C7$ double bond in rPSB11). The magnitude of such effect has been calculated²³ and the corresponding values are displayed in Fig. 3.2B bottom. An additional effect that also changes $\Delta E(S_1-S_0)$ and $\Delta E(S_2-S_1)$ in the same direction as increasing the conjugated chain length is by modulating the BLA geometry. This effect can be inferred comparing Fig. 3.2B top, with 3.2B bottom. The top panel shows energies for the S_1 equilibrium structure of a constrained planar (i.e. untwisted) conjugated chain. At a planar geometry, the S_1 electronic structure of PSB5 is dominated by a Φ_{CT} electronic configuration and, therefore, its equilibrium geometry features an almost inverted BLA pattern. Similar modulating, but more complex, effects are expected when looking at the energies of the 1Bu/1Ag and 2Ag/1Bu CIs suggesting that the reaction path leading from the FC point to the CIs may also be modulated by the environment surrounding the chromophore.




Figure 3.3. Conical intersections for PSB5. A. Relative stability of the Franck-Condon states and low-lying 1Bu/1Ag and 2Ag/1Bu conical intersections of PSB5 for both isomerization around $C5=C6$ and $C7=C8$. Energies were computed at the 3-root-SA-CASPT2(IPEA=0)//2-root-SA-CASSCF/6-31G* level of theory. The dashed lines are guides for the eye. The schematic representations of the geometrical and mixed S_1/S_0 electronic structure of the conical intersections are also given. The dashed

segment connecting the two radical centers indicates that these electrons are coupled. Data from ref. 43.

Above we have exclusively focused on the 2Ag/1Bu CI associated with the twisting about the central C5=C6 double bond of PSB5. However, similar CIs also exist which are associated with the twisting about other double bonds in the chromophore. As shown in Fig. 3.3 for the 2Ag/1Bu CI mediating the C7=C8 double-bond isomerization, such CIs have different relative energies. Therefore, in general, a mixture of isomerized photoproducts (different stereoisomer of PSB5) may be expected, which will depend not only on their relative energies but also on the details of the reaction coordinates leading to the corresponding CI. It can be hypothesized that the chromophore environment (i.e., again, the protein cavity or the solvent cavity and counterion) controls the isomerization selectivity by stabilizing a specific CI and/or favoring a specific reaction coordinate. Indeed, as we will review in Section 3, the cavities of retinal proteins favor single isomerizations about either the C13=C14 or the C11=C12 double bond of their retinal chromophores. However, secondary isomerization products (e.g. involving the C9=C10 and C7=C8 double bonds) have been reported.




Figure 3.4. Schematic representation of the S_0 and S_1 potential energy surfaces plotted along the BP of the CI. The structures and BP vectors correspond to the CI of the PSB11 chromophore of bovine rhodopsin. A loop located on the BP and centered at the CI is shown. A projection of the loop on the S_0 potential energy surface goes from a Φ_{DIR} region (green) to a Φ_{CT} region (brown) and then returns to a Φ_{DIR} region. The Newman projection on the top-right defines the HOOP internal coordinate as corresponding to the dihedral angle representing the displacement from the "ideal" equilibrium position (dashed segments). Adapted with permission from refs. 56 Copyright 2012 The American Association for the Advancement of Science and 57. Copyright 2006 Taylor & Francis.

3.2 Structure of the photochemical and thermal reaction paths. A deeper understanding of the possible effects of the protein cavity on the non-adiabatic S_1/S_0 transition driving a double-bond photoisomerization calls for the geometrical and electronic characterization of the branching plane (BP) of the CI.^{15,58-60} This is the plane (also called **g-h** plane)⁶⁰ defined by the molecular modes corresponding to the gradient difference and derivative coupling vectors computed at the CI. These vectors are introduced more fully in Section A.I.1. A pictorial representation of the BP and intersecting potential energy sur-

faces associated with the S_0 and S_1 states of the rPSB11 chromophore is given in Fig. 3.4. One of the vectors is dominated by a combination of twisting deformation about the reactive C11=C12 double bond and a HOOP wag. The other vector corresponds to a BLA stretch.⁵⁷ An analysis of the S_0 and S_1 charge distributions along the BP reveals a relationship between the electronic structure of the S_0 and S_1 states in the CI region and in the FC region. The main point is that, consistently with a quantum mechanical theorem called the geometric phase,⁶¹ the S_1 state changes its electronic character along a small loop centered at the CI and residing on the BP. More specifically, in rPSB11, its character passes two times (e.g. roughly every 180°) from a Φ_{CT} character typical of the S_1 state at the FC point, to the Φ_{DIR} character associated with the S_0 state at the FC point. In Fig. 3.4 these different regions of the S_1 potential energy surface are colored in brown and green, respectively. The geometric phase theorem explains why an opposite electronic state change is observed for S_0 , namely when one moves along the same loop. Therefore, on the S_0 potential energy surface there are regions dominated by a Φ_{CT} character rather than the Φ_{DIR} (or Φ_{COV}) character expected considering the S_0 state in the FC region.




Figure 3.5. Schematic representation of the structure of the potential energy surfaces driving the ultrafast double-bond photoisomerization and thermal isomerization of rPSB11 in Rh. A. S_1 potential energy surface driving the photochemical isomerization. The changes in the charge distribution of the chromophore during S_1 evolution and after relaxation to S_0 are also shown with bubbles. The bubbles represent positive (blue) and negative (red) charges on the atoms of the chromophore framework. The curly arrows indicate the intramolecular electron transfer process driving the charge translocation occurring upon photoexcitation and also during the progression and decay of the excited state. The top structure corresponds to the S_1 structure of the chromophore in the vicinity of the CI (the framed region corresponds to the "double cone" seen in Fig. 3.4). The associated electron transfer is the one occurring during the hop from S_1 to S_0 . B. Mechanism of the S_0 thermal isomerization. This transformation is controlled by two transition states (TS_{CT} and TS_{DIR}) located along the BLA

coordinate and on opposite sides with respect to the CI. TS_{CT} has a charge-transfer character (Φ_{CT}) and therefore a 1Bu-like electronic structure. In contrast, TS_{DIR} has the diradical character (Φ_{DIR}) of a 1Ag-like electronic structure that becomes a closed-shell covalent electronic structure at the reactant (Rh) and product (bathoRh) geometry. The bubbles represent positive and negative charges on the atoms of the chromophore framework at the corresponding transition states. The cross-section along BLA (computed at the 2-root-SA-CASPT2//CASSCF/6-31G* level of theory) provides information on the energy difference between TS_{CT} and TS_{DIR} and shows that in Rh the transition state controlling the isomerization is TS_{CT} . Adapted with permission from refs. ⁵⁶ Copyright 2012 The American Association for the Advancement of Science and ⁵⁷. Copyright 2006 Taylor & Francis.

The qualitative structure of the S_0 and S_1 potential energy surfaces driving the rPSB11 chromophore evolution is given in Fig. 3.5A. In order to drive an ultrafast chemical reaction it is not enough to have a CI with a suitable geometrical and electronic structure and BP. The S_1 potential energy surface must also feature a barrierless (or nearly barrierless) reaction path connecting the FC point to the CI. The properties of such a path (i.e. reaction coordinate, slope and electronic character) may be controlled by the interactions with the protein environment. In principle, such interactions may slow down or even stop the isomerization if an energy barrier is created between the FC point and the CI. However, this is not the case for the S_1 potential energy surface illustrated in Fig. 3.5A, which shows a barrierless potential energy surface consistent with that of Rh.²¹

In protonated Schiff bases that are free to isomerize, there are specific molecular structure changes describing the path towards the CI (e.g. BLA, HOOP, and C11=C12).^{21,27} In the following sections we will see that these same changes also describe the excited state relaxation of other biological chromophores whose S_1 state is dominated by a charge-transfer excitation Φ_{CT} of their π -systems. Such a path is described by progression along multiple coordinates (or modes), and cannot be effectively visualized in a 2D representation like the one of Fig. 3.5A. In fact, already a 3D representation would reveal that a CI is actually not an isolated point but it is part of a N-2 dimensional "line" of CIs called intersection space or IS (see also Fig. 2.3). While here, we do not further discuss the relationship between the S_1 isomerization coordinate and the IS, in Appendix II we show that it has been possible to map the S_1 paths connecting the FC structures of different isomers of a minimal rPSB model to the corresponding CIs as well as the position of these CIs within an IS segment.⁶² This information is useful for discussing the dynamics of the S_1 deactivation which does not occur by following a single "static" path but, rather, by following different trajectories possibly reaching different IS points.

In Fig. 3.5B we display a schematic representation of the S_0 potential energy surface of Rh. As we will discuss in Section 3, the features of this surface determine if the thermal isomerization of the Rh chromophore may or may not compete with its photoisomerization by generating "thermal noise." Indeed, consistently with Scheme 3.1, such competition depends on the activation barrier E_a^T associated with an S_0 transition state. However, it has been shown that, as a consequence of the potential energy surface topography imposed by the Rh CI, the

thermal isomerization may occur via two geometrically and electronically distinct transition states: TS_{CT} , characterized by the Φ_{CT} configuration, and TS_{DIR} , characterized by the Φ_{DIR} electronic configuration.⁵⁶ Note that dominating transition state structure (i.e. the one associated with the lower E_a^T value) would be also responsible for the thermal back-isomerization of the chromophore to the original reactant.




Figure 3.6. Non-adiabatic trajectory for bovine rhodopsin released from the FC point with zero initial velocities (FC trajectory). The trajectory is computed at the 2-root-SA-CASSCF/6-31G*/AMBER level of theory.²¹ A. S_1 and S_0 energy profile computed along the trajectory. Notice that at ca. 100 fs the S_1 state crosses the S_0 state. At this point the system hops from the S_1 to the S_0 state and continues its motion along the S_0 potential energy surface. B Geometrical changes along the S_1 branch (i.e. until the hop point) of the same non-adiabatic trajectory. The plotted geometrical parameters contribute to one of the BP vectors (C10-C11-C12-C13 twisting in black open squares and HOOP in blue open circles). We also report the coupled C8-C9-C10-C11 twisting (dashed curve) forming the S_1 bicycle-pedal reaction coordinate²² discussed in the introduction. Notice that immediately after the hop the C8-C9-C10-C11 dihedral twisting inverts direction (not shown) and the bicycle-pedal motion is ultimately aborted.²¹ C. Changes in the charge distribution along the same trajectory. The parts of the trajectory dominated by a Φ_{CT} electronic structure are marked in brown while the one dominated by a Φ_{DIR} electronic structure are marked in green. The QM/MM model used is based on a computer time-saving CASSCF(12,12)/6-31G*/AMBER level of theory, with an active space comprising the full π -system and with methyl groups and part of the β -ionone ring in the MM subsystem to reduce the size of the QM subsystem. Figures are adapted with permission from ref. ²¹. Copyright 2007 National Academy of Sciences.

3.3 Time evolution of the excited-state population. Above we have focused on the structure of the S_1 potential energy surfaces controlling the chromophore photoisomerization. However, a barrierless reaction (i.e. not controlled by a transition state) can only be satisfactorily described by looking at the evolution of the geometrical and electronic structure of the system as a function of time. In Fig. 3.6 we report the results of a single non-adiabatic trajectory calculation providing information on such process for Rh.²¹ As shown in Fig. 3.6A,

the time evolution of the S_1 and S_0 energy profiles, S_1 - S_0 gap. In Ref. ²¹ the authors also report the oscillator strength evolution (proportional to the transition charge-transfer character and intensity at the corresponding wavelength). The same calculation provides information on the evolution of the associated geometrical parameters and on the followed reaction coordinate which has already been introduced in Fig. 2.3. In Fig. 3.6B we display the evolution of the twisting deformation about the reactive C11=C12 double bond, unreactive C9=C10 double bond, and HOOP parameters. In this way it has been possible to document the "aborted" bicycle pedal reaction already mentioned in the introduction. The same calculation also provides information on the evolution of the charge distribution and, therefore, Φ_{CT} or Φ_{DIR} character of the chromophore electronic structure during the S_1 motion and after the hop to S_0 . Notice that such a hop does not necessarily occur at exactly the CI point but, rather, as mentioned in the previous section and detailed in Sections A.I.3 and A.I.4, the hop occurs along the "volume" surrounding the CIs belonging to an IS "segment".

In Fig. 3.6C it can be seen how the initial Φ_{CT} character evolves according to the process schematically anticipated in Fig. 3.5A.²¹ Notice that a wider region bearing a Φ_{CT} character in S_0 implies a wider Φ_{DIR} dominated region on S_1 . This wider S_1 region would imply a charge evolution significantly different from the one of Fig. 3.6C. In particular, after an initial charge transfer from the Schiff base to the hydrocarbon region, the charge partially gets back to its original position and the fully twisted chromophore would thus have a diradical character.

The trajectory of Fig. 3.6 only describes the motion of the molecule after a vertical transition from its S_0 equilibrium structure to S_1 and with no initial velocities. Accordingly, such a trajectory only provides *qualitative* information on the dynamics of the photo-excited population (also called vibrational wavepacket), as one misses the statistical description of the motion prepared, for instance, by a laser pulse. Such a population includes a large number of molecules differing in structural displacement and velocities and consistent with the Boltzmann distribution at the experimental temperature. It is thus the dynamics of the population that one needs to simulate to provide parameters that can be directly compared with the experiment and, most importantly, to determine QY values. As schematically illustrated in Scheme 3.1, the vibrational wavepacket time evolution can be simulated by computing a finite set of non-adiabatic trajectories each associated with a structure and velocity extracted from a simulated Boltzmann (or Wigner^{63,64}) distribution. Such calculations have actually been reported for Rh by various groups.^{32,34}

The simulation of the population dynamics with a suitable number of trajectories can be used to simulate transient spectroscopic quantities and compare them with the results of pump-probe transient spectroscopy experiments. For instance, the simulation of transient emission and absorption spectra is schematically illustrated in the inset of Fig. 3.7A which primarily report the energy profile computed along five different

non-adiabatic trajectories launched from structures and velocities selected from some initial Boltzmann distribution.³² More specifically, it is possible to trace, along entire sets of trajectories, the evolution of different energy gaps and oscillator strengths between different states, as also schematically illustrated in the inset.




Figure 3.7. Population dynamics of Rh simulated via non-adiabatic trajectories. A. S_1 and S_0 energy profiles along five non-adiabatic (surface-hop) trajectories belonging to a Boltzmann distribution. The black circle represents S_1 to S_0 hop points. The inset above provides a schematic representation of S_1 to S_0 transient emission and S_1 to S_n (with S_n being an upper excited state) and S_0 to S_1 transient absorption events as a function of time. B. Observed (panel a) and computed (panel c) transient electronic spectra of bovine rhodopsin. The simulated transient spectra (blue for stimulated emission and yellow for absorption) have been obtained by computing 40 semi-classical trajectories. The observed and simulated changes in emission (positive)/absorption (negative) at specific wavelengths, are given in panels b and d, respectively. Adapted from ref. ³². Copyright 2010 Nature Publishing Group.

In a more recent study by Polli et al.⁴⁰ the photoisomerization quantum yield of isorhodopsin (isoRh), a secondary photoproduct of the photoisomerization of bovine rhodopsin featuring a rPSB9 chromophore (see Fig. 3.8A), was simulated using a larger set of semi-classical trajectories. The results pro-

vided an overview of different competing processes occurring during the relaxation of the initial isoRh S_1 population. As displayed in Fig. 3.8A, part of the population simply returns to a vibrationally excited form of the original S_0 reactant. Part of the population undergoes a single isomerization event at either the C11=C12 double bond, generating a new intermediate with 11-*cis* and 9-*cis* double bonds (termed rPSB911), or at the C9=C10 double bond to give bathorhodopsin (containing a rPSBAT chromophore). Finally, a fraction of the population generates Rh (containing a rPSB11 chromophore) via a simultaneous isomerization of the C11=C12 and C9=C10 double bonds.

In principle, the analysis of the resulting trajectory ensemble not only provides a test for the validity of the mechanistic information extracted by minimum energy path (MEP) computations and single-trajectory probing discussed above, but also can be used to directly simulate the evolution of the signals obtained via transient absorption spectroscopy and compare with experiment. For instance, as shown in Fig. 3.7B, the simulation of time-resolved absorption spectra has been reported for Rh. The matching between experimental and simulated spectra³² provides a validation of the mechanistic hypothesis originated with previous investigations. In particular, it confirms the involvement of a barrierless relaxation path and decay in the vicinity of a CI (supported by the progressively red-shifting wavelengths associated with the observed stimulated emission signal) during the internal rPSB11 photoisomerization process. Such studies employing an ensemble of trajectories therefore can be used to compute more reliable excited-state lifetimes ($\tau_{\text{cis} \rightarrow \text{trans}}$), compared to single-trajectory calculations.

The calculation of hundreds, if not thousands, of non-adiabatic trajectories allows QY computations in the way anticipated in Scheme 3.1. These are calculated as the N_p/N_t ratio between the number N_p of trajectories reaching the S_0 potential energy surface valley centered on the photoproduct energy minimum (e.g. for Rh, this would be the bathoRh intermediate) and the total number of non-adiabatic trajectories hopping from S_1 to S_0 . However, presently, the accurate computation of QY is limited by the computational cost associated with running many trajectories at an accurate level of theory. Indeed, only indicative values have been reported on the basis of less than 100 trajectories.^{32,40} For instance, the QY of rPSB11 isomerization in Rh was estimated using 40 trajectories and found to be 0.61 compared to a 0.64 experimental value.³²

3.4 Quantum yields and isomerization stereochemistry.

The example above indicates that non-adiabatic trajectories can be employed to investigate the selectivity (see Fig. 3.3 and related text) of double-bond photoisomerizations. In fact, selectivity is quantitatively described by QYs of isomerizations controlled by distinct and competing CI channels. For isoRh a representation of the structures of these CIs (CI_1 and CI_2) are given in Fig. 3.8B. CI_1 leads to isoRh reconstitution with a 0.6 QY, indicating that this is the major channel in isoRh photochemistry. However, the same channel also leads to a 0.2 QY of the derivative featuring the PSB911 chromophore and produced by following an aborted bicycle-pedal reaction coordinate ultimately leading to C11=C12 isomerization exclusively.

In contrast, the CI_2 channel is leading to a very limited (0.04 QY) production of Rh produced via two parallel double-bond isomerizations. This same channel mainly produces bathoRh (featuring the rPSBAT chromophore) through a single C11=C12 isomerization with ca. 0.15 QY. Such isomerization is driven by progression along the same aborted bicycle-pedal reaction coordinate but followed in the opposite direction.




Figure 3.8. Simulated photoisomerization of isoRh. A. Geometrical progression of the C9=C10 and the C11=C12 dihedral angles. The numbers indicate the average reaction times and final configurations. The + sign indicates a CW twisting motion while the - sign CCW twisting. The paths turn from green to red with the S_1 to S_0 surface hop at one of the two CIs. B. Distribution of S_1 to S_0 hop times at the two CI_1 and CI_2 decay channels. The QM/MM model used is based on a CASSCF(10,10)/6-31G*/AMBER level of theory, with an active space not comprising the full π -system and with methyl groups and part of the beta-ionone ring in the MM partition to reduce the size of the QM subsystem. Both figure adapted with permissions from ref. ⁴⁰. Copyright 2014 John Wiley and Sons, Inc.

As mentioned above, double-bond isomerization selectivity is determined by differences in QYs obtained, as anticipated in Scheme 3.1, via non-adiabatic trajectory calculations. These calculations are important because they provide direct access to a molecular-level description of the statistical process leading to high or low QYs. However, they do not provide direct access to a QY theory for ultrafast reactions. In fact, while several hypotheses and even experimental verifications^{37,65} of these hypotheses have been reported, the molecular factors controlling QYs in rhodopsins are still not established.

According to Weiss and Warshel³⁶ and Mathies and coworkers,¹⁰ for fast impulsive (rather than diffusive) dynamics there

exists a correlation such that a larger velocity of the S_1 population moving toward the CI correlates with a higher reaction QY. Although the existence and relevance of such relationship, which is based on a Landau-Zener model^{38,66-68} of the S_1 decay, is still under investigation,^{69,70} it is apparent that a decrease in the time required to reach the CI enforces a coherent S_1 dynamics. While a coherent dynamics regime appears to be necessary for the control of the QY value,⁷¹ the molecular-lever mechanisms that could exploit such a regime is still substantially unknown.

As mentioned above, QYs can be defined as N_p/N_t (i.e. the number of trajectories reaching the photoproduct P divided by the total number of trajectories launched). We now examine a set of results aimed at the understanding of the dependence of the photoisomerization "direction" (i.e. either toward the photoproduct or toward the reactant) of single non-adiabatic trajectories on critical molecular coordinates. The central idea is that the time evolution of certain coordinates determines in which way the reactive double bond reconstitutes immediately after the S_1 decay.^{27,39} The stereochemistry of such reconstitution depends on: (i) the nature of the electronic wave function, and (ii) the overlap between the fragment π -orbitals containing the paired electrons, which, in turn, depends on the chromophore geometry.




Figure 3.9. Schematic representation (adapted from ref. ²⁷ Copyright 2011 American Chemical Society) of the geometrical coordinates modulating the overlap between the two π -systems (represented by p-orbitals at C2 and C3 centers) that interact to form the isomerizing double bond of a retinal chromophore. These coordinates can also be thought of as vibrational modes with a given excited-state frequency. The frequency is low for the torsion mode (i.e. the C1-C2-C3-C4 dihedral angle), larger for the HOOP mode (i.e. the difference between the H-C2-C3-R and the C1-C2-C3-C4 dihedral angles) and even larger for the d_{CC} mode (d_{CC} is part of the BLA mode mentioned above). R=H for the isomerization of the C11=C12 double bond in animal rhodopsins while R=Me for the isomerization of the C13=C14 double bond in most microbial rhodopsins.

In Fig. 3.9 we introduce the molecular modes affecting the π -overlap across the photoisomerizing double bond and in Fig. 3.10 we introduce the relationships between these modes and the stereochemical directionality of the double-bond photoisomerization. We start by considering a structure where the

HOOP and torsion are close to -90° (see the central structure of Fig. 3.10A). At this highly twisted structure the overlap of the π -systems of the C1C2 and C2C3 fragments (represented by p-orbitals at the C2 and C3 centers respectively) is close to zero. We assume that this structure corresponds to a S_1 point close to a CI and therefore about to hop from S_1 to S_0 . We also assume that the torsion is slow with respect to the HOOP, consistently with the associated frequencies of these modes (see Fig. 3.9 bottom).




Figure 3.10. Coordinates affecting the isomerization direction and QY. A. Effects of the HOOP mode on the double-bond photoisomerization direction (see text). The central structure refers to a situation in which the HOOP and torsion modes are close to -90° . B. Schematic representation of hypothetical effects of the change in HOOP frequency or initial conditions on the photoisomerization direction. The left diagram represents a reference situation. In part adapted from ref. ²⁷. Copyright 2011 American Chemical Society.

We then look at the effect of the vibrational deformation towards negative (see Fig. 3.10A top) and positive (see Fig. 3.10A bottom) HOOP values, which increases the orbital overlap between the two π -systems. One finds that negative and decreasing HOOP values prompt formation of the E reactant (large b-b' and a-a' overlap) while positive and increasing values prompt formation of the Z product (large b'-a and b-a'

overlap). Notice that these rules are valid for the M-helical central structure of Fig. 3.10A and associated counterclockwise isomerization motion (i.e. a motion where the negative C1-C2-C3-C4 value is increasing going from -180° to -90°). The rules would invert for the corresponding P-helical structure and clockwise isomerization featuring positive rather than negative HOOP and torsion. We conclude that the isomerization directionality (i.e., towards reactant or product) may be controlled by the HOOP direction of change at the hop point. These results have been computationally verified in several works^{39,40,72,73} where it is clearly shown that the majority of the trajectories follow the above rules not only in terms of velocities (i.e. sign of the HOOP velocity in degrees/fs) but also HOOP acceleration which is proportional to the HOOP gradient as reported in Fig. 3.11.³⁹




Figure 3.11. Dynamics of the 11-*cis* to all-*trans* photoisomerization of Rh. A. Time evolution of the torsion (i.e. the C10C11C12C13 dihedral angle) coordinate. The green lines correspond to motion in S_1 and the red lines denote S_0 motion. B. Velocities of the Torsion and HOOP coordinates (HOOP=HC11C12H-C10C11C12C13, see also below). Green triangles (red triangles) denote trajectories yielding the all-*trans* photoproduct (reactive trajectories). The bar graphs count the number of events in the corresponding left and right halves of the diagrams. Notice that, consistently with the rules discussed in the text and illustrated in Fig. 3.10A, the reactive trajectories correlate with negative HOOP velocities. Adapted with permission from ref. ³⁹. Copyright 2011 Royal Society of Chemistry and related supporting information.

In Fig. 3.10B we show how, according to the above rules/theory, the E (C1-C2-C3-C4 ca. 180° or -180°) to Z (C1-

C2-C3-C4 ca. 0°) photoisomerization directionality can be affected by the HOOP frequency and phase. Such variations may be imposed by a change in the retinal chromophore environment, such as those imposed by mutations of the protein, thus influencing the QY magnitude and isomerization selectivity. As stressed above, the hop occurs when the two orbitals are nearly orthogonal (i.e. when the structure is close to a CI as mentioned above). In the reference diagram on the left of Fig. 3.10B, the HOOP torsion oscillates for more than a period before entering the hop region (dashed vertical red line) then it enters it in a situation where the HOOP value is increasing (going to less negative values and thus displaying a positive velocity) while the C1-C2-C3-C4 torsion value is constantly decreasing (i.e. going from the E isomer reactant where the C1-C2-C3-C4 angle is nearly -180° as we are now starting from the E form of the chromophore, to the CI that has a C1-C2-C3-C4 angle close to -90°). In this situation the HOOP has a negative velocity and would hinder the formation of the Z isomer. Thus, the majority of the trajectories are expected to lead to the E isomer reactant even if a non-negligible number of trajectories may lead to product. Fig. 3.11 shows a computational demonstration of this effect for Rh where the C1-C2-C3-C4 torsional angle discussed above is now the C10-C11-C12-C13 angle of the rPSB11 chromophore and the isomerization proceeds from Z (starting close to 0°) to E.

The directionality of the non-adiabatic trajectory discussed above can be modified by changing either the frequency or phase of the HOOP coordinate relative to the torsion coordinate. These changes are schematically illustrated by the diagrams on the right of Fig. 3.10B. Changing the frequency (see central-right diagram) to a lower value allows the HOOP mode to reach the CI region while it is becoming more negative, and therefore acquires a positive velocity. According to the above rules, this has the consequence of leading to reactive trajectories. The same effect can be achieved by shifting the phase of the HOOP mode (see bottom-right diagram in Fig. 3.10B). In conclusion, one way of controlling QYs would be to "engineer" specific combinations of frequency and phase changes in the HOOP mode.

Recently research work has been carried out to explore the effect of external forces applied to a chromophore (e.g. such as in a polymer chain incorporating one photochemically reactive unit).⁷⁴ The authors report non-adiabatic trajectory simulations showing a marked and counterintuitive increased in double-bond E to Z photoisomerization quantum yield of a reduced model of the rPSBAT chromophore when pulling forces are applied axially to the terminal C and N atoms of the molecule.⁷⁵ The effect is successfully interpreted in terms of the phase relationship seen above.

3.5 Summary: Biological chromophores and their modulation.

On the basis of the rPSB results discussed in the previous sections we may conclude that Nature has implemented ultrafast double-bond isomerizations through different "selection processes". For instance, the chromophore is selected in such a

way to provide a low-lying CI (or IS segment) which could be accessed easily after having absorbed a photon of visible wavelength and could preferentially mediate the isomerization of a specific double bond (e.g. at the center of the conjugated chain). The chromophore must also provide a space-saving reaction coordinate capable to cope with spatial constraints as the isomerization may have to occur in a tight molecular cavity. Other selection processes are related to the "modulation" of the spectroscopic and photochemical properties of the chromophore. Therefore, a chromophore with a spectroscopic/reactive excited state featuring a large charge-transfer character is likely to be very sensitive to the environment electrostatics and, therefore, to the amino acid composition of the hosting protein. These modulable properties ultimately enable the optimization of the biological function including light sensitivity. In fact, above we have reviewed computational evidence showing that the charge-transfer character may be involved in the modulation of both static properties, such as excitation energies (e.g. λ_{\max}), reaction path slopes, conical intersection stability and thermal isomerization barriers, as well as dynamic properties, such as excited state lifetime and, quantum yields. Similarly, the chromophore geometrical "manipulation" (e. g. the pre-twisting of single and/or double bonds) may be involved in excitation energies (e.g. λ_{\max}) modulation as well as in the selection of the reactive double bond.

4. RETINAL PROTEINS

In Sections 2.1, 2.2 and 3.1, we have introduced Rh as a prototypical system for understanding the mechanism of double-bond isomerization in photobiology. As a result, we have extensively discussed the *cis-trans* isomerization of the C11=C12 double bond in Rh, the geometric and electronic structure of its CI, the reaction path followed during photoisomerization, and we also presented trajectory calculations simulating the isomerization of Rh. However, rhodopsins are a very large family of photoreceptor proteins found in all domains of life, where they serve a diverse array of functions.⁷⁶ In humans alone, rhodopsins are known to be responsible for color vision, dim light (scotopic) vision, and non-visual light perception for regulation of circadian rhythm through a pigment called melanopsin.⁷⁷ As also mentioned earlier, not all rhodopsins have the same photochemistry and photophysics. In this section, we will present computational studies on other types of rhodopsins, and attempt to draw a comparison between different types of rhodopsins. This section is divided into subsections focusing on conical intersections (Section 4.1), reaction paths (Section 4.2), and trajectories (Section 4.3).

4.1 Conical intersections.

In Section 3.1 we have discussed the geometrical and electronic structure of the CI of Rh as a reference vertebrate visual pigment. As shown in Fig. 4.1A, the structure and stereochemistry of the CIs driving the rPSB11 isomerization in squid rhodopsin (SqRh) and in human melanopsin (hMeOp), an invertebrate visual pigment and a vertebrate non-visual pigment, respectively, were found to be similar to the Rh CI. The comparison of these CI structures with the corresponding S_0 equi-

librium structures (for Rh see the top-right structure in Fig. 4.1A), provides information on the rPSB11 changes occurring before S_1 decay. It is shown that, in all three cases, rPSB11 undergoes the same clockwise (CW) twisting of the C11=C12 reactive double bond accompanied by a nearby counterclockwise (CCW) twisting of the C9=C10 double bond.²⁴ The sense CW and CCW rotation is defined by the motion of the forward or backward carbon atoms of the isomerizing double bond (e.g. C13 and C10 atoms in the Newman projection on the right of Fig. 4.1A).

The CI structures of animal rhodopsins can also be compared with those of microbial rhodopsins. More specifically, while the CI structures of the rPSBAT or rPSB13 chromophores of archaea and eubacteria rhodopsins^{78,79} drive a C13=C14 rather than C11=C12 isomerization, the documented S_1 structural changes are similar to the aborted bicycle-pedal coordinate driving the isomerization in animal rhodopsins. For instance, the sensory rhodopsin from the eubacterium *Anabaena* (ASR), which exists in two forms (ASR_{13C} and ASR_{AT}) featuring a 13-*cis*,15-*cis* rPSB13 and 13-*trans*,15-*trans* rPSBAT chromophore respectively,^{80,81} displays CIs characterized by a large twisting of the C13=C14 and C15=NH bonds with a minor twisting of the C11=C12 bond (see Fig. 4.1B).⁷⁹ The value of the twisting suggests that C15=NH will revert, as expected for an aborted process, to the original conformation immediately after the decay to S_0 .

In spite of their differences, the CIs of Rh and ASR appear to drive isomerization processes consistent with alternated CCW and CW twisting deformations along the conjugated chain. In fact, the stereochemistry of the ASR_{13C} and ASR_{AT} CIs can be described in terms of a C13=C14 CCW and a C15=NH CW twisting. This is consistent with the hypothesis that, in general, the C9=C10 and C13=C14 twists in a CCW direction while the C11=C12 and C15=NH twists in a CW direction. Such hypothesis is supported by the CI structure computed for the archaea light-driven proton pump bacteriorhodopsin (bR) that, in its light-adapted form (bR_{LA}), features a 13-*trans*, 15-*trans* rPSBAT chromophore. In fact (see Fig. 4.1C), the bR CI is consistent with a bicycle pedal mechanism involving a C11=C12 CW and a C13=C14 CCW twist with a minor CW twist along C15=NH. In conclusion, while the regiochemistry of the bicycle-pedal isomerization seems to be distinct for animal and microbial rhodopsins, the direction of the twisting motion is found to be consistent at least for the few CIs currently investigated.

4.2 Reaction paths.

4.2.1 Photochemical reaction paths in animal and microbial rhodopsins. The MEP driving the photoisomerization of the rPSB11 or rPSBAT chromophores have been computed for both gas-phase reduced models of the chromophore as well as for QM/MM models of different rhodopsins. We will begin by examining important features of the MEP profile for the photochemical conversion of the visual photocycle intermediate bathoRh to Rh. This back-isomerization process has been studied experimentally⁸² (and recently by using time-resolved spectroscopy^{83,84}) as well as computationally (see for instance ref. ⁸⁵), because it is one of the fundamental processes that could limit the sensitivity of a photoreceptor. This has been illustrated in Scheme 3.1 by a dashed vertical arrow represent-

ing a photon absorption from the photoproduct. Its detailed investigation is therefore of interest. What we report below is only a partial account of the work focusing on reaction paths.

The analysis of the S_1 MEP of this process (see Fig. 4.2A), provides a static description of the photoisomerization mechanism. It also provides a basis for interpreting the results of non-adiabatic trajectory calculations (see Section 4.3) in terms of forces (the slopes of the S_1 to S_0 potential energy surfaces) driving the chromophore changes. These forces control the progression of the S_1 population towards the CI and, after decay, its relaxation towards either the primary S_0 photoproduct or the original reactant.⁸⁵




Figure 4.1. Rhodopsin conical intersection structures. A. Superposition of the CI structures computed for a vertebrate visual pigment (bovine rhodopsins, Rh), an invertebrate visual pigment (squid rhodopsin, SqRh) and for a human non-visual pigment (human melanopsin, hMeOp). The Rh S_0 equilibrium structure is also shown in the top-right corner for reference. Adapted partly from ref.²⁴. Copyright 2014 National Academy of Sciences. B. Comparison between the photoisomerization stereochemistry of Rh and the eubacterial *Anabaena* sensory rhodopsin (ASR). According to the corresponding CI structures, the aborted bicycle pedal isomerization occurs along the C9=C10-C11=C12 moiety in Rh and the C13=C14-C15=NH in ASR. Geometric data from ref.²³. C. Schematic representation of the CI structure of the light-adapted form of bacteriorhodopsin (bR) indicated as bR_{LA}. The structure is consistent with an aborted bicycle pedal motion along the C13=C14-C15=NH moiety. Geometric data from ref.⁸⁶.

Consistently with the data presented in the right part of Fig. 4.2A, the MEP starting from bathoRh initially develops along the central C11=C12 double bond turning it into a stretched C11-C12 single bond. As anticipated above, this motion is part of a more complex BLA stretching coordinate. Such relaxation leads rapidly to a mixed coordinate incorporating a HOOP wagging deformation of the two hydrogen atoms linked to the carbons of the reactive double bond.²⁷ As a reminder, HOOP is defined as the difference between the angle β , corresponding to the H-C11-C12-H dihedral angle, and the angle α corre-

sponding to the skeletal C10-C11-C12-C13 dihedral. Due to the S_1 forces initially acting along the C11=C12 double bond and H-C11-C12-H dihedral, the HOOP wagging changes more rapidly than the skeletal C11=C12 twisting, reaching its maxima at a point where the skeletal deformation is minimal. This rapid change along the MEP is explained in terms of the S_1 anti-bonding character between the two π -systems overlapping through the C11-C12 bond. Such a character is caused by the Φ_{CT} -dominated S_1 electronic structure which, effectively, shifts the spin-pairing of the electrons of one position along the chromophore's conjugated chain. Consequently, a force acting on the nuclei develops along the critical BLA stretching, HOOP wagging and C11=C12 twisting modes.²⁷ As reported above (Fig. 2.3, Fig 3.9 and related text), the fastest evolving coordinates are the ones that more rapidly decrease the overlap: the BLA (dominated by the C11=C12 expansion) that increases the distance between the centers of the two π -systems and HOOP (dominated by the change in the angle β) that rapidly increases the angle between the orbital centers. In conclusion, while, as evident from Fig. 4.2A the S_1 potential energy constantly decreases along the MEP coordinate of bathoRh, the coordinate itself involved several different modes and is therefore highly curved.




Figure 4.2. Structure of the MEP driving the photoisomerization of different rhodopsins. A. Photoisomerization of bathoRh to Rh. This intermediate has a largely distorted rPSB11 chromophore that reverts to rPSB11 regenerating Rh. The left part displays the energy profiles with structures illustrating the main geometrical changes along the MEP. It also provides a definition of the dihedral angles α and β , which define the HOOP and C10-C11-C12-C13 twisting coordinates. The right part describes the coupling of the C10-C11-C12-C13 dihedral angle of the isomerizing double bond with the HOOP angle and C11-C12 length (notice that the x-axis here is the torsion and not MEP coordinate. However, since the C10-C11-C12-C13 twisting is monotonically increasing along the MEP, the diagram also loosely represents the HOOP and stretching geometrical progression along the MEP from reactant to the bathoRh CI). The relationship between the progression along the MEP and the overlap between the π -systems (represented by hybrid p-orbitals) describing the bond breaking/making of

the C11=C12 bond is also given. Adapted from ref. ²⁷. Copyright 2011 American Chemical Society B. Approximated MEP computed in terms of a relaxed scan along the C12-C13-C14-C15 dihedral angle describing the photoisomerization of the chromophores of the eubacterial ASR_{AT} (left) and ASR_{13C} (right). C. Light-adapted (left) and dark-adapted (right) equilibrium structures of the chromophores of the archaea proton pump bR.

The energy profiles along the S₁ MEP computed for vertebrate, invertebrate, eubacteria and archaea rhodopsins are all substantially barrierless.^{21,24,79,86} However, the MEP computed for ASR_{AT} and bR_{LA} and featuring a rPSBAT chromophore show a flat S₁ energy profile immediately following BLA relaxation.^{79,86} This flat region has been proposed to be responsible for the observed longer S₁ lifetimes and photoproduct appearance times with respect to rhodopsins hosting either the rPSB11 or rPSB13 chromophore.^{79,86} A comparison between the energy profiles computed along an approximated MEP (i.e. a relaxed scan) is given in Fig. 4.2B for the case of ASR_{AT} and ASR_{13C}.

The analysis of the S₁ MEPs described above suggests that the stereochemistry of the S₁ relaxation of vertebrate, invertebrate, eubacteria, and archaea rhodopsins is the consequence of a small twisting of the initial S₀ equilibrium structures.^{21,24,79,86} Such twisting would be responsible for initiating the relaxation in specific CW or CCW directions, ultimately leading to a specific CI. For instance, as shown in Fig. 4.1A, the S₀ equilibrium geometry of Rh includes a rPSB11 with the C11=C12 bond twisted by ca. 6° in the CW direction. Such a small twist in the chromophore S₀ equilibrium structure is also observed for SqRh and hMeOp, which both display a 10° twist around the C11=C12 bond.²⁴ Similarly, the microbial rhodopsins ASR_{AT}, ASR_{13C}, and bR_{LA}, show an S₀ equilibrium structure with a C13=C14 twist of 8°-11° in the CCW direction (see Fig. 4.2B and 4.2C), consistent with the result of the MEP that shows a CCW isomerization coordinate.²³

The S₁ MEPs computed for ASR_{AT} and ASR_{13C} both describe a twisting of the reactive C13=C14 bond occurring in the CCW direction. When considering that ASR is photochromic (see Fig. 4.3A) such that ASR_{AT} and ASR_{13C} are interconverted via irradiation at two different wavelengths,^{80,81} Strambi et al.⁷⁹ hypothesized that the ASR chromophore operates like a light-driven single-molecule unidirectional rotor.⁸⁷ The corresponding rotation mechanism has been documented by following the S₁ MEPs reported in Fig. 4.2B and by computing the K and L photocycle intermediates of both the ASR_{AT} and ASR_{13C} forms.⁷⁹ The K and L structures are the result of MEPs that describe the S₀ relaxation from the CIs of the ASR_{13C} and ASR_{AT} towards the ASR_{AT} and ASR_{13C} photoproducts, respectively. The mechanistic picture emerging from such calculations is shown in Fig. 4.3B, where the stereochemistry of the relevant double-bond regions is assigned using both the M and P helicity labels and the CW and CCW labels describe the twisting direction associated with either light or thermally induced isomerizations.




Figure 4.3. Unidirectional photochromic interconversion mechanism of ASR derived from calculations. A. The intertwined photocycles of ASR_{AT} and ASR_{13C} as represented by the C12-C13-C14-C15-NH-Cε chromophore moiety in different photocycle intermediates. The wavelength values refer to observed⁸⁰ and computed (in bold) absorption maxima of each intermediate. In each state, the isomerizing bond is highlighted in color. B. The photochromic cycle of ASR. Solid-line curved arrows indicate alternating CCW and CW isomerizations about the C13=C14 and C15=N double bonds, respectively. The helicities of the C15-NH-Cε and C12-C13-C14 moieties (dashed curved arrows) are assigned by determining whether the helicity of the corresponding H(C14)-C15-NH-Cε and C12-C13-C14-(C15)H fragments is M or P. Adapted with permission from ref. ⁷⁹. Copyright 2010 National Academy of Sciences.

The rPSBAT chromophore was studied in the chimeric channelrhodopsin C1C2 more recently by two different research groups.^{88,89} Both studies were based on relaxed scan calculations using the CASPT2//CASSCF methodology and reached similar conclusions. Consistently, two pathways were determined with opposite rotating C13=C14 bonds. One pathway was found to have a low or no barrier in the excited state, while the other one was showing a larger barrier explaining two different time constants for excited-state lifetimes from transient absorption measurements.⁸⁹

4.2.2 Relationship between thermochemical and photochemical isomerization paths in rhodopsins. As discussed in Section A.I.1, the S₀ potential energy surface of rPSB11 may host two transition states (TS_{DIR} and TS_{CT}) controlling the same thermal isomerization. This has also been demonstrated for gas-phase reduced models of rPSB11.^{43,90} Fig. 4.4 presents the TS_{DIR} and TS_{CT} structures computed in a bovine Rh QM/MM model.⁵⁶ TS_{DIR} features the structure expected for a homolytically broken double bond, with two radical centers

delocalized along orthogonal π -systems. Such a transition state was also located by Nemukhin and co-workers in a model of Rh employing DFT to describe the chromophore.⁹¹ The TS_{DIR} charge distribution correlates with that of the S_0 rPSB11 reactant with a +0.98e charge localized in the Schiff base region. In contrast, TS_{CT} has most of its charge (+0.90e) located on the β -ionone region, thus more closely resembling the charge distribution of the vertically-excited S_1 rPSB11 (compare the corresponding bubble diagrams in Fig. 3.5B). It was found that TS_{CT} has a computed activation energy of 34 kcal/mol and lies 11 kcal/mol in energy below TS_{DIR} . While these energies depend on the level employed in the calculation, an increase in the level of theory invariably leads to a larger stabilization of TS_{CT} . Therefore, it has been found that TS_{CT} has full control of the thermal isomerization.⁵⁶

To understand why both TS_{CT} and TS_{DIR} exist in Rh, and how a single reaction could be mediated by two different transition states, we need to revisit concepts from Section 3.2 where we discuss the effect of the conical intersection on the topography of the ground state. There, we have discussed how the S_0 and S_1 states change their electronic character along a small loop centered at the CI and residing on the BP. In other words, in rPSB11, the S_0 electronic character passes two times (e.g. roughly every 180°) from a Φ_{DIR} character to a Φ_{CT} character. Due to the peaked topography of the CI and the fact that TS_{CT} and TS_{DIR} lie on opposite sides of the CI along the BLA coordinate, this means that TS_{CT} and TS_{DIR} are dominated by different electronic characters, Φ_{CT} and Φ_{DIR} , one associated with the S_1 state at the FC point and one associated with the S_0 state at the FC point, respectively. Therefore, *the existence of the low-lying CI in Rh allows the existence of a transition state, TS_{CT} , with the same electronic character as the first excited state of Rh.* Such a transition state would not exist if it were not for the existence of the nearby CI.




Figure 4.4. Comparison of the structures at the S_0 transition states TS_{DIR} and TS_{CT} and at the CI for Rh. The approximated CASSCF/6-31G*/AMBER level of theory has been employed to compute the transition state geometry. Energy differences and barriers were computed at the CASPT2/CASSCF/6-31G* level. The electronic structures are represented by the corresponding Lewis structure (only the N-containing moiety is represented). The brown region corresponds to the region dominated by

the Φ_{CT} electronic configuration. Adapted with permission from ref. 56. Copyright 2012 The American Association for the Advancement of Science.

How does isomerization through TS_{CT} occur? In Fig. 4.5A we display a graphical representation of the change in π -electron density along the thermal isomerization path for Rh when passing through the lower energy transition state TS_{CT} . It is apparent from inspection of the structure that along the longest path segment leading to TS_{CT} , C11=C12 remains in a “locked” Φ_{DIR} dominated state featuring two π -electrons coupled across the breaking bond. However, due to the electronic nature of the region surrounding TS_{CT} , the electronic structure suddenly changes to a Φ_{CT} configuration as it is evident by the gap in π -electron density at C11=C12. This leads to unlocking of the C11=C12 double bond, similar to how it is unlocked in the S_1 state of Rh prompting photoisomerization. Of course, once the molecule passes the TS_{CT} moving towards bathoRh, the electronic structure acquires its original bonded electronic character allowing for the formation of the C11=C12 double bond in a new all-*trans* configuration.

The sudden change in the electronic character is quantitatively described in Fig. 4.5B. The positive charge on the β -ionone-containing moiety of rPSB11 is shown in terms of an area diagram. It is evident that in the TS_{CT} region, the positive charge on such a moiety suddenly increases and then decreases after the transition state. The same analysis demonstrates that the alternative and higher-energy isomerization path passing through TS_{DIR} displays a change in charge distribution substantially opposite to the one seen for TS_{CT} .

The nature of the transition state controlling thermal isomerization could have important implications for visual sensitivity. The similarity between the Φ_{CT} -dominated electronic structure of FC and TS_{CT} , together with the results indicating that TS_{CT} is controlling the thermal isomerization, provides a direct link between the rate of thermal isomerization (controlled by the energy of TS_{CT}) and the absorption wavelength (controlled by the FC S_1 energy). This link is illustrated in Fig. 4.6A. The diagram (see the brown color position along the curves indicating a dominating Φ_{CT} character) explains the link between E_a^{T} and $\Delta E(S_1-S_0)$ and, in turn, between $-\log k$ (where k indicates the kinetic constant of the thermal isomerization reaction) and $1/\lambda_{\text{max}}$. Accordingly, any opsin red-shifting the absorption (i.e. decreasing the $\Delta E(S_1-S_0)$ value) would simultaneously decrease E_a^{T} . This behavior, deduced from quantum mechanical calculations, appears to be consistent with the “Barlow correlation,” a prediction made by Barlow in the 1950s^{56,92} and observed in a series of experiments⁹³⁻⁹⁵ that demonstrate that indeed a relationship between the λ_{max} value and the thermal activation kinetic constant (k) in different visual pigments exists. The results above provide a basis for the development of a pigment-noise theory. Such a theory would hold that, in a series of related pigments (e.g. the rod visual pigments) the thermal noise (i.e. the thermal isomerization competing with the photochemical isomerization) would decrease when passing from red-light perceiving pigments to blue-light perceiving pigments. In other words, higher visual sensitivity may be achieved by pigments that absorb in the blue. This could be one of the reasons why dim-light vision is tuned to the blue edge of the visible spectrum.⁵⁶

The computational results reported above suggest that a Barlow-like relationship can be reproduced by QM/MM models of rhodopsins. This is illustrated schematically in Fig. 4.6B where a negative charge represents the hypothetical effect protein environment in two different visual pigments carrying the rPSB11 chromophore and absorbing light of shorter and longer wavelength, respectively. The bottom-left panel shows that the negative charge is distant from the β -ionone-ring moiety and therefore the S_1 FC and the S_0 TS_{CT} structure are only slightly stabilized with respect to the surrounding regions of the S_1 and S_0 potential energy surface, respectively. On the other hand, the top-left panel shows a situation where the negative charge is closer to the same β -ionone-ring moiety. In this situation the S_1 FC structure will be stabilized with respect to S_0 , leading to a red-shifted absorption maximum. At the same time, TS_{CT} will also be stabilized leading to a lower barrier and faster thermal isomerization rate. By interpreting the change in the position of the negative charge in terms of a change in the chromophore cavity, one achieves a molecular-level understanding of the Barlow relationship. Of course, a behavior not consistent with the Barlow correlation, would be observed if the isomerization was controlled by TS_{DIR} .

A number of computational results have been reported that support the mechanism reviewed above. In Fig. 4.7A the $-\log k$ vs. $1/\lambda_{max}$ correlation has been computed by modeling a set of Rh mutants incorporating two different types of chromophores. These comprise 5 bovine Rh mutants (hosting rPSB11, sometimes referred to as the A1 chromophore) and 6 derivatives featuring the longer wavelength absorbing 3,4-dehydroretinal (A2) chromophore. The calculated correlation can be compared directly to experimental data collected by Ala-Laurila et al.⁹⁴




Figure 4.5. Evolution of the electronic structure along TS trajectories in Rh. A. Schematic representation of the S_0 potential energy surface driving the isomerization of the protein embedded rPSB11 chromophore and of the progression from Rh to bathoRh along a TS trajectory. The three rPSB11 structures display the calculated shape of the π -electron density along the conjugated framework for the trajectory passing through TS_{CT} . B. Change in S_1 and S_0 potential energy (red and blue lines, respectively) and charge distribution (orange area) along the two TS trajectories passing through TS_{CT} and TS_{DIR} . Each TS trajectory is a composite trajectory defined by releasing two classical trajectories from the optimized TS with minimal velocities towards the reactant and product, respectively, but then reversing the reactant trajectory to generate the plot shown.

A correlation of the same type has been obtained by looking at the computed E_a^T and $1/\lambda_{max}$ values for a set of more distant (in the sense of the sequence diversity) set of pigments with respect to Rh mutants. The set comprises four visual pigments from cottoid fish from lake Baikal (see Fig. 4.7B), which are also compared with bovine Rh. In this case the apparent linear correlation between the two quantities has also been analyzed in terms of two effects. The first is related to a modulation of the chromophore geometry, while the second is a pure modulation of the protein electrostatic field acting on the chromophore. It was demonstrated that the modulation in the electrostatic field is mainly responsible for the correlation between E_a^T and $1/\lambda_{max}$. The validation of a Barlow-type correlation in the lake Baikal fish flock provides an explanation for the fact that the species with the most blue-shifted absorption reside at the abyssal zone of the lake (400-1500 m depth) where the quantity of available light is at its minimum. In this situation, the visual pigment must have a high thermal isomerization barrier to limit the thermal noise. On the other hand, the species thriving in the maximally illuminated littoral zone of the lake (1-5 m depth) have red-shifted absorption maxima.

Fig. 4.7C shows that a Barlow-type correlation may also exist for distant rhodopsins that do not even serve the same biological function. In fact, using the same type QM/MM model, it has been demonstrated that the computed E_a^T and $1/\lambda_{max}$ values for a small set of vertebrate, invertebrate and microbial rhodopsins may also be directly proportional, again having the most red-absorbing member of the set featuring the lowest thermal isomerization barrier. In conclusion, the existence of a Barlow correlation (i.e., a link between absorption wavelength and thermal isomerization rate) appears to be present in all types of rhodopsins.




Figure 4.6. Relationship between $\Delta E(S_1-S_0)$ and E_a^T in rhodopsins. A. One-dimensional representation of the relationship between the energy profiles and electronic structure of the S_1 and S_0 isomerization paths when TSCT is controlling the S_0 process. The brown segments along the paths are dominated by a Φ_{CT} configuration. The green segments are instead dominated by a Φ_{DIR} (diradical or covalent) configuration. B. Schematic representation of the simultaneous effect of the protein environment on the $\Delta E(S_1-S_0)$ and E_a^T values. The negative charge in the “red” and “blue” panels schematically represents the electrostatic effect of the cavity hosting the chromophore.




Figure 4.7. Barlow-like correlation in rod visual rhodopsins. A. Simulation of the $-\log(k)$ vs. $1/\lambda_{max}$ relationship. By using a set of TSCT E_a^T values computed at the CASPT2//CASSCF/6-31G* level of theory and the same pre-exponential factor used in Luo et al.⁹⁵ it has been possible to simulate the Barlow relation for 11 rod pigments using the $\log k = \log A(E_a^T, T) - E_a^T/RT$ expression (full circles). The computed relation is compared with experimental data (crosses), which are collected in ref.⁹⁴. The figure is adapted with permission from ref.⁵⁶. Copyright 2012 The American Association for the Advancement of Science. B. Computed thermal isomerization barrier E_a^T plotted as a function of $1/\lambda_{max}$ for Cottoid fish rod pigments with the rPSB11 chromophores in the protein (black circles), in the gas phase (gray triangles) and in the protein with no point charges (red squares). Straight lines indicate the ideal linear relationship. Adapted with permission from ref.⁹⁶. Copyright 2016 Nature Publishing Group. C. Computed values of the EaT barriers controlling the thermal isomerization of hMeOp, sqRh, Rh (featuring the rPSB11 and rPSB_{13C} chromophore, respectively), and the two forms of the eubacterial rhodopsin ASR_{AT} and ASR_{13C} plotted as a function of $1/\lambda_{max}$. Adapted with permission from ref.²³. Copyright 2015 National Academy of Sciences.

4.3 Trajectories.

4.3.1 Rhodopsin. The first simulation of the photoisomerization of the dim-light visual pigment Rh was reported in 1976 by Warshel.²² This pioneering effort introduced the space-saving reaction coordinate hypothesis. The simulated trajectory pointed to a rPSB11 isomerization mechanism dubbed bicycle-pedal (BCP in Fig. 4.8A) because of the simultaneous twisting of two double bonds resembling the motion of the pedals of a bicycle. The mechanism originally proposed by

Warshel involved isomerizations at C11=C12 and C15=NH and therefore is distinct from the version described in the introduction based on more recent QM/MM models. In fact, recent studies point to a BCP at C11=C12 and C9=C10. Furthermore, it is aborted upon decay at the CI yielding a so called non-reactive or aborted BCP (NBCP) where only the C11=C12 completes the isomerization, but not C9=C10.

In a seminal study, Warshel employed an empirical force field method (QCFF/PI)^{97,98} that provided the S_1 and S_0 gradients necessary to propagate a non-adiabatic trajectory starting on S_1 . At that time, a crystallographic structure of rhodopsin was not available, and therefore the steric interactions with the protein matrix were modeled by imposing geometric constraints on the chromophore atoms. It is remarkable to note that Warshel has predicted this reaction to complete in 200 fs before such an ultrafast timescale became accessible experimentally. Other trajectory calculations by Birge and Hubbard were favoring a one bond flip (OBF) isomerization at C11=C12.⁹⁹ They predicted also a longer excited-state lifetime of ca. 2.0 ps. Using an improved protein-environment model Warshel and co-workers refined their isomerization model.¹⁰⁰ The original mechanism was modified by Warshel et al. such that only one double bond, C11=C12, isomerized successfully and a second one, C9=C10, becomes significantly twisted but then reverts back to the original configuration. This is essentially the NBCP mechanism documented by modern QM/MM models based on multiconfigurational quantum chemical methods.

In course of development of quantum chemical methods and improvement of the computational hardware more accurate molecular dynamics simulations become accessible. However, even more important was the availability of the crystal structures as a starting point for trajectory calculations. The trajectory calculations are the closest simulations to experiment because it considers the effect of kinetic energy in contrast to MEP calculations. Further, molecular dynamics simulations are free of bias and do not need *a priori* knowledge of the reaction mechanism as the outcome is determined by the initial conditions. Hence, with the advance of QM/MM trajectories various proposed mechanisms of the photoisomerization could be tested (Fig. 4.8). Such mechanisms evolved from chemical intuition or static quantum chemical calculations, e.g. Liu et al. suggested a volume-conserving isomerization of neighboring single and double bonds in the so called “hula-twist” (HT in Fig. 4.8A) mechanism.¹⁰¹⁻¹⁰³ More recently, a “double bicycle-pedal”⁸⁶ or “folding-table”¹⁰⁴ (FT in Fig. 4.8A) mechanism was proposed for bR.

The mechanisms above have mainly been invoked to explain the ability of the rPSB chromophore to isomerize in a space-saving way inside a tight binding pocket that restricts large-amplitude motions of the embedded chromophore. However, in addition to the steric consideration, electronic factors also play an important role in determining the mechanism of isomerization. The physical origin of the preferred excited-state BCP-like relaxation is schematically illustrated in Fig. 4.8B, where it is shown that such a mechanism is driven by the Φ_{CT} electronic structure of the excited-state chromophore. In the same figure we also show how this mechanism is different for

polyenes where the isomerization is driven by an excited state with diradical Φ_{XDIR} electronic structure (see Section 3.1).




Figure 4.8. Overview of various isomerization mechanisms. A. one bond flip (OBF), bicycle pedal (BCP), non-reactive or aborted bicycle pedal (NBCP), hula twist (HT) and folding table (FT). B. Relationship between electronic structure and facile motion in a charge-transfer excited state (CT) of an rPSB and in an excited diradical state (XDIR) of a polyene. See also Section 3.1.

Rh was the major focus of trajectory calculation for several reasons: (i) the crystal structure became available in 2000,⁴¹ and (ii) shortest excited state lifetime leading to short computational times. The exact details of the computational methodology varied in the studies in several aspects. A broad range of the quantum chemical methods was used, starting from restricted open-shell Kohn-Sham (ROKS) method³⁰ to wave function based methods. The latter methods were the preferred choice and included semi-empirical floating occupation molecular orbital configuration interaction (FOMO-CI)^{34,105} but used mainly CASSCF^{21,26,27,29,32,106}. To compensate for the lack of dynamic electron correlation in CASSCF, Frutos et al. have scaled the forces linearly to qualitatively match those obtained at the CASPT2 level.²¹ This approximation was supported by the finding that only the steepness of the excited-state potential was different between the energy profiles described at the CASSCF and CASPT2 levels of theory. More recently, Martinez and coworkers have introduced an empirical correction in a so-called α -CASSCF approach that scales the splitting between state-specific and state-averaged CASSCF.¹⁰⁷ The correction was determined to match the multistate CASPT2 reference energy and gradients.

Despite the differences in the computational setup the majority of the trajectories were found to reach an S_0/S_1 degeneracy on the scale of 100 fs,^{21,27,29,32} and to form predominantly the bathoRh intermediate. The reaction coordinate was composed of the sequential inversion of BLA and torsion of the C11=C12 bond as already determined in MEP or relaxed scan calculations discussed in section 4.2. However, also the 9-*cis* isomer (isoRh) and a genuine BCP product were found as by-products in some trajectories^{26,29} which can only be observed in molecular dynamics simulations. Another new insight gained from trajectory simulations was demonstrated by Ishida and coworkers¹⁰⁶ on Rh and IsoRh. The results showed vibrational coherence only in the case of Rh but not for isoRh. Further, the authors explained the longer excited-state lifetime of isoRh with respect to Rh. In particular, it was found that the

former stays longer on the excited state with respect to Rh because of back-and-forth twisting of the dihedral angle in S_1 , while in Rh this doesn't occur (see Fig. 4.9). Two amino acids were found to be responsible for this extended S_1 lifetime: Tyr268 and Thr118. Furthermore, the authors proposed a new mechanism: the "wring-a-wet-towel" motion. However, since this mechanism is defined by rotation of C9-C10 and C11-C12 in opposite directions it appears to be a variant of the BCP mechanism.




Figure 4.9. Change in length of the reactive bond with respect to the corresponding dihedral angle along: A. Rh trajectories and B. isoRh trajectories leading to the same bathoRh intermediate. Full triangles indicate hop points. Adapted from ref. ¹⁰⁶. Copyright 2012 American Chemical Society.

Further insight into the isomerization in IsoRh was provided by Garavelli and coworkers.⁴⁰ Their QM/MM molecular dynamics simulations have located two deactivation pathways starting from rPSB9. One path was leading to a CI that produced a di-*cis* photoproduct while the other path reached a different CI that can generate the rPSB11 and rPSBAT photoproducts. The latter pathway was found to have an increased interaction with Tyr268, in line with the work of Ishida and coworkers,¹⁰⁶ and therefore a longer lifetime on the excited state than the former pathway. Hence, the authors interpreted the presence of two different isomerization directions to be responsible for the biexponential decay of IsoRh.

4.3.2 Bacteriorhodopsin and other microbial rhodopsins.

The rPSBAT chromophore in bR and microbial rhodopsins in general, has a longer excited state lifetime than rPSB11 that is thought to originate in the presence of an excited state barrier. The longer simulation time and the accurate assessment of the barrier present a challenge for accurate excited state dynamics simulations. The first trajectories were computed by Birge and coworkers using semiempirical molecular dynamics.¹⁰⁸ Their calculations of the rPSBAT to rPSB13C isomerization in bR were performed at a time before full structural data for bR

became available. They predicted that a transoid minimum on S_1 may trap some of the excited-state population before decay. The first high-resolution (1.55 Å) crystal structure of bR became available in 1999, thus providing a more realistic starting point for simulations. In 2001 Warshel and Chu¹⁰⁹ used a QM/MM setup with the Empirical Valence Bond method¹¹⁰ to describe the S_1 potential of the chromophore in the QM subsystem. In their simulations, they also accounted for the effect of polarization of the protein environment by the QM atoms.

The first simulation of rPSBAT in bR carried out with a QM/MM model based on *ab initio* multiconfigurational quantum chemistry was reported two years later by Schulten and co-workers.¹¹¹ The QM subsystem in their calculations consisted of a reduced model, a three double-bond portion of the retinal from C11 to C ϵ , and described at the CASSCF level of theory. Consequently 6 π -electrons in 6 π -orbitals were included in the active space. The remaining atoms of the retinal chromophore, together with the protein, were calculated with a classical empirical force field.

Morokuma and coworkers¹¹² studied bR and rPSBAT in methanol using the ONIOM scheme for QM/MM molecular dynamics and CASSCF as an electronic structure method. The trajectories were computed in bR as well as in methanol that served a reference system. The NBCP mechanism was found in both environments. A different comparison of the environmental effect on rPSBAT photoisomerization was carried out by Punwong et al.¹¹³ who also studied the gas-phase reaction. The QM subsystem was described at the semi-empirical FOMO-CI¹⁰⁵ level of theory. The isomerization in methanol was found to be much longer than in the gas-phase. The reason for this longer time is an S_1 barrier which slows down the isomerization. They found that the electrostatic (not the steric) interactions with methanol are the primary cause of this torsional barrier. More specifically, the barrier originates from the solvent being oriented to stabilize the Franck-Condon region of the chromophore, and therefore isomerization requires the solvent to reorient as the charge distribution changes during torsion on the excited state, slowing down photoisomerization compared to the isolated rPSBAT chromophore. Punwong et al. also discuss the photoisomerization quantum yield in the gas-phase and solution. They agreed with the hypothesis of Martínez and co-workers¹¹⁴ that the CI topography has an important role in determining the branching ratio of product vs. reactant formation (i.e., the photoreaction quantum yield).¹¹³

In order to compare to different microbial rhodopsins Punwong et al.¹¹⁵ simulated the excited-state dynamics of bR and halorhodopsin (hR) using the same QM/MM model as mentioned above.¹¹³ The authors found that bR has much shorter excited-state lifetime than hR. The different isomerization rates was attributed to different electrostatic stabilization of the positive charge of rPSBAT on S_1 in hR and bR.¹¹⁵

4.3.3 Comparative studies of animal and microbial rhodopsins.

A comparison of the S_1 dynamics of Rh and isoRh has been reported by Ishida and coworkers¹⁰⁶ and, as seen in Section 3.3 and 4.3.1, by Garavelli and coworkers.⁴⁰ In both cases the slower isomerization of isoRh with respect to Rh, has been successfully simulated. Their results indicate that the

steric hindrance imposed by the environment on the C9=C10 bond of rPSB9, but not on the C11=C12 double bond of rPSB11, is the cause for the different isoRh and Rh dynamics. Since isoRh and Rh have the same protein cavity, these findings suggest that the selection of a specific rPSB chromophore isomer is an important factor for the modulation of the rhodopsin function.

While the rPSB9 chromophore is mainly known as a byproduct of the in PSB11 photoisomerization in Rh, distinct rPSB chromophore isomers characterize the dark states of microbial and animal rhodopsins (also called type I and type II rhodopsins).¹ In fact, type I and type II rhodopsins are characterized by the rPSBAT and rPSB11 chromophore, respectively. Both chromophores are responsible for the activation of the protein function that is triggered by the photoisomerization of the C13=C14 and C11=C12 double bonds. However, the fact that rPSB11 is thermodynamically unstable with respect to rPSBAT indicates that rPSB11 might have been selected by biological evolution to carry out an otherwise inaccessible function. To investigate whether such function exist, a recent study has compared the photoisomerization mechanisms of rPSBAT and rPSB13 in the microbial rhodopsin ASR (see above and Section 4.2.1) and of rPSB11 in Rh using trajectory computations.²³ The simulations unveil electronic effects achievable in rPSB11 but not in rPSBAT that may explain the higher photoisomerization speed reported for animal rhodopsins.

Consistently with previous studies,²¹ it was assumed that during the 200 fs simulation time a FC trajectory (see Section 3.2) describes the average evolution of the corresponding S_1 population. This has been partially assessed by running small sets of trajectories starting with different initial conditions and comparing them with the single FC trajectory^{21,24,32}. As shown in Fig. 4.10A the ASR_{AT} trajectory remains on S_1 for the entire duration of the trajectory whereas ASR_{13C} reaches the CI after ca. 150 fs. The estimated S_1 lifetimes are consistent with the available transient spectroscopy measurements¹¹⁶ and computations¹¹⁷ showing that the ASR_{AT} photoisomerizes at least five times more slowly than ASR_{13C}. Moreover, ASR_{13C} (see Fig. 4.10B) seems to react in a time-scale that is only slightly slower than that computed for Rh^{21,24} (Fig. 4.10C), consistently with the observations.^{32,70}




Figure 4.10. QM/MM trajectories of ASR_{AT}, ASR_{13C}, and Rh computed at the two root state average scaled-CASSCF/Amber (black lines) level of theory and corrected at the CASPT2²¹ and XMCQDPT2 levels.¹¹⁸ A. S_0 (filled diamonds), S_1 (filled triangles), and S_2 (filled circles) CASPT2//CASSCF/Amber energy profiles along the ASR_{AT} trajectory.


The main out-of-plane (deviation larger than $\pm 5^\circ$) dihedral angles of the chromophore S_0 equilibrium structure are given at the top. The vertical dashed arrow represents weak fluorescence. B. Same data for ASR_{13C} with representation (at Top) of the bicycle pedal motion driving the isomerization on S_1 in which the bond order is reversed. The vertical dashed arrow represents weak fluorescence. C. Same data for the Rh trajectory. D. View of the XMCQDPT2 (solid line) and CASPT2 (dotted line) S_1 and S_2 energy profiles (Upper) and oscillator strengths (Lower) along the ASR_{AT} trajectory. Adapted with permission from ref. ²³. Copyright 2015 National Academy of Sciences.

The slower photoisomerization of ASR_{AT} is also consistent with its observed larger fluorescence quantum yield.⁷⁰ In fact, starting 25 fs after excitation, the authors compute an average oscillator strength in the 0.8-1.6 range (see Fig. 4.9D). Taking the average $\Delta E_{S_1-S_0}$ values over the 175 fs interval, they predict a fluorescence λ_{\max} of 669 nm computed λ_{\max} values are comparable with the observed values (i.e., a λ_{\max} of 700-710 nm with a shoulder around 650 nm).⁷⁰

It is apparent from inspection of the energy profiles in Fig. 4.10A and 4.10B that both S_1 and S_2 are involved in the isomerizations of ASR_{AT} and, partially, of ASR_{13C}. Indeed, the S_1 and S_2 states become nearly degenerate ca. 25 fs after photoexcitation. However, while ASR_{AT} travels along the degeneracy for the full duration of the trajectory, ASR_{13C} abandons the degeneracy region ca. 50 fs after entering it and then evolves toward the CI. In contrast, no degeneracy is found for Rh, which evolves toward the CI directly after photoexcitation. A similar behavior is found for other animal rhodopsins that feature the rPSB11 as their chromophore (e.g. SqRh and hMeOp). These data support a relationship between the mag-

nitude of the estimated S_1 lifetime (defined as the time it takes to reach the CI, τ) and the extent of the S_2/S_1 mixing. Inclusion of more excited states in the calculations does not change this conclusion.

The τ values of ASR, together with those previously reported for Rh, sqRt, and hMeOp²⁴ provide information on a dynamic factor that may be implicated in the rhodopsin function (e.g. light sensitivity). Whereas an increase of τ may be associated with a decrease in slope of the S_1 potential energy surface accelerating progression toward the CI, in ASR_{AT} and other rPSBAT-hosting rhodopsins the occurrence of the S_2/S_1 degeneracy changes the dynamics. The S_1 relaxation would thus become slower and more diffusive with the chromophore exploring the rugged potential energy surface shaped by several S_2/S_1 avoided crossings (see inset in Fig. 4.1B).



Scheme 4.1. Comparison of isomerization paths in rPSB11-based and rPSBAT-based rhodopsins. A. Schematic representation of the photochemical (solid arrows) and thermal (dashed arrows) isomerization paths for rPSB11 in Rh. Adapted with permission from ref. ²³. Copyright 2015 National Academy of Sciences. B. The same representation for rPSBAT in ASR_{AT}. The vertical dashed arrow indicates weak fluorescence. Adapted with permission from ref. ²³. Copyright 2015 National Academy of Sciences. In the inset we display C. the previously reported three-state model for the PSBAT-hosting Archaeal bacteriorhodopsins (bR).¹¹⁹

The results above indicate that whereas a two-state isomerization model (Scheme 4.1A) applies to fast (<150 fs) reacting systems incorporating the rPSB11 or even rPSB13 chromophores, a three-state model (Scheme 4.1B) would apply to slower rhodopsins hosting rPSBAT. A three-state model has been previously reported¹²⁰ for the rPSBAT-hosting Archaeal bacteriorhodopsins (bR), which displays a slow and complex S_1 dynamics.¹²⁰ However, in such early three-state models, the S_2 diradical state ultimately drives the progression toward the CI (Fig. 4C, Inset). Although this seems to be the case in polyenes,¹²¹ it is inconsistent with the structure of protonated Schiff base CIs.¹²² Instead, the three-state model based on the more recent work of Luk et al.²³ involves the S_2 state only along part of the isomerization path, where it dips down and results in a slowing down of the S_1 isomerization dynamics. Such a three-state model may also apply to other rPSBAT-hosting rhodopsins such as bR and Channelrhodopsin. In fact, this would be consistent with the reported bR⁸⁶ and, recently, Channelrhodopsin⁸⁸ reaction paths. It would also be consistent with the similar dynamics observed for ASR_{AT} and light-adapted bR (hosting exclusively rPSBAT)^{116,123} as well as for ASR_{13C} and the corresponding rPSB13-containing bR. Finally, the same three-state model may also control the S_1 dynamics of unsubstituted and certain substituted rPSBAT chromophores in solution that display S_1 picosecond lifetimes.¹²⁴ The presence of an S_2/S_1 degeneracy in solvated rPSBAT models

has been proposed¹²⁵ and held responsible for the observed solution dynamics of rPSBAT.¹²⁶ Similar data have been reported for rPSB11,^{49,127} suggesting that the Rh cavity has an important role of removing S_2/S_1 degeneracies along the photoisomerization pathway in order to optimize photoisomerization lifetimes (and, therefore, possibly also optimize photoisomerization quantum yields).

4.4 Summary: Conserved CIs and reaction coordinate variability. Above we have reported on the results of a set of comparative studies carried out to establish which mechanistic aspects of rhodopsin photoisomerization are conserved or undergo variation in different rhodopsins. It is found that the geometrical and electronic structures of the CIs are conserved and consistent with a 1Bu/2Ag CI reached via a NBCP type of motion (see Fig. 4.8). The nature of the motion along the reaction coordinate also appears to be conserved and starts, in general, with a BLA relaxation which then immediately couples with HOOP motion. The reactive double-bond skeletal isomerization motion activates after the initial relaxation. We have also pointed out that the stereochemistry of the isomerization is, for the examined rhodopsins, of the CW type for C11=C12 and C15=NH and CCW for C9=C10 and C13=C14. Such stereochemistry appears to be consistent with the documented pre-twisting of specific double bonds of the chromophore. We can therefore hypothesize that the degree and direction of pre-twisting imposed by the protein cavity determines which double bond is selected for the reaction and in which direction, CW or CCW, it will twist.

The mechanism summarized above needs to be further detailed for the case of the reaction coordinate of an inverse photoisomerization reactions. For bathoRh to Rh photoisomerization, which starts from a largely CW pre-twisted bathoRh intermediate, the reaction coordinate follows a reaction path which is almost exactly inverted with respect to the Rh to bathoRh path. Therefore, notice that for a directly inverse (with "directly" we mean starting from the primary photo-product of the reaction) process the C11=C12 bond rotates CCW and the C9=C10 CW. This situation is markedly different from the one of the "direct" and "inverse" reactions occurring in the cyanobacterial rhodopsin ASR. In this case, we have documented reaction coordinates which are part of a rotary motion driven by the absorption of two-photons and where the isomerizing double bond moves always in the same CCW direction. This is due to the fact that ASR_{AT} and ASR_{13C} are not directly interconverted but require additional thermal isomerization steps. In fact, they differ in the conformation of their side chain and stereochemistry of the C15=NH bond.

In contrast with the above conserved features, the reviewed results indicate that there are differences between the energy profiles and excited state lifetime of rhodopsins featuring the rPSB11 and rPSBAT chromophore. In fact, wild-type rhodopsin with the rPSBAT chromophore (ASR_{AT} and bR_{LA}) feature much flatter energy profiles. In Section 4.3.3 we have provided evidence supporting the involvement of a different number of electronic states for these type of rhodopsins. rPSB11 rhodopsins only involve two electronic states (S_1 and S_0) while the rPSBAT mechanism dominating the photoisomerization of microbial rhodopsins also involves the unreactive S_2 state. We

have seen that in these last rhodopsins S_2 partially mixes with S_1 leading to a flatter or, better, rugged S_1 potential energy surface consistent with a slower excited state progression towards the CI.

The reviewed results show that each documented CI may come together with two transition states (TS_{CT} and TS_{DIR}) on the S_0 potential energy surface. The one featuring charge-transfer character, TS_{CT} , is energetically favored and controls the protein thermal isomerization. Interestingly, the corresponding isomerization coordinate is very similar to the NBCP coordinate driving the excited state isomerization discussed above but has a different BLA value which display a larger degree of single-bond/double-bond inversion. Finally, it has been found that the similar charge-transfer characters (documented by looking at the charge translocation) of the FC and TS_{CT} structures provide a natural and direct explanation for observed and simulated Barlow-like relationships.

5. PHOTOACTIVE YELLOW PROTEIN

The photoactive yellow protein (PYP) is a 125 amino acid cytoplasmic blue-light photoreceptor controlling the photoavoidance response of the salt-tolerant bacterium *Halorhodospira halophila*.¹²⁸ PYP belongs to the Xanthopsin protein family and is its most studied member.¹²⁹ It is also the structural prototype for the PAS (PER-ARNT-SIM) class of signal transduction proteins.¹²⁹⁻¹³² PYP contains a deprotonated 4-hydroxycinnamic acid (or *p*-coumaric acid, pCA) chromophore linked covalently to the γ -sulfur of Cys69 via a thioester bond (see Fig. 5.1). Thus, in contrast to the cationic rPSB chromophore found in rhodopsins, PYP has an anionic chromophore where the phenolic oxygen forms short hydrogen bonds with nearby tyrosine (Tyr42) and glutamic acid (Glu46) residues (see Fig. 5.1B). Upon blue-light irradiation, PYP enters a fully reversible photocycle involving several intermediates formed on timescales ranging from a few hundred femtoseconds to seconds (see Fig. 5.2). Although a full understanding of the details of the PYP photocycle is still lacking, it can be divided into three major events: (1) PYP undergoes photoisomerization of the anionic chromophore to transform from pG (i.e. the starting structure) to yield the pR intermediate, which is a structure with a twisted C10=C11 bond (see Fig. 5.2B) and a red-shifted absorption, (2) protonation of the anionic chromophore and structural relaxation leads to the pB intermediate, which has a blue-shifted absorption with respect to the starting conformation; and (3) the recovery of the initial structure. The various intermediates are shown in the photocycle in Figure 5.2.

The detailed photoisomerization mechanism of the PYP chromophore has been debated for years. The light-induced dynamics of several models of pCA have been studied experimentally and computationally both in gas-phase¹³³ and in solution.^{134,135} In the following sections, we focus our discussion on the computational studies,¹³⁶⁻¹⁴⁰ which are considered complementary to their experimental counterparts. Specifically, in Section 5.1 we discuss the geometric and electronic structures of the CIs of pCA that mediate double-bond isomerization along different competing pathways. In Section 5.2 we review the proposed isomerization paths and mechanisms. Finally, in

Section 5.3 we report on trajectory studies and compare the results with the corresponding experimental observations.




Figure 5.1. A. Crystallographic structure of PYP in the pG state (PDB ID: 2ZOI). The protein backbone is drawn in ribbon representation with the p-coumaric acid chromophore (pCA) and relevant side chains in the licorice representation. B. Schematic representation of the hydrogen bond network formed by residues Tyr42 and Glu46. These residues, together with the positive charge on Arg52, help to stabilize the negative charge on the pCA phenolate. Hydrogen bonds are represented by dashed lines.




Figure 5.2. A. Photocycle of PYP from the perspective of time-resolved crystallography. Approximate timescales are given. The framed part of the photocycle corresponds to the photoisomerization, which occurs on a picosecond timescales that this present contribution reviews focusing on. Adapted with permission from ref. ¹⁴¹. Copyright 2016 The American Association for the Advancement of Science. B. Structure of I_T , pR_0 as compared with pG .

5.1 Conical intersections.

The isomerization of the PYP chromophore has been studied using the isolated chromophore (pCK^-) as a model.^{137,139,142} These studies suggest the existence of two possible pathways for the decay of pCK^- from S_1 to S_0 : α -twist and β -twist (see Fig. 5.3). The corresponding sloped CIs are located in the vicinity of S_1 minima whose structures are shown at bottom of Fig. 5.3. The α -twist path induces a complete transfer of the negative charge from the phenolic ring to the alkene fragment where the C9-C10 bond length is found to be 1.47 Å. In contrast, the β -twist induces only a partial charge transfer in the same direction leading to a stretched single bond (which has a bond length of 1.47 Å) rather than to a double-bond twisting. Such preliminary gas-phase studies suggest that protein residues or solvent molecules forming hydrogen bonds with the phenolic oxygen are likely to play an important role in the selection of the isomerization pathway, and drive the S_1 relaxation along the twisting coordinate.

More recent studies have characterized the S_1/S_0 CI reached via a twisting about the C10=C11 double bond, corresponding to the β -twist of pCK^- . The CI has a C9-C10-C11-C12 dihedral angle of 85°, compared to 170° at the equilibrium geometry.

The CI is geometrically close to the experimentally observed S_0 intermediate I_T (see Fig. 5.2), which also has a twisting angle of 85° in the crystal structure.¹⁴³ In the same computational study,¹⁴⁰ it is shown that the optimized intermediate I_T has a C9-C10-C11-C12 angle of 80°. However, the experimental assignment of the I_T structure is not without controversy.^{144,145} While the computational study by Wei et al.¹⁴⁰, which was performed at the CASSCF level, report on a twisted I_T intermediate, DFT calculations^{144,146} indicate that the I_T structure is structurally closer to the fully isomerized *cis* product with a C9-C10-C11-C12 angle of 21°. Kaila et al.¹⁴⁴ suggested that such conflict is the consequence of the arbitrary choice of loose restraints in the X-ray data refinement. However, Jung et al.¹⁴⁵ pointed out that the discrepancies might be due to the experimental conditions and crystal preparation, even if the same time-resolved X-ray crystallography were employed. While the exact origin of discrepancies remains to be understood,¹⁴⁵ the most recent femtosecond time-resolved X-ray study shows no evidence for such a ~90° twisted intermediate, but finds an isomerization that leads directly to a ~30° twisted structure (pR_0 in Fig. 5.2B).¹⁴¹




Figure 5.3. Scheme of the pCA chromophore, the pCK^- chromophore analogue and the corresponding excited-state minimum energy configurations of the analogue: (a) α -twisted S_1 minimum and (b) β -twisted S_1 minimum. Adapted from ref. ¹³⁹. Copyright 2009 American Chemical Society.

5.2 Reaction paths.

In order to understand the photochemical properties of the PYP chromophore, the S_1 potential energy surface of the pCK^- gas-phase model was studied at the CASSCF level.¹³⁹ As anticipated above two excited-state minima were identified (see Fig. 5.3): the α -twisted minimum, which features a 90° twisted bond adjacent to the phenol ring, and the β -twisted minimum, which features a 90° twisted ethylenic bond. In pCK^- one finds a negligible barrier for reaching the α -twisted S_1 minimum from the FC region, whereas there is a significant barrier for reaching the β -twisted minimum. Hence, in the gas-phase, the main relaxation channel on S_1 would proceed along the α -twist coordinate. Similar results have been found at the CC2 level of theory.¹⁴⁷ Since isomerization in the protein proceeds exclusively along the β -twist, these results suggest that the PYP apoprotein controls the isomerization path making the β -twist the energetically favorable reaction coordinate. Another study

by Gromov et al. at the CC2 level focused on a pCK⁻ chromophore where the phenolic oxygen is stabilized by a number of water molecules.¹³⁸ Their study demonstrates that hydrogen bonding with water molecules can modulate the photoreaction pathways through charge stabilization of the phenolic oxygen to also make the β -twist pathway accessible.

As anticipated above, time-resolved X-ray crystallography studies, which provide information on the PYP light-induced dynamics, were employed to resolve the controversy over the PYP isomerization mechanism.^{141,143,148} Such studies^{143,148} suggested that there are two possible pathways for the photoisomerization. These are the HT mechanism suggested by Liu and Asato¹⁴⁹ and the BCP mechanism²² originally proposed for the rPSB chromophore in rhodopsin. The bonds involved in the HT mechanism in PYP are the C9-C10 and C10=C11 bonds. Instead, the BCP path is characterized by simultaneous C10=C11 and C12-S13 twisting (note that C12-S13 has a partial double-bond character because it is a thioamide), and is the only mechanism observed in the time-resolved X-ray study by Pande *et al.*¹⁴¹

The HT and BCP mechanisms have been investigated in PYP by computing the corresponding relaxed scans. Unlike animal rhodopsins, which feature a barrierless relaxation with no S₁ intermediate, here a shallow S₁ minimum is located for PYP with a C9-C10-C11-C12 dihedral angle of 155.5°.¹⁴⁰ This is consistent with a less reactive potential energy surface and therefore with the slower (550 fs) photoisomerization compared to the animal rhodopsin (e.g. Rh). However, it is comparable to the microbial rhodopsins with a rPSBAT chromophore (e.g. bR) displaying a flatter S₁ potential energy surface (see Section 4.3.3 above). In fact, as reported for bR, the S₁ relaxation continues downhill after the minimum and reaches the CI. The energy profile of the computed path is consistent with the low fluorescence quantum yield observed in PYP^{150,151} which can be assigned to the located S₁ minimum.

Inspection of the computed S₁ relaxation coordinate indicates that, even though the hydrogen bond between the phenolate oxygen of the chromophore and the Glu46 residue is weakened by the decreased negative charge, it still helps to restrain the phenyl ring from twisting along the C9-C10 bond (i.e. the one that would be involved in the HT mechanism). This C9-C10 twisting goes from -3.6° (at the FC point) to 3° (at the S₁ minimum) to 33° (at the CI). In order to continue such deformation on S₀, the chromophore has to overcome a 7 kcal/mol barrier to break the hydrogen bond between the chromophore and Glu46. On the other hand, along the S₁ relaxation following a BCP mechanism, the twisting of C10=C11 goes from 170° (at the FC point) to the 155° (at the S₁ minimum) and finally to 85° (at the CI). Simultaneously the twisting of the C12-S13 bond goes, as expected for such mechanism, in the opposite direction from -177° (at the FC point) to -168° (at the S₁ minimum) to -147° (at the CI). Upon relaxation to S₀ the twisting about the C12-S13 bond has to overcome a smaller S₀ barrier (4.4 kcal/mol) with respect to the HT and does not require the breaking of the hydrogen bond with Glu46. This suggests that the BCP mechanism is more favorable than the HT mechanism, even though both mechanisms are feasible. The smaller barrier for the BCP mechanism (see Fig. 5.4)

shows consistency with the shorter lifetime determined experimentally. The BCP mechanism leads to the formation of I_{CT} (~1.7 ns) while the HT mechanism leads to the formation of pR₁ (lifetime = ~3 ns).¹⁴³

The conclusions above are based on relaxed scans, which only provide an approximate description of the reaction coordinate. An improved description can be achieved with a minimum energy path or trajectory studies that incorporate the effects of the light-induced dynamics, as discussed below.

5.3 Trajectories.

In order to investigate the photoinduced dynamics of the isomerization of the PYP chromophore, trajectory studies have been carried out in the gas-phase,^{137,139} in solution,¹³⁸ as well as in the protein.¹³⁶ Semi-classical trajectory calculations by Groenhof and co-workers¹³⁹ with non-adiabatic transitions described by Robb's approximation (see Section A.I.3) followed the progression of gas-phase pCK⁻ on the S₁ potential energy surface.

These molecular dynamics simulations in the gas phase confirm the result of reaction-path calculations. Indeed, it is found that, in the gas phase, the trajectories reach the S₁ minimum along the α -twist coordinate. It is also found that the S₁/S₀ IS lies far away from the S₁ minimum making the S₁ decay inefficient. Again, this demonstrates the importance of the environment in controlling the mechanism and timescale of the S₁ decay of the pCA chromophore.




Figure 5.4. Stationary points from a relaxed scan along the dihedral angle of C9-C10-C11-C12 on the S₁ potential energy surface. This coordinate describes the photoisomerization of PYP. The scan shows the location of stationary points up to the formation of first ground-state intermediate I_T that is formed upon hopping to ground state. The barriers in the ground state are shown on the right side for the Hula-twist or bicycle-pedal mechanism starting from the I_T intermediate to the next step. The numbers indicate CASPT2 energies in kcal/mol where the blue color highlights the S₁ state and red the S₀ state.¹⁴⁰

To examine the effect of hydrogen bonding on the PYP chromophore, a simple water cluster model was employed to perform QM/MM trajectory calculations.¹³⁹ The QM subsystem was treated at the CASSCF level. The results demonstrate that the predominant S₁ decay channel involves a α -twist (88%)

rather than a β -twist (22%) mechanism. However, this is an important change from the gas-phase where the decay is exclusively via the α -twist. This is explained by the fact that the water molecules hydrogen bonded to the carbonyl oxygen would stabilize the α -twisted minimum while those hydrogen bonded to the phenolate oxygen would stabilize the β -twisted minimum, thus modulating the excited-state pathways. The transient stabilization of the charge distribution would bring the IS close to these minima causing a faster decay. A faster decay in solution has also been documented by Martínez and coworkers using *ab initio* multiple spawning (AIMS) calculations.³⁴ The results, which are consistent with the CC2 reaction path studies^{138,147} mentioned in Section 5.2 above, also indicate that the S_1 dynamics and decay timescale are modulated by hydrogen bonding. Naseem *et al.*¹⁵² further studied the PYP chromophore in solution and found that there is an increase in pK_a in the *cis* form but not in the *trans* form due to electrostatic repulsion between the coumaryl tail and the phenolate ring moieties. This aids the protonation of the *cis* form in the next step of the photocycle.

In order to understand the effect of the PYP protein environment, Groenhof *et al.*¹³⁶ carried out QM/MM trajectory computations at the CASSCF level with Robb's surface hopping algorithm. They showed that the S_1 pCA chromophore rapidly decays to S_0 through a β -twist motion instead of the α -twist motion as reported in the gas-phase. The authors have attributed it to hydrogen bonding between the phenolate ion with the nearby Tyr42 and Glu46 residues. Similar to the effects in the water environment discussed above, these hydrogen bonds enhance S_1 decay from the β -twisted minimum. Moreover, these simulations also demonstrated that the same hydrogen bonds remain intact along the isomerization coordinate, which follows a NBCP mechanism consistently with the reaction path studies. At the surface hop, there is a ca. 90° twist at the C10=C11 bond while the adjacent C12-S13 bond is twisted by ca. 50° . The simulations show a 200 fs fluorescence lifetime and 30% isomerization quantum yield and are therefore close to the corresponding experimental values (550 fs¹⁴¹ and 35% quantum yield¹⁵¹).

In contrast to the solution environment, the protein in PYP causes the chromophore to isomerize exclusively along the β -twist coordinate, with none of the excited state population undergoing an α -twist.¹³⁶ This indicates that, apart from hydrogen bonding to the phenolate part of the chromophore, there is also a large effect of the protein residues on tuning the isomerization mechanism of the chromophore. In particular, the positively charged guanidinium moiety of Arg52 plays an important role in biasing the relaxation towards double-bond (β) isomerization. Other studies³⁴ have confirmed that when a small positive point charge (+0.2) is placed in the position where the nitrogen atom (N ϵ in Fig. 5.1) of the Arg52 residue is, the α -twist motion is suppressed while the β -twist is pro-

motored, while a negative point charge would produce the opposite effect.³⁴ Such an effect has been experimentally assessed by producing the Arg52Gln mutant. It was shown that Arg52Gln can still enter the photocycle, although with a lower rate and quantum yield.¹⁵³ The conclusion is that Arg52 is important but not essential for photoactivation.

In an attempt to simulate the effects described above, QM/MM trajectory calculations of the PYP Arg52Gln mutant were performed by Groenhof *et al.*¹⁵⁴ They find that without a positively charged Arg52, the predominant S_1 relaxation follows an α -twist rather than a β -twist coordinate. It was also found that in the Arg52Gln mutant, the hydrogen bond between the carbonyl oxygen of the chromophore and the backbone amino group of Cys69 is broken. However, the chromophore rapidly forms three new hydrogen bonds involving the amino groups Tyr98 and Asp97 as well as a water molecule that enters the chromophore pocket from the solution. These hydrogen bonds stabilize the negative charge on the alkene moiety and lead to a rapid decay of the chromophore to S_0 . In conclusion, the Arg52Gln mutant of PYP has a similar decay mechanism as the chromophore in water. Even though the α -twist path does not produce the *cis* diastereoisomer of the chromophore, the computations show that it causes variations in the hydrogen bond network and the flipping of the thioester group. However, the same processes were also computationally demonstrated to occur during the β -twist relaxation. It has thus been proposed¹⁵⁴ that the flipping of the thioester group and not the β -twist is the key step in the photocycle activation. This would be consistent with the fact that, as it is experimentally observed, the PYP Arg52Gln mutant is still photoactive.¹⁵⁵

5.4 Controversies regarding low-barrier hydrogen bonding and photoisomerization mechanisms. All the PYP mechanistic studies discussed so far are based on models featuring with protonated Arg52 residue and an anionic chromophore. However, an alternative scenario has been debated where Arg52 is deprotonated and the anionic chromophore is protonated to become a neutral one. More specifically, this would require a low-barrier hydrogen bonding (LBHB) in PYP LBHB is a non-standard hydrogen bond network, which was originally proposed to possess covalent bond-like character.¹⁵⁶ As opposed to standard hydrogen bonding, where the hydrogen is covalently bonded to one heteroatom and more weakly interacting with the other, LBHB is a stronger intermolecular interaction where the hydrogen is more equally shared between two heteroatoms. LBHB between the Arg and the chromophore ion pair would cancel out, in part, the negative charge of the chromophore and the positive charge on Arg52 yielding a more neutral charge environment. This controversy was triggered by a neutron diffraction study by Yamaguchi *et al.*¹⁵⁷ suggesting the existence of a neutral, LBHB bounded chromophore with a neutral Arg52 residue in PYP. In fact, the existence of LBHB allows the stabilization of the anionic pCA chromophore without the need of a charged Arg52. However, Saito *et al.*¹⁵⁸ suggested that there is no LBHB in PYP based on QM/MM calculations and electrostatic pK_a calculations. Later, Nadal-Ferret *et al.*¹⁵⁹ suggested that, PYP displays LBHB in solid phase while, it reverts back to a regular hydro-

gen bonding in solution. Further studies by Graen *et al.*¹⁶⁰ suggested that a single PYP model in vacuum is not necessarily a good model to describe the experimental results measured in solid phase as they found that a model with multiple copies of PYP, which would better describes the PYP in crystals, show no LBHB. The discrepancy is caused by the fact that the real environment around PYP is sufficiently isotropic due to the high ionic strength in crystals but a single PYP in vacuum is anisotropic, which is not a good model to reflect the solid phase of PYP. Moreover, this study has re-examined the neutron densities found by Yamaguchi *et al.*¹⁵⁷ and suggested that the large differences in experimental neutron densities at the side chain of Arg52 do not exclude the possibility of a protonated Arg52. Even though most of the studies point towards a fully protonated Arg52 with no LBHB, the effect of LBHB versus deprotonated Arg52 on the S_1 dynamics of PYP has yet to be studied computationally. This is particularly interesting as it has also been proposed, on the basis of ultrafast stimulated Raman studies, that the initial S_1 relaxation involves conversion from LBHB to a normal hydrogen bonding before the photoisomerization occurs.^{161,162}

The controversy over the photoisomerization mechanism of PYP is also still unresolved. The study by Jung *et al.*¹⁴³ on the picosecond timescale has suggested that the photoisomerization of the double bond can proceed via two structurally distinct *cis* intermediates through both the HT and BCP mechanisms. However, studies by Pande *et al.*¹⁴¹ at the femtosecond resolution suggest that the photoisomerization proceeds exclusively through the BCP mechanism which is consistent with the simulations from Groenhof *et al.*¹³⁶

Both HT and BCP mechanisms would ultimately lead to pB (i.e. the *cis* distereoisomer of the chromophore) and generate the signaling state of PYP, even though the BCP mechanism would be the major pathway. However, a recent experimental study¹⁶³ of a different PYP from *Leptospira biflexa* (Lbif) suggested that there is a bifurcation in the photocycle of PYP. Lbif PYP shows an apoprotein different from the most widely studied PYP from *Halorhodospira halophila* (Hhal) even if it has the same pCA chromophore. The study suggests that the long-standing inconsistency in the existence of a bifurcation in photocycle at room-temperature is due to a small population of intermediates only found in the most computationally studied Hhal PYP. These results will likely motivate computational studies of Lbif PYP to see whether any bifurcation exists leading to different isomerization mechanisms.

5.5 Summary: A possible bifurcating photocycle. The results of different simulations of the PYP photoisomerization are, in general, consistent with the experimental observations. They suggest an S_1 BCP relaxation mechanism leading to a CI. Even though the HT mechanism is not observed in trajectory simulations inside the protein, the existence of a bifurcation cannot presently be ruled out, especially with the recent experimental finding on a different apoprotein that could lead to detour in the photocycle.

Unlike rhodopsins, PYP has an anionic chromophore with a relatively small apoprotein. This has led to the controversies regarding low-barrier hydrogen bonding. Due to its small pro-

tein size, the chromophore is partially exposed. This would make a single PYP in vacuum not a good model to reflect the solid phase of PYP. These findings suggest that it is important to have a model that has a good description of the chromophore environment that is consistent with the experimental conditions in order to give an accurate simulation.

At any rate, the mechanism that controls the primary steps of the photocycle of PYP is a double-bond photoisomerization, regardless whether HT or BCP mechanism is followed. Then, it undergoes a breaking of a hydrogen bond between the chromophore and protein backbone that triggers a proton transfer from the protein to the phenolate oxygen of the chromophore. This leads back to the pB structure and the signaling state of PYP. Since both HT and BCP would lead to the same pB, the BCP mechanism must follow an aborted isomerization, just as in Rh.

PYP is unique among natural photoreceptors because it forms large and stable crystals that are suitable for X-ray diffraction. This system has therefore provided an excellent opportunity not only to simulate the time-resolved isomerization computationally, but also to “see” it experimentally with time-resolved X-ray crystallography. PYP therefore has led the way for the application of time-resolved X-ray crystallography in photobiology, and has presented a unique and exciting opportunity to compare more directly the worlds of QM/MM simulations and X-ray crystallography. Despite this, there is still more to be done to better understand the details of the isomerization mechanism of the PYP chromophore and the formation of the signaling state, mainly due to understanding the differences between the gas phase protein and solid state as indicated above.

6. PHYTOCHROMES

6.1 Phytochromes: CI’s, reaction coordinates, and trajectories. Phytochromes are a family of photoreceptor proteins that bind a linear tetrapyrrole as a chromophore. They are characteristic among photoreceptors for their absorption in the red, which leads to a photoproduct with absorption in the far-red edge of the visible spectrum. Phytochromes are photochromic and can be switched between the red absorbing (P_r) and far-red absorbing (P_{fr}) states. While originally discovered in higher plants where they play an important role in regulating circadian rhythms, members of the phytochrome family of proteins have also been found in bacteria, cyanobacteria, and fungi where they regulate a variety of biological functions.¹⁶⁴ An example of a prototypal plant phytochrome is shown in Fig. 6.1A. Not all phytochromes share the same architecture, although the photosensory module of most canonical phytochromes consists of three conserved domains; PAS, GAF, and PHY domains.¹⁶⁵

The chromophores of phytochromes are methine-bridged linear tetrapyrroles (known as bilins) that are bound to a cysteine residue of the apoprotein through a thioether group, as shown in Fig. 6.1B. Phytychromobilin ($P\Phi B$) acts as the chromophore of plant phytochromes, while in bacterial and fungal phytochromes the chromophore is phycocyanobilin (PCB) or biliverdin (BV). Despite these small differences in the chro-

mophore structures, spectroscopic studies have indicated that phytochromes of plants, bacteria, and fungi all share a common photocycle, shown in Fig. 6.2. In the dark-adapted state, most phytochromes exist in the P_r state. Resonance Raman (RR) studies^{166,167} indicate that photoexcitation of chromophore at ca. 660 nm initiates a double-bond isomerization of the methine double bond that bridges rings C and D (the C15=C16 bond in Fig. 6.1C), leading to the Lumi-R intermediate. After a series of thermal relaxations of both the chromophore and the protein, the P_{fr} active state is reached, which is called "active" because it triggers the biological function. The protein can then be reverted back to the P_r state by exciting in the far red (at ca. 730 nm) to give Lumi-F (although this step may also occur thermally on a much longer time-scale). Back-conversion does not follow the same mechanism as the forward process, but rather goes through different intermediates; Lumi-F and Meta-F.




Figure 6.1. A. Crystal structure (PDB ID: 4OUR) of a plant phytochrome B dimer from *Arabidopsis thaliana* showing the PAS, GAF, and PHY domains as well as the PΦB chromophore. Adapted with permission from ref. ¹⁶⁸. Copyright 2014 National Academy of Sciences. B. The structures of Phytochromobilin (PΦB), phycocyanobilin (PCB) and biliverdin (BV) in the ZZZssa (i.e. 4-Z, 10-Z, 15-Z, 5-syn, 9-syn, 14-anti) conformation. Note that the three chromophores differ only at rings A and D. Only one resonance structure is shown. C. Model chromophore commonly used in gas-phase studies of PΦB, PCB, or BV. The reactive bond in phytochromes is indicated with an orange curved arrow. Important atoms are numbered to indicate bonds that are free to photoisomerize in the gas-phase.




Figure 6.2. The phytochrome photocycle. The first step of the photocycle, the *Z* to *E* isomerization of P_r to Lumi-R, occurs on a ps timescale and exhibits multi-exponential kinetics with both an ultrafast (2-3 ps) and one order of magnitude slower (30-40 ps) component.¹⁶⁹

The details of the photoinduced *Z* to *E* isomerization from P_r to Lumi-R, as well as the *E* to *Z* isomerization from P_{fr} to Lumi-F, have slowly begun to emerge over the past two decades. The understanding of the photochemistry of phytochromes has been largely driven by experimental spectroscopic studies employing RR^{166,167,170-173} NMR,^{174,175} as well as, more recently, from the analysis of the X-ray crystal structures^{168,176-178} of the P_r and P_{fr} intermediates. Many of these studies have been largely aided by theory as well. For example, theory has been instrumental in the assignment of RR spectra,^{166,172,173} and in the interpretation of polarization resolved femtosecond pump-probe spectroscopy.^{179,180} However, in contrast to the case of rhodopsins and PYP, there are very few computational studies attempting to determine the photoisomerization mechanism of phytochromes, e.g. by optimizing excited state reaction pathways or by calculating semi-classical trajectories. As we have done in previous sections, we will focus exclusively on such studies aimed at modeling the isomerization process in phytochromes. However, due to the limited number of studies, this section will not be divided into subsections related to funnels, paths, and dynamics as we have done in previous sections. There are a number of reasons why such computational studies have been relatively scarce:

1. The large size of the chromophore prevents the use of many of the quantum chemical methods discussed in Appendix I. For example, a CASSCF calculation that includes the full π -electron system in the active space would require an active space of 28 electrons in 25 orbitals, which is not tractable due to the factorial scaling of computational cost with the size of the active space. For this reason, electronic structure methods used to study phytochromes are less computation-

ally demanding, and therefore often more approximate. For instance, this often means using single reference methods like CIS or TD-DFT that are unable to treat the non-adiabatic regions of potential energy surfaces correctly,^{181,182} or using approximations that do not account for static and dynamical electron correlations in a balanced way (as explained in Section A.I.1). It is therefore difficult to achieve a solid description of the mechanistically important reaction paths and CIs, for instance, found on the excited-state potential energy surface.

2. Crystal structures of phytochromes have only become available in the last decade. The first crystal structure of a phytochrome was published in 2005 for the bacterium *Deinococcus radiodurans*,¹⁷⁶ while the first crystal structure of a plant phytochrome was only resolved recently in 2014.¹⁶⁸ This means that early investigations of the photoisomerization of the phytochrome chromophore were confined to the gas-phase, or an implicit solvation model such as the polarizable continuum model (PCM) as an approximation to account for the environmental effect of the protein.¹⁸³⁻¹⁸⁵ An added difficulty was that, for a long time, even the conformations of the chromophores in P_r and P_{fr} were unknown due to lack of a crystal structure. While there has been, for the most part, a general consensus that the photoisomerizations occurred at C15=C16, it was not clear if a single bond isomerization (e.g. a hula-twist mechanism similar to the one introduced in Section 4.3.1) would have occurred simultaneously or in tandem. In fact, before crystal structures of plant phytochromes became available, it was predicted that $P\Phi B$ is in the *5-Z,anti-10-Z,syn-15-Z,anti* (*ZZZasa*) configuration on the basis of RR studies interpreted using DFT gas-phase frequency calculations.¹⁷² This conclusion was also supported by SAC-CI calculations,¹⁸⁶ leading several early computational studies to adopt the *ZZZasa* configuration for computational studies. SAC-CI, short for "symmetry adapted cluster-configuration interaction", is a method that uses an exponential ansatz for the wave function parametrization similar to the EOM-CC (see Section A.I.1), but it uses a different exponential operator than the one used in canonical coupled cluster.¹⁸⁷⁻¹⁸⁹ It was only confirmed recently that $P\Phi B$ in plant phytochromes is in the distinct *ZZZssa* configuration and therefore different from the one originally hypothesized.¹⁶⁸
3. It has recently been found that, in several phytochromes, both the P_r and P_{fr} states do not exist in one homogeneous conformation but in two conformers (or sub-states) that quickly interconvert (e.g. see ref. ¹⁹⁰ and references therein). This complicates their modeling, particularly because the two sub-states are expected to have different photophysics/photochemistry (e.g. see ref. ¹⁸⁰), and therefore both need to be modeled to achieve a comprehensive understanding.

Unlike rhodopsins and PYP, phytochromes also have a more complex multi-domain protein structure, potentially making QM/MM studies difficult. For example, as displayed in Fig. 6.1A, a plant phytochrome monomer typically comprises three domains: GAF, PAS, and PHY. Rhodopsins and PYP are more compact.

Despite the difficulties outlined above, there have been a number of attempts to computationally investigate the photoisomerization mechanism of phytochromes. Most of these studies have been confined to gas-phase chromophore models, which serve as preliminary studies to understand the excited-state behavior of the chromophore in the absence of any external perturbation. This approach is similar, historically, to the case of rhodopsins where studies were first conducted with gas-phase calculations on reduced models of rPSB,^{20,191,192} only to evolve to full QM/MM simulations later on.^{21,193}

Early studies by Durbeej, Borg, and Eriksson¹⁹⁴⁻¹⁹⁶ of the phytochrome chromophore used a slightly reduced model of PΦB, where they replaced the C3 thioether linkage, the propionic carboxyl groups at C8 and C12, and the methyl groups at C2, C7, C13 and C17 by hydrogen atoms (see Fig. 6.1C) to compute a relaxed scan around the C15=C16 isomerizing bond. They used the CIS method to optimize the S₁ state at different constrained C15=C16 dihedral angles, and TD-DFT to compute the energies of the S₀ and S₁ states along these scans. Their studies indicated that the PΦB has a significantly lower rotation barrier when it isomerizes exclusively along the C15=C16 bond, as opposed to a hula-twist mechanism where an adjacent single bond also isomerizes.¹⁹⁴ However, since such studies investigated the gas-phase chromophore, it is unclear if the same mechanism will apply in the protein where there are steric constraints restricting large-amplitude motions of the chromophore. Durbeej *et al.* also found that deprotonation of the chromophore at the pyrrole nitrogen of ring B or ring C leads to a higher rotation barrier than if the chromophore remains protonated,¹⁹⁵ which indicated that the chromophore does not become deprotonated before isomerizing.

Later, Durbeej¹⁹⁷ used the same reduced model to compare the S₁ isomerization barriers around the C4=C5, C10=C11, and C15=C16 bonds (i.e. around the methine double-bonds bridging rings A-B, B-C, and C-D, respectively). This time, he used TD-DFT instead of CIS for S₁ optimizations. He also tried modeling the chromophore both in the gas-phase and in PCM. He found that the second excited state, S₂, which has doubly excited character with respect to S₀, is not accessed along the isomerization coordinate in either the gas-phase or PCM models. More interestingly, he found that the chromophore prefers to isomerize around the central C10=C11 bond in the gas-phase, while the isomerizations of the terminal double-bonds C4=C5 and C15=C16 have higher barriers in both gas phase and PCM. Since protein-bound PΦB isomerizes selectively along the C15=C16 bond,^{166,167} this indicates that the protein plays an important role in this selectivity. This is in line with the protein crystal structures, which indicate that the protein cavity forms a tight binding pocket around rings A, B, and C while ring D is less tightly bound. The gas-phase results were

later extended to PCB and BV,¹⁹⁸ indicating that the conclusions above are valid in bacteria, cyanobacteria, and fungal phytochromes as well. A few years later, Zhuang, Wang, and Lan¹⁹⁹ performed nonadiabatic dynamics simulations with on-the-fly surface hopping using the same reduced model as Durbeej to investigate the reactivity of the PΦB chromophore in the gas-phase. The dynamics were run with the semi-empirical OM2/MRCI method, but single-point calculations at the ADC(2)/TZVP level were used for verification. They found, in agreement with Durbeej,¹⁹⁷ that isomerization occurred exclusively around the methine bridge linking rings B and C, with no trajectories isomerizing around C4=C5 nor C15=C16. However, rather than isomerizing around C10=C11, most trajectories actually isomerized around the C9-C10 bond with only a few trajectories isomerizing around C10=C11. Recently, Falklöf and Durbeej²⁰⁰ investigated the isomerization of protein-embedded BV in a bacteriophytochrome using a full QM/MM model. They again performed a relaxed scan around the C4=C5, C10=C11, and C15=C16 torsions of the BV chromophore inside the protein. Their calculations confirmed that steric interactions with the protein pocket are the reason for the selectivity of the photoisomerization around C15=C16, because they restrict the isomerization at the A-B and B-C bridges. This is the only QM/MM isomerization study of phytochromes, so there is still very limited understanding of the atomic-level protein interactions controlling selectivity and photoisomerization mechanism.

Note that, unlike rPSB, excitation of the PΦB chromophore does not immediately lead to a significant change in dipole moment,¹⁹⁵ meaning that no significant charge transfer occurs upon vertical excitation. The excitation involves a HOMO to LUMO transition, with both orbitals localized on the central B and C rings.²⁰¹ The reason for this is most likely that the A and D rings are somewhat twisted (by ca. 45°)²⁰¹ with respect to the B and C rings, decreasing the degree of delocalization of the frontier orbitals along the full π-conjugated framework (i.e. forming a locally excited state). However, along the C15=C16 isomerization path, a large and sudden change in dipole moment does occur when the chromophore is ca. 60° twisted out of plane, indicating that a charge-transfer character has to develop along the S₁ path before decay.¹⁹⁵ This is similar to protonated Schiff bases, where twisting localizes the HOMO and LUMO π orbitals on either side of the isomerizing bond inducing a transfer of the positive charge from the Schiff base moiety to the opposite molecular end (see Fig. 3.5A).

As mentioned in the caption of Fig. 6.2, pump-probe experiments on BV-binding phytochromes have shown that they exhibit multi-exponential decay dynamics, with both fast (2-3 ps) and slow (30-40 ps) components.¹⁶⁹ In 2009, Altoè and coworkers²⁰¹ subsequently published a computational study where they attempted to explain the observed dynamics by using a gas-phase reduced model of the chromophore. They used the CIS electronic structure method for S₁ optimizations, but reaction paths were computed with CASPT2//CASSCF with a 10-electron 11-orbital active space. The reduced (10,11)

active space space was chosen based on the convergence of the vertical spectroscopic properties upon a systematic increase. They found that the gas-phase chromophore, upon excitation, initially relaxes from the Franck-Condon region to a shallow S_1 minimum. From that minimum, they performed a relaxed scan along four different rotation coordinates: C4=C5, C9-C10, C10=C11, and C15=C16 and found that isomerization of the C9-C10, C10=C11, and C15=C16 bonds are competitive because they have similar S_1 barriers, while C4=C5 is the only bond that is not likely to isomerize. Their calculations also indicate that isomerization at C15=C16 does not lead to a CI, but rather to a twisted intramolecular charge-transfer (TICT) state that has an S_0 - S_1 energy gap of 17.4 kcal/mol at the CASPT2//CIS level (or, when they attempted to re-optimize it at the CASSCF(10,11) level of theory, an energy gap of 5.4 kcal/mol with CASPT2//CASSCF). This twisted intermediate has its positive charge localized on the D ring, as opposed to the vertical excited state that had its charge localized still on the B and C rings. Such a TICT state would then have to decay to the ground state via a nearby sloped CI (see Fig. 6.3). On the other hand, isomerization around the C9-C10 or C10=C11 bonds leads directly to a peaked CI.




Figure 6.3. A TICT is a minimum on the excited-state potential energy surface reached by following a torsional deformation accompanied by a charge transfer. A TICT appears in a vicinity of a sloped conical intersection (see right panel) instead of a peaked conical intersection (left panel), and can decay to the ground state either by fluorescence or by accessing the nearby sloped CI.

Based on these observations, the authors²⁰¹ hypothesized that the isomerization of the C9-C10 or C10=C11 occurs on an ultrafast timescale, and therefore these processes correspond to the fast 3 ps component of the pump-probe experiments (grey panel in Fig. 6.4). Even if the protein restrains the isomerization motion, they propose that deactivation does not lead to an isomerized product, but returns back to the original P_r reactant. The other path, which is the biologically relevant C15=C16 isomerization, leads to a TICT that is longer lived and decays via a sloped CI which is higher in energy with respect to the minimum (see Fig. 6.3 right). Thus, the isomerization through this channel occurs on a slower time-scale. The authors assign this path to the experimentally observed slow ~30 ps component, consistently with pump-probe experiments indicating that the slow component corresponds to the biologically relevant C15=C16 isomerization (White panel in Fig. 6.4).¹⁶⁹




Figure 6.4. The top panel summarizes the mechanistic picture emerging from CASPT2//CIS calculations by Altoe *et al.*²⁰¹. The fast decay channel (~3 ps) corresponds to internal conversion via an aborted isomerization of the C9-C10 or C10=C11 bond. The slower decay channel corresponds to the biologically relevant C15=C16 isomerization. Geometries for the Franck-Condon point (FC), the planar excited-state minimum (PLA), and the TICT (D) are also shown. C-C-C-C Dihedrals (red) and bond lengths in Ångstrom (black) are labeled for exocyclic bonds. Figure and geometries from ref.²⁰¹. Copyright 2009 American Chemical Society.

6.2. Summary: The protein environment selects the isomerizing C15=C16 bond. Among photoreceptors that undergo double-bond isomerization, phytochromes and other related bilin-binding photoreceptors have been the least studied so far. There are several reasons for this, but chief among these reasons is the size of the chromophore (making the application of QM methods expensive and, consequently, limiting the understanding of its electronic structure) and the scarce availability of experimental data such as crystal structures until somewhat recently. While QM/MM studies on phytochromes are slowly starting to emerge, QM studies on chromophore models have paved the way to understanding the photochemistry and photophysics of the chromophore.

One of the main results emerging from theoretical simulations is that the chromophore in the gas phase isomerizes preferen-

tially around the central C9-C10 or C10=C11 bonds, but the phytochrome protein environment modulates the selectivity, most likely through steric interactions, and causes the isomerization to occur along C15=C16 instead. Experimentally, pump-probe experiments on BV-binding phytochromes have shown that they exhibit multi-exponential decay dynamics with a slow (30-40 ps) and fast (3 ps) component, so computational studies have also posited that this could be due to competing C15=C16 and C9-C10/C10=C11 isomerization in the protein.

With new emerging QM methods and more efficient algorithms, computational studies on phytochromes are almost certain to increase in the near future. Because of the size and extent of conjugation of their tetrapyrrole backbone, bilins absorb naturally in the red and can emit in the far-red as well. The BV chromophore is also naturally occurring in mammalian cells due to breakdown of heme.²⁰² This has important implications for their potential use in optogenetics as well as bioimaging; since mammalian tissue is more transparent at far-red and near-IR wavelengths, the development of optogenetics tools or fluorescent proteins that absorb/emit in the far-red or infra-red is highly desirable for deep-tissue processes. Recent studies have demonstrated that this is a practical and promising possibility.^{203,204} This is almost certain to further catalyze the use of computer simulations to understand the photoisomerization of BV and related chromophores in different proteins.

7. OTHER PROTEINS SHOWING DOUBLE-BOND ISOMERIZATION

7.1 Rhodopsin mimics.

To systematically study the correlation between structure, absorption wavelength and photochemical and thermal reactivity in retinal proteins, the intracellular lipid binding protein family (iLBP), and in particular the human cellular retinoid binding protein II (hCRBP II) and human cellular retinoic acid binding protein II (CRABP II), have been employed as targets for protein engineering.^{54,205-207} Tolerance to mutations and the large binding cavity of iLBPs allowed flexibility in the redesign of the protein and enabled the binding of a range of different ligands.²⁰⁸ More specifically, the relative rigidity of the iLBP backbone structure makes it an ideal scaffold for protein engineering because it is more likely that mutations will not result in structural rearrangement,²⁰⁸ allowing the effects of specific mutations to be compared directly. These systems have been engineered to bind the rPSB chromophore such that it is almost fully embedded within the binding pocket. This is an important property of these systems because otherwise exposure to aqueous media would buffer electrostatic changes and result in the loss of effectiveness of the protein in modulating the photophysics of the embedded chromophore. Furthermore, hCRBP II and CRABP II are water-soluble proteins instead of membrane proteins and thus much easier to crystallize.

Over the past few decades, Borhan, Geiger, and coworkers have prepared and investigated several iLBP-based rhodopsin mimics. These authors exploited the ease of crystallization of

these engineered protein and, consequently, have access to detailed structural information via crystallographic methods. Although, most of their reported studies focus on spectroscopy and color tuning,⁵⁴ one recent report²⁰⁹ presenting a reengineered CRABP II system that displays both photochemical and thermal isomerization of the embedded retinal chromophore (see Fig. 7.1A). Before this study, it has been difficult to demonstrate that photoisomerization of the protein-bound retinylidene chromophore (usually rPSBAT) occurs in these types of engineered systems. More specifically, Borhan, Geiger and coworkers have reported on an iLBP rhodopsin mimic where the chromophore specifically isomerizes at the protonated Schiff base -CH=NH- double bond both thermally and photochemically. This isomerization was characterized at atomic resolution by quantitatively interconverting the isomers in the crystal (in the solid state), both thermally and photochemically.²⁰⁹ This event is accompanied by a large pK_a change of the imine similar to the pK_a changes observed in bacteriorhodopsin and visual opsins during isomerization.

Although there is presently no computational study on the type of isomerizing rhodopsin mimics described above, different groups have attempted to reproduce the observed change in absorption maximum upon mutations of an hCRBP II system.^{55,210} In one case the mechanism of the photoisomerization process has also been investigated²¹¹ in a CRABP II mimic, but this is in a different system from the one where the isomerization has been detected experimentally. In this study, the authors investigated the viability of rPSBAT photoisomerization from all-*trans* to 11-*cis* and to 13-*cis*. Since the protein pocket of CRABP II is adapted to bind the rPSBAT chromophore (i.e., the all-*trans* isomer), the computed isomerized products were predicted to be unstable. Indeed, the optimized geometries of rPSB11 and rPSB13 are significantly distorted, with dihedrals -27° and -9° twisted out of plane, respectively. These deformations extend to other parts of the chromophore backbone. From Fig. 7.1B, it can be seen that the isomerization leads, in both cases, to large skeletal changes that mainly involve the N-containing segment of the chromophore. This happens even though the β -ionone ring sticks out of the protein pocket and would be, in principle, allowed to rotate.

Further computational studies in this area promise to reveal how to construct rhodopsin mimics undergoing photoisomerization and, hopefully, showing photochromism and bistability. This could be done, for instance, by constructing and studying comparatively models of rhodopsin mimics prepared in the lab and shown to be either photochemically reluctant or active like the one of ref. ²⁰⁹.




Figure 7.1. Artificial rhodopsin mimics. A. Photoconversion of a CRABP II rhodopsin mimic in the crystalline state from ref. ²⁰⁹. Copyright 2016 American Chemical Society. The metastable *cis*-imine structure is obtained from a crystal that was UV irradiated for 30 min. Above it is displayed the crystal structure of the *trans*-imine which can be obtained via visible light irradiation of the *cis*-imine structure. B. Structures and main dihedral angles (deg) for the computed 11-*cis* (left) and 13-*cis* (right) primary photoproducts. The backbone dihedral values display deformations that deviate from planarity (0 or 180°) by more than 5°. At the bottom, we report the superposition of these two isomerized chromophores with PSBAT in PSB-KLE-CRABP II. Adapted from ref. ²¹¹. Copyright 2013 American Chemical Society.

7.2 Fluorescent proteins.

Double-bond isomerization in photobiological systems is by no means exclusive to rhodopsins, phytochromes, and the PYP chromophore. Fluorescent proteins (FPs) are another example of proteins where photoinduced double-bond isomerization of the chromophore can take place. Photoinduced processes, including double-bond isomerization, in FPs have been reviewed recently by Acharya *et al.*²¹² FPs have a rich photophysics where several excited-state processes may occur. This includes isomerization, proton transfer, and even electron transfer. The interplay of these excited-state processes is a subject of many theoretical and experimental studies in different types of FPs.²¹²

FPs have revolutionized the field of bioimaging,²¹³ and since the discovery of the Green Fluorescent Protein (GFP)²¹⁴ a palette of FPs with colors spanning almost the entire visible spectrum have been prepared by mutagenesis.²¹⁵ However, although the fluorescent properties of FPs have been widely exploited in biotechnologies, the biological function of wild-type FPs is poorly understood. Therefore, whereas double-bond isomerization triggers biological responses in rhodopsins, phytochromes, and PYP, double-bond isomerization in FPs is a secondary and often undesirable process. This is especially true when it comes to biotechnological applications, because double-bond isomerization competes with fluorescence and may lead to quenching. On the other hand, it has been shown that double-bond photoisomerization in FPs can be exploited for specific purposes, such as in the development of a new generation of photoswitchable FPs such as Dronpa (see Fig. 7.2).

The majority of FP chromophores are made up of an imidazolinone ring conjugated with another aromatic ring via a methine bridge. In GFP and in Dronpa, the aromatic ring is phenolate (Fig. 7.2B) which is reminiscent of the phenolate already seen in the PYP chromophore.




Figure 7.2. Dronpa, a photoswitchable fluorescent protein. The crystal structure of Dronpa²¹⁶ is reported on the left. The chromophore of Dronpa can isomerize between an "on" *cis*-isomer (green structure) and "off" *trans*-isomer (grey structure). The crystal structure is shown in the "on" state, with the chromophore in its *cis* conformation. The Lewis formulas are adapted with permission from ref. ²¹⁷. Copyright 2014 Elsevier.

The methine bridge in the GFP chromophore is the only segment of the π -conjugated framework that is not locked in a ring and can therefore rotate around the single bond (ϕ) or double bond (τ). Rotation along either ϕ or τ usually leads to a twisted excited-state intermediate (TI_ϕ and TI_τ).^{218,219} In early computational studies of a GFP model chromophore, a minimum energy CI was found along a hula twist coordinate ($MECI_{hula}$) that involves the out-of-phase twisting of both ϕ and τ , but in the gas-phase it lies high in energy relative to TI_ϕ and TI_τ . More recent studies have found low-lying CIs in the gas phase that are twisted only along ϕ ^{220,221} and τ ²²⁰ ($MECI_\phi$ and $MECI_\tau$, respectively). The relative energetics and geometries of such intermediates and minimum energy CIs has been the subject of numerous computational studies.^{218-220,222,223} However, it is clear that the solvent environment,^{219,222,223} protein environment, protonation state of the chromophore,²²¹ as well as chromophore substituents all have a substantial impact on the relative energetics of these CI points and the pathways leading to them. Therefore, the photophysics of the chromophore is highly dependent on such factors.




Figure 7.3. Schematic representation showing key structures on the excited-state potential energy surface of a gas-phase GFP chromophore (top-right Lewis formula model). After excitation, the chromophore relaxes from the Franck-Condon point (FC) to the fluorescent state (FS). From the FS, the molecule may evolve by twisting along ϕ or τ . Twisting around ϕ is controlled by a barrier TS_ϕ and leads to a twisted intermediate with $\phi \approx 90^\circ$ (TI_ϕ). Twisting along τ is controlled by a barrier TS_τ and leads to a twisted intermediate with $\tau \approx 90^\circ$ (TI_τ). A conical intersection ($MECI_{hula}$) is reached along a hula-twist motion involving both torsion along ϕ and τ , but this CI is energetically high in energy. Other minimum energy conical intersections were located closer in energy to the twisted intermediates that are distorted along a bond-stretching/pyramidalization coordinate with respect to the intermediates ($MECI_\phi$ and $MECI_\tau$). This figure was adapted with permission from ref. ²¹⁸. Copyright 2004 American Chemical Society and updated to account for results presented in ref. ²²⁰.

Twisting around either ϕ or τ is a major decay pathway in many FPs, leading to twisted S_1 intermediates (TI_ϕ and TI_τ) with nearby low-lying CIs. This is partly also responsible for phenomena such as emission intensity variations, on-off blinking, and photoreversible switching between light and dark states.²²⁴⁻²²⁷ When the chromophore is not held rigidly in the protein binding pocket (e.g., in gas-phase or solution), the barriers for rotation (TS_ϕ and TS_τ) are small, which explains why fluorescence quantum yields of GFP chromophore models *in vacuo* or in solution are several orders of magnitude smaller than in the protein.²²⁸⁻²³⁰ Conversely, restricting the motion of the GFP chromophore by placing it in a rigid environment increases the barriers to rotation (TS_ϕ and TS_τ) and leads to brighter fluorescence.^{231,232} Steric interaction is not the only effect at play here, since hydrogen-bonding and electrostatic interactions with the proteins likely play a very important role in influencing the ease of rotation around ϕ or τ .

In the vast majority of FPs, the chromophore is in the *cis* conformation.²³³ On the other hand, the *trans* conformer is usually not fluorescent, with a few notable exceptions. This possibly stems from the fact that the *trans* form of the chromophore is tilted or twisted along ϕ or τ to relieve a steric clash between the imidazolinone carbonyl oxygen and the phenol ring, making the *trans* form more conformationally flexible than the *cis* form.²³⁴ The presence of a long-lived transient dark *trans* state can be detrimental to the functionality of FPs in single-molecule studies,²³⁵ because it contributes (or is a major source of) blinking and intensity variations. However, the same phenomenon can be exploited in photoswitchable FPs if

it can be controlled. One notable example, mentioned above, is Dronpa,²³⁶ where the native state is bright (*cis* form) but can be switched off (to the *trans* isomer) by illuminating with intense blue light, or switched back on again with weak near-UV light. Dronpa is an example of a negative photoswitch because it is bright in its native state and can be turned off with light. There are also several examples of positive photoswitches, such as Padron²³⁷ and kindling FPs, that are dark in their native state (typically, in the *trans* conformation) but their fluorescence can be turned on momentarily by photoisomerization to the light state.²³⁷ The exact mechanism of photoswitching in such photoswitchable FPs is a subject of debate,²¹² in particular with regards to the protonation states of the chromophore. This has complicated the computational investigation of isomerization in such systems. However, the photoswitching occurs on an ultrafast timescale,²³⁸ which indicates that a low-lying CI must be accessed during photoisomerization. For example, a study by Morokuma and co-workers²³⁹ employing a CASSCF/AMBER model of Dronpa found that, regardless of the protonation state of the chromophore, isomerization occurs via a one-bond flip around τ rather than a hula-twist CI. On the other hand, a more recent study by Morozov and Groenhof²⁴⁰ found qualitatively different results. They investigated a Dronpa mutant using twelve QM/MM semi-classical trajectories computed at the SA(2)-CASSCF(6,6)/3-21G/AMBER level of theory and found that while some of the trajectories (8 out of 12) did not isomerize at all in 50 ps, four of the trajectories decayed via a hula-twist mechanism involving twisting of both ϕ and τ . Such a hula-twist CI was also optimized by the same authors at the XMCQDPT2 level of theory.²⁴⁰

7.3 Twisting about a double bond as a deactivation channel in biomimetic or biological ring-locked double bonds.

As discussed in several of the sections above, not all twisting around a double bond leads to an isomerized product. Since isomerization quantum yield is never 100%, there are non-reactive (or aborted) isomerization events that lead back to the original reactant. Such aborted isomerization events contribute to internal conversion, since they simply depopulate the excited state at a twisted CI only to regenerate the original reactant.

Sometimes, such an aborted isomerization is the only possibility. For instance, double bonds locked in a ring smaller than an eight-membered ring often cannot isomerize due to steric constraints imposed by the ring. Cyclooctene is the smallest cycloalkene that is stable in the *trans* conformation at room temperature.²⁴¹ *trans*-cycloheptene may exist at lower temperatures, but at room temperature isomerizes quickly to *cis*-cycloheptene through a low-energy pathway that involves formation of a diradical dimer.²⁴²




Figure 7.4. Three examples showing ring-locked double bonds. *cis*-locked

rPSB11 employs a five-membered ring to prevent isomerization around the C11=C12 bond resulting in completely different photophysics. NAIP molecular switches are rPSB11 mimics that employ rings to induce selectivity in double-bond isomerization (to mimic the steric effect of the protein environment in rhodopsin). Finally, HBDI-like switches mimic the HBDI chromophore/fluorophore of GFP but have only one isomerizable double bond.

The fact that smaller cycloalkenes cannot form isomerized *trans* products has been exploited in many instances for locking double bonds to prevent their isomerization. One example of this presents itself in *cis*-locked rhodopsins, which are artificial constructs where the C11=C12 bond of the rPSB11 chromophore is locked in the *cis* conformation by a five-membered ring (Fig. 7.4). *Cis*-locked bovine rhodopsin has been prepared in the lab and has an excited-state lifetime of ca. 85 ps, which is almost three orders of magnitude longer than the lifetime of wild-type (unlocked) rhodopsin.^{243,244} Since *cis*-locked rPSB11 in bovine rhodopsin cannot isomerize along the C11=C12 bond, computational studies have indicated that it decays by twisting of an adjacent C9=C10 double bond instead.²⁴⁵

The use of rings to lock double bonds in a certain configuration has also been useful for the design molecular switches, where all double bonds but one are incorporated in a ring to prevent their isomerization.²⁴⁶ This makes it possible to construct light-driven molecular switches that isomerize selectively along one specific exocyclic double bond, such as in the case of the N-alkyl-indanylidene-pyrrolinium (NAIP),^{18,247-249} Fluorenylidene-Pyrroline,²⁵⁰ and benzylidene-pyrroline rPSB11 mimics.²⁵¹ The same concept has also been used to synthesize HBDI mimics called HBDI-like switches.²⁵²

corresponding structure of the CI of uracil (structure from ref. ²⁵³) is also shown at the top right, with the dihedral representing its torsional deformation labeled. Figure adapted with permission from ref. ²⁴⁶. Copyright 2014 John Wiley and Sons, Inc.

While ring-locked double bonds may not be able to isomerize, they may still undergo internal conversion via a twisted CI similar to the case of an aborted isomerization. One example in proteins is found in the putative fluorophore of bacterial luciferase, which is a derivative of a flavin mononucleotide (FMN). In particular, the fluorophore is an N(5)-ethyl 4a-hydroxy flavin (Et-FIOH). While EtFIOH is strongly fluorescent in the bacterial luciferase protein,²⁵⁴ the same fluorophore in solution has very weak fluorescence.²⁴⁶ A joint computational and experimental study found, using a methyl FIOH model (Me-FIOH, Fig. 7.5A), that the first singlet excited state (which is the fluorescent state) is repulsive and does not have a fluorescent minimum (Fig. 7.5B). Instead, from the vertical Franck-Condon point, it evolves sequentially along a BLA mode (similar to that observed in other conjugated organic chromophores such as rPSB11) followed by an out-of-plane deformation of the pyrimidine ring. The deactivation BLA and torsional deformation route is barrierless on the excited state, and leads to a twisted CI with the pyrimidine twisted almost 90° with respect to the rest of the flavin plane. The documented Me-FIOH deactivation mechanism and CI structure is in some ways similar to internal conversion of pyrimidine nucleic acid bases (i.e. cytosine, thymine, and uracil), which also follow a BLA-torsional deformation deactivation mechanism.^{253,255,256} However, there are some distinct differences between the case of Me-FIOH and pyrimidine bases, as shown in the comparison with uracil in Fig. 7.5B. For example, the BLA pattern is almost opposite in the pyrimidine ring of Me-FIOH and in uracil. Moreover, the uracil CI has a twisted C=C double bond (orange dihedral in Fig. 7.5) that involves a high degree of pyramidalization of one of the carbon centers and can be accessed via a facile hydrogen out-of-plane twisting. The Me-FIOH CI, on the other hand, is only partially twisted along the corresponding C=C bond but instead is mainly twisted along an adjacent N=C bond (red dihedral in Fig. 7.5). This twisting mode involves a large structural deformation due to movement of the entire pyrimidine ring with respect to the original flavin plane, which explains why such a mode is not accessed in a binding cavity and fluorescence is much higher in the protein.

8. CONCLUSIONS AND PERSPECTIVES

Photobiological systems are incredibly complicated, and a comprehensive understanding of their mechanism requires a full understanding of electronic and nuclear processes occurring on a range of timescales. Currently, a single model alone cannot possibly tell a full story. Therefore, photoreceptors have been the subject of numerous experimental and theoretical studies investigating different aspects of their mechanisms. In this review, we have focused on a relatively narrow topic; the QM/MM modeling of double-bond isomerization in rhodopsins, photoactive yellow proteins and phytochromes. Even with such a specific topic, it has been impossible to be exhaustive due to the volume of publications tackling these systems. Therefore, as mentioned in the introduction, the present review




Figure 7.5. Aborted double-bond "isomerization" in a flavin derivative. A. Structure of Me-FIOH. B. CASPT2//CASSCF S_0 and S_1 energy profiles for a solvated Me-FIOH model (the solvent is accounted for implicitly with PCM) along a minimum energy path calculation connecting the FC to a CI. The bond lengths for bonds that change by more than 0.05 Å during the MEP (in black) and the dihedral angle representing the ring deformation (in red) are labeled for the FC structure, the CI, as well as for the structure reached after an initial relaxation along a stretching mode. The

is not fully comprehensive but revolves around the background and research interests of the authors. More precisely, it reflects the progress made by a number of computational chemistry schools that started to study photochemical reactions in the 1990s. One of these schools, which focused on the development and application of the CASSCF method, emerged from a collaboration between Fernando Bernardi and Michael A. Robb²⁵⁷ and has contributed significantly to the computational results reviewed above. However, the theoretical background required for studying such reactions and, in particular, the double-bond photoisomerization of conjugated molecules, was laid out by numerous earlier studies. For instance, the electronic states driving the photoisomerization of a model rPSB had already been described by Josef Michl and Vlasta Bonačić Koutecký in the late 1980s.^{19,258} Michl and coworkers also originally highlighted the pivotal role of CIs in organic photochemical reactions and tested these hypotheses using configuration interaction computations.^{17,258} Even earlier than that, 2013 Nobel laureate Arieh Warshel provided the theoretical background for understanding the effect of the environment on the rPSB chromophore by developing, implementing and applying, for the first time, QM/MM technologies to derive mechanistic information. Thus, effectively, Warshel initiated the transition from computational photochemistry to computational photobiology using semi-empirical quantum chemical methods before practical *ab initio* multiconfigurational quantum chemical tools became available.

In the 90s, several groups carried out methodological and applicative research in computational photochemistry (e.g. CI optimization methods, reaction path calculations, and trajectory propagation) that made the studies reviewed here possible. Such work is only marginally covered here, but has been the subject of a recent review.¹⁷ For instance, Schlegel and coworkers developed an effective method for locating CIs.²⁵⁹ Yarkony and coworkers²⁶⁰ developed rigorous methods and theories for locating and characterizing CIs.^{60,261} Martínez and coworkers have reported on practical CI optimization methods and non-adiabatic dynamics approaches and used them to investigate several biological chromophores.¹¹⁴ Morokuma and coworkers have reported an effective branching plane calculation method.²⁶² Authors such as Schulten²⁹ and Birge⁹⁹ were among the first to perform semi-classical trajectory computations for Rh, bathoRh,⁸⁵ and bR.^{108,263,264} Needless to say, such progress in photobiology was dependent on progress on the theoretical photochemistry front. For instance, Domcke,^{256,265} Köppel,²⁶⁶ Cederbaum,²⁶⁷ and Martínez²⁶⁸ and coworkers, among others, played a vital role in rigorously demonstrating how ultrafast decay occurs at CIs. The demonstration, by quantum dynamics calculations, that few vibrational degrees of freedom are sufficient for irreversible radiationless decay at CIs led to a fundamental change of paradigms in radiationless decay theory. This understanding has been central to photochemistry in general and complemented the above-mentioned computational demonstrations that low-lying CIs are ubiquitous and common features of the excited states of sizable organic and biological molecules.

In conclusion, several authors created the scientific and technical basis for the computational investigation of ultrafast photochemical reactions in light-responsive proteins and, there-

fore, for the understanding of the biological function of double-bond photoisomerization. In the following subsections we draw some general conclusions (Section 8.1) and highlight directions (Section 8.2) that, in the authors' opinion, would be important for evolving from a scientific understanding towards the design of new molecular materials. In fact, until recently, theoretical chemists working on biological photoreceptor proteins have mainly focused on method development with the aim of exploring and uncovering the molecular mechanisms underlying protein function. Today we are witnessing a change of paradigm as some of the same scientists who studied the mechanism of photobiological systems are becoming more and more like engineers who can use computations and lab models to design for new functions. However, this transformation requires integration of methods for predicting and analyzing conformational changes, hydrogen bond networking, protein-chromophore interactions and thermal/photochemical reactivity before revealing new alleys towards the engineering of unprecedented systems.

8.1 General conclusions.

The experimental investigations of light-responsive proteins operating through double-bond isomerization have examined different "stages" of the primary photoproduct formation. When referring to Scheme 3.1 these are: light absorption and initial relaxation, reaction coordinate, excited-state intermediate and/or CI structures, and, finally, reaction time scales and quantum yields. In this review, we have tried to discuss selected groups of computational investigation addressing one or more of these stages in different proteins. Here, we will try to summarize the general mechanistic information that has resulted from these investigations.

8.1.1 Initial relaxation. The initial event in the "activation" of the examined biological chromophores corresponds to an allowed π - π^* excitation which may be described as a charge-transfer state. In cationic chromophores such as rPSB or P Φ B, this implies a partial translocation of the positive charge away from the C=N Schiff base (in rPSB) or C=N pyrrole bond (in P Φ B) while, in anionic chromophores such as in pCA, this implies a translocation of the negative charge away from the phenolic oxygen. In the framework of the Franck-Condon approximation, such instantaneous change in the electronic structure creates a non-equilibrium nuclear configuration that then relaxes out of the FC point. One of the first results of the computational work presented above has been the description of such a motion that is dominated by the expansion of the double bond and contraction of the single bond (the BLA coordinate). This implies the "unlocking" of the double bonds of the system as the primary effect of photon energy absorption. Notice (see Section 3.1) that in neutral and symmetric conjugated systems such as polyenes, these states would not promptly generate a charge translocation and therefore an inversion between double bonds and single bonds but, rather, a charge separation producing structure with almost equivalent single and double-bond lengths.

The computational investigations reviewed here have also indicated that the BLA relaxation can be effectively coupled to other molecular modes by the chiral protein environment. One example is the protein-induced stabilization of a chromophore conformation with a pre-twisted double bond. This would effectively couple the BLA mode with specific torsional molecular modes resulting in a multimodal initial excited-state relaxation. For instance, we have seen that in certain cases, such as that of animal rhodopsins, the initial relaxation effectively couples the BLA and the HOOP hydrogen wagging motion. A few femtoseconds later this bimodal relaxation couples with the twisting deformation of the carbon framework and, effectively, initiates the evolution along the isomerization coordinate. It is thus apparent that the coupling described above is central to determining the stereoselectivity and regioselectivity of the isomerization. This can be sterically or electrostatically imposed by pre-twisting a specific double bond of the chromophore and/or inducing an asymmetric force field (i.e. potential energy surface shape) that results in stereoselectivity/regioselectivity of the isomerization. Such steric and electrostatic effects result in the selection of the C11=C12 or C13=C14 bonds instead of the C9=C10 bond in rPSB, the β instead of α isomerization in pCA, or the C15=C16 bond instead of the other double bonds in P Φ B. On the other hand, computational studies have shown that the pCA chromophore in gas phase or solution is more likely to isomerize along α ,^{138,139,147} while P Φ B *in vacuo* is more likely to isomerize along C9=C10 or C10=C11.¹⁹⁷⁻¹⁹⁹ Therefore, the protein environment has a large control on the stereoselectivity of photoisomerization for different chromophores. Understanding this control has already been the subject of much interest for the computational community, because a better understanding of such control is central to the engineering of artificial proteins.

8.1.2 Reaction coordinate. Comparison of excited-state reaction paths and trajectories of different isomerizing chromophores (e.g. rPSB, pCA) reveals that the bicycle-pedal or aborted bicycle-pedal mechanism is commonly encountered in photobiology. This is because such a coordinate is space-saving, but also favorable if one considers the electronic character of the excited state of these chromophores. In fact, the initial relaxation along BLA naturally unlocks two nearby double bonds. In contrast, a hula-twist coordinate would require the twisting deformation of a double and a single bond simultaneously, thus increasing the energy cost necessary to achieve such a space-saving motion. Of course, when a bicycle-pedal motion cannot take place, such as in P Φ B or in the HBDI chromophore of fluorescent proteins featuring a single double bond located outside a cyclic framework, the hula-twist reaction coordinate may become operative if leading to a favorable space saving motion. The hula-twist reaction coordinate may become competitive with the bicycle-pedal coordinate when there is a penalty in undergoing bicycle-pedal type motion, as in pCA where the bicycle pedal type motion requires breaking the hydrogen bond between the carbonyl oxygen of the chromophore and the backbone amide in a nearby amino acid. Notice that in polyenic hydrocarbon systems, the initial relaxation along the locally excited (charge separated) π

- π^* state has a high probability of populating a lower energy diradical electronic state with different potential energy surface features. As we have discussed in Section 3.1, such electronic structure leads to a kink-type conical intersection that forces evolution along a hula-twist-like excited-state reaction coordinate and, ultimately, conical intersection.

Even if the bicycle-pedal reaction coordinate seems to be the most favorable, it is often aborted upon decay leading to the effective isomerization of a single double bond. The selection of the aborted isomerization has not been thoroughly investigated but, in rPSB is connected with the amplitude of the twisting deformation. For instance, in Rh a ca. 90° twisting about the reactive C11=C12 bond is accompanied by just a 60° twisting about the adjacent C9=C10 bond, which, therefore, has the largest probability of reverting back to its original configuration. In the bicycle-pedal reaction coordinate of PYP, the chromophore also has a ca. 90° twisting about the alkene moiety while the adjacent C=S bond is only ca. 50° twisted.

8.1.3 Excited-state dynamics and decay at the conical intersection. Trajectory calculations have largely confirmed the character of the excited-state electronic and geometrical progression indicated by reaction coordinate studies. On the other hand, new information has been provided on the factors that control the excited-state lifetime (i.e. the photoisomerization speed) and isomerization quantum yields. Animal rhodopsins have emerged as the fastest photoisomerizing proteins with estimated excited-state lifetimes as low as 60 fs²⁴ and experimentally detected lifetime as low as 70 fs in bovine rhodopsin³² (in fact, in a recent paper Miller and co-workers emphasized the possibility that the initial photoproduct could form in an even shorter time-scale).¹⁴ This speed has been attributed to factors related to the slope of the barrierless potential energy surface connecting the FC point to the CI. However, it has also been documented that a prerequisite for achieving such a speed is the energetic separation between the potential energy surfaces of the first excited state (i.e. the spectroscopic π - π^* charge-transfer state) and the higher electronic states such as the nearby diradical excited state. This separation guarantees a continuous acceleration towards the CI without entering a rugged and flat potential energy surface where the nature of the electronic character of the excited state may vary. This last factor has been proposed to explain the different dynamics observed for animal rhodopsins and microbial rhodopsins.²³ The reason for these differences is related, of course, to the less distorted all-*trans* rPSB isomer in microbial rhodopsins which features the isomerization of the C13=C14 rather than C11=C12 bond.

A second reason for observing a slower double-bond photoisomerization dynamics in proteins is the formation of transient excited-state intermediates whose reactivity is restrained by an excited-state barrier or by the existence of a sloped rather than peaked CI. This has been computationally found in the pCA chromophore of PYP and the P Φ B chromophore of phytochromes. In P Φ B, the intermediate appears to correspond to a TICT state featuring a complete charge transloca-

tion and the same 90° twisted geometrical structure of the CI but located energetically above it.

Finally, as documented in Section 3.4, the analysis of the result of non-adiabatic trajectory calculations has started to clarify the relationship between reaction speed and quantum yield. This appears not to be related to a simple proportionality relationship as a ballistic picture of the isomerization motion would suggest. In contrast, the currently emerging picture is focusing on the achievement of a phase relationship between the different vibrational modes populated immediately after photoexcitation and therefore on an extremely short time-scale. Such a control would require an ultrafast excited-state dynamics of the excited-state population whose function would be to guarantee the conservation of the molecular level (not necessarily population level) coherence between the modes that contribute to the reaction coordinate (e.g. the HOOP and the skeletal reactive torsional motion). According to this mechanistic model, a productive trajectory will have to display a certain phase relationship at the moment of decay. A faster excited-state motion (or a slower one) may result in a phase relationship allowing exclusively the return to the initial reactant double-bond configuration and, statistically, in a quantum yield decrease. Presently, such a mechanism has been considered exclusively in animal rhodopsin. Microbial rhodopsins may not utilize it (e.g. when using the rPSBAT chromophore), but instead could utilize other mechanisms for achieving high quantum yields that are applicable to situations where the excited-state motion is slower and less coherent.

Finally, in an ultrafast regime where vibrational coherence is maintained during the entire excited state evolution, photo-product formation seems to occur according to an oscillatory mechanism, which appears to be connected/determined by the HOOP-like motion occurring along the reaction coordinate. This has been reported for an abridged model of ASR_{13C},⁷³ a microbial rhodopsin isomer for which vibrational coherence has been observed.⁷⁰

This type of behavior and its significance is presently investigated using more realistic models (e.g. a 5 double-bond chromophore model) as highlighted in the graphical TOC of the present review.




Figure 8.1. Oscillatory behavior in the photoproduct formation of an abridged ASR_{13C} model (ASR_{13C}-M). Top. QM/MM model of ASR_{13C} with a PSB3-like chromophore and a reduced chromophore cavity. The curly arrow indicates the reactive C13=C14 double bond. Bottom. Time evolution of 400 reactive (red) and nonreactive (green) hop points plotted as a function of the variable α and β and of the time. The Newman diagrams provide the definition of the geometrical parameter affecting the overlap between the p-orbitals centered at C13 and C14 and describing the double-bond reconstitution. Adapted from ref. ⁷³. Copyright 2016 American Chemical Society.

8.2 Present limitations and future perspectives.

8.2.1 Advancements in electronic structure methods. Most of the computations reviewed in Sections 4-7 have been possible thanks to decades of developments leading up to modern electronic structure methods, such as those discussed in Appendix I. Ideally, to study the mechanism of photobiological systems one requires methods that can: (a) quantitatively describe the relative energies of ground and excited-state potential energy surfaces, (b) describe the near-crossing and crossing regions of potential energy surfaces, and (c) be computationally affordable. When one wants to describe transitions through crossings and avoided crossings, accurate non-adiabatic couplings are also needed. CASSCF is one of the methods that most prominently has made possible, and still makes possible,²⁶⁹ the investigation of biological chromophores. The CASSCF implementations by Michael A. Robb²⁵⁷ and by the member of the Swedish school led by Bjorn Roos had, as far as the present authors are concerned, a major impact. As a consequence, computer packages such as GAUSSIAN²⁷⁰ and MOLCAS,²⁷¹ allowing the integration of state-average CASSCF gradients and non-adiabatic couplings into QM/MM technologies, have been applied widely. However, CASSCF does not correctly describe reaction barriers and relative energies between electronic states (i.e., point *a* above). For this reason, the CASPT2 method, developed by Roos and coworkers, had, and still has,²⁷² a paramount importance for correcting CASSCF energies. However, CASPT2 gradients are still too expensive to be systematically employed in the type of studies describe above (point *c*). Furthermore,

CASPT2 suffers from some artifacts when it comes to describing near-crossing regions (point *b*), although this has been remedied by more recent extensions to the method. Such drawbacks notwithstanding, this is the reason for the popularity of the CASPT2//CASSCF protocol in photobiology.

The issues above indicate that there is a need for electronic structure methods that can yield accurate but affordable QM/MM models. The further development and efficient implementation of analytical gradients for CASPT2 and multi-state multi-configurational methods such as MRCI,²⁷³⁻²⁷⁵ MS-CASPT2,²⁷⁶ XMCQDPT2,¹¹⁸ QD-NEVPT^{277,278} and XMS-CASPT2^{279,280} will certainly impact trajectory calculations and mechanistic studies. In fact, recently, Shiozaki and coworkers have reported both analytical gradients²⁸¹ and non-adiabatic couplings²⁸² computed at the XMS-CASPT2 level. Meanwhile, many research groups continue to push the boundaries of electronic structure theory as methods continue to be developed, improved, and made more efficient. Some of these methods are already used to study photobiological systems, while others could very well be used in the near future. TD-DFT has been widely used, although primarily for geometry optimizations and spectroscopic studies due to its limitations when it comes to describing near-intersection regions. However, several developments addressing these near-intersection issues, such as DFT-configuration interaction methods (e.g., CIC-TDA-TD-DFT,²⁸³ CDFT-CI,^{284,285} SI-SA-REKS,^{182,286} and spin-flip TD-DFT,^{287,288} may result in more mechanistic studies that employ DFT. DFT methods are also being combined with multi-configurational methods such as CASSCF to account for the missing static correlation energy component.²⁸⁹ Coupled-cluster approaches such as linear-response CC and EOM-CC are also used due to their high accuracy and systematic improvability, and their application to photobiological systems is likely to increase as more and more efficient implementations become available.^{290,291} This is also true for ADC methods.²⁹²⁻²⁹⁴ Quantum Monte Carlo methods^{295,296} are also emerging as players in the photobiology community due to their favorable computational scaling and ability to account for electron correlation. These are just a few examples, out of many, of established or emerging electronic structure methods for studying biological chromophores.

While CASSCF and CASPT2 are used routinely for systems such as rPSB and the pCA chromophore, certain biological chromophores are still too large for such methodologies. This is in spite of recent progress in parallelization and fast algorithms such as the Cholesky decomposition.^{297,298} Therefore we often find that larger chromophores, such as PΦB or carotenoids, are not as well studied as rPSB or pCA. For example, a CASSCF calculation on PΦB that includes the full π -electron system in the active space would require an active space of 28 electrons in 25 orbitals, which is beyond the current computational limit of CASSCF. Although a reduced active space could be used,²⁰¹ this could lead to complications in the course of dynamics because the relative importance of selected configurations can vary in time as the geometry changes. However, development of new approaches built on the CASSCF concept, such as RASSCF²⁹⁹, GASSCF³⁰⁰, and DMRG^{301,302}, could help overcome such problems.

In order to increase the quality of a QM/MM model, not only is the electronic structure method describing the QM subsystem important but also the description of the interaction between QM and MM subsystems. In fact, the search for more accurate and computationally effective QM/MM models is another important technical challenge that computational photochemistry and photobiology faces. This calls for the development of more realistic force fields to describe the protein and its interaction with the QM subsystem. The importance of polarizable embedding (Section A.I.2.1) has been pointed out in several pilot studies applied to the calculation of excitation energies in photoreceptor proteins.³⁰³⁻³⁰⁵ However, while several polarizable embedding implementations exist and have been used to study spectroscopic properties, there are no mechanistic or dynamics studies using polarizable embedding yet for photoreceptors. Further implementations are needed to realize a combination of polarizable embedding with non-adiabatic dynamics. The same can be said for DFT embedding schemes, which are also powerful and accurate but need to be further developed before they can be used in mechanistic/dynamic studies of photoreceptors. Finally, another promising approach for combined quantum/classical description is Effective Fragment Potential (EFP).³⁰⁶ In this approach, a potential for solvent molecules is generated by calculating several parameters from *ab initio* calculations. The extension to proteins (macromolecule EFP, or mEFP) is very promising for future applications to photoreceptors.^{307,308}

One brute-force attempt for overcoming the problem of the interaction between the chromophore and its environment is that of extending as much as possible the size of the QM subsystem. This can be through calculations including clusters of amino acids surrounding the chromophore in the QM subsystem, all the way to including the entire protein in QM. The latter requires exceptional computing power not currently provided through conventional machines. However, hardware developments have made such calculations accessible. Already few years ago, it has been shown that video gaming machines containing two consumer graphical cards can outpace a state-of-the-art quad-core processor workstation by a factor of more than 180 for Hartree-Fock energy and gradient calculations.³⁰⁹ Such performance enhancements make it possible to run large-scale Hartree-Fock and DFT calculations, which typically require hundreds of traditional processor cores, on a single graphical processing unit (GPU) workstation. For instance, benchmark molecular dynamics simulations are performed on molecular systems such as an aspartic acid molecule solvated by 147 waters (457 atoms, 2014 basis functions) at the HF level at a rate of 0.7 ps of computed dynamics per day on a single desktop GPU.^{309,310} The further development and application of this technology, especially for post-Hartree Fock methods, is expected to have a big impact on the field.^{311,312}

8.2.2 We need to look at electron, not only nuclear, dynamics. In Fig.1.4 and Section 3.4 we have discussed the importance of phase relationships between different molecular modes. Such a relationship is established within few femtoseconds after photoexcitation but ultimately has an impact on how excited-state dynamics unfold and therefore on the point when molecules hop from the excited state to the ground-state

potential energy surface. In order to completely understand the onset of such phase relationships we must answer questions related to the validity of the Franck-Condon principle and exact separation of nuclear and electronic motions. In fact, the charge-transfer electronic structure achieved upon photoexcitation involves a re-organization of the electron cloud and, possibly, an intimate coupling between such reorganizations and the onset of nuclear motions (e.g. the early BLA and the HOOP motions discussed for visual pigments). This type of coupled electronic and nuclear relaxation have already been computationally studied in ionization processes.³¹³ The same type of electronic-nuclear coupling may affect the “hop” event itself, influencing the nuclear motion and, possibly, the excited-state lifetime and quantum efficiency upon passage through the CI region.³¹⁴ These phenomena are connected with the rapidly advancing field of attosecond spectroscopy that could allow the observation of such coupling phenomena in biological chromophores on the timescale of the electronic response (the attosecond timescale).^{315,316} In molecules, attosecond pump-probe spectroscopy enables investigations of the prompt charge redistribution and localization that accompany photoexcitation processes, where a molecule is lifted from the ground Born-Oppenheimer potential energy surface to one or more excited surfaces, and where subsequent photochemistry evolves on femtosecond or longer timescales.

8.2.3 We need to investigate the coupling between isomerization and other processes. The engineering of genetically encodable molecular “machines” powered by light requires the understanding and simulation of coupling between double-bond photoisomerization and other chemical processes such as proton transfer. For instance, it is established that, in fluorescent proteins like GFP,³¹⁷ the excited-state HBDI chromophore deprotonation is preceding the formation of the fluorescent intermediate while in the Dronpa (Section 7.2) proton transfer is coupled to double-bond photoisomerization.³¹⁸ Similarly, in rhodopsins, the ground-state deprotonation of the chromophore follows the isomerization process at some stage along the photocycle.³¹⁹ While these functionally important processes are usually seen as sequential, little is known about the actual mechanism and timescale of the coupling. For instance, in rhodopsin the possibility that the isomerization and even the excited-state charge translocation occurring along the reaction coordinate³²⁰ could destabilize the hydrogen bond network and trigger successive events has been proposed and investigated both experimentally and computationally, (for a recent work see³²¹) but still with no conclusive picture.

On the other hand, coupling between proton transfer and electron transfer (PCET) has been shown to be implicated in a number of photobiological systems such as, most recently, in blue light using flavin adenine dinucleotide (BLUF) proteins.³²² In this case, free-energy simulations elucidated the role played by the active site conformations before and following photoexcitation.

8.2.4 Quantum chemistry and molecular evolution. In Sections 4.2.3 and 4.3.3 we have shown that QM/MM models can be employed for comparative investigations of the thermal and photochemical reactivity of rhodopsins. These studies belong

to research lines where quantum chemistry is employed for solving biophysical problems. The further extension of this approach to other light-responsive proteins, such as the ones treated in Sections 4-6, would certainly bring new progress. On the other hand, the authors of the present review believe that, in the near future, quantum chemical studies should focus on the possibility to model large sets of light-responsive proteins. This would be of basic importance for the development of computational photobiology, as several biological problems require the comparison of large sets of proteins. For instance, QM/MM models can be constructed for proteins representing the nodes of a phylogenetic tree and, therefore, protein sequences presently extinct (i.e. ancestral proteins). In this way the unavoidable overlap between, for instance, quantum chemistry and molecular evolution studies should lead to new research lines. This includes the search into the past history of a protein, but also information on how to engineer it (a chemical target connected with the preparation of new materials) or, simply, learning about how a certain extant protein has been achieved as a function of time (a biological target connected with evolution). In other words, QM/MM models would allow the user to virtually “resurrect” a protein in the computer, thus providing insight into the evolutionary history of molecules, a window on the past revealing how adaptive specializations may have occurred in Nature (see also the Lake Baikal fish rhodopsin example⁹⁶ in Section 4.2.3). Furthermore, this same information could provide inspiration for the engineering of highly efficient light-converting biomimetic systems such as the one seen in Section 7.1. In the more extreme view anticipated in Section 7.3, one can learn from such studies how to engineer efficient components for light-driven molecular devices including coupled chromophore/optical cavities or coupled chromophore/plasmons.^{323,324}

Rhodopsins, PYP's and phytochromes appear to be excellent targets for understanding how evolution has gradually taken control of such elementary, but versatile, process as light-induced double-bond isomerization. For instance, the visual rhodopsins of invertebrate and vertebrates are known to possess extraordinary adaptations for maximizing photosensitivity under low light conditions. For this reason the laboratory resurrection and functional characterization of a 200 million year old archosaur rod visual pigment has been reported.³²⁵ However, due to the effort and expense associated with laboratory manipulation, this experimental approach is limited to small numbers of inferred sequences, making the detailed description of important evolutionary transitions and the testing of evolutionary hypotheses difficult. Even if a sufficiently large number of ancestors could be resurrected in the lab, the related structural information could be difficult to achieve. In fact, this would require sufficient expression and the availability of crystal structures. These difficulties can be significantly decreased by *in silico* screening of the many, potentially interesting, ancestor structures. Such screening would be based on the fast production of consistent QM/MM models of rhodopsin proteins as well as of a set of mutants. The study of the corresponding model would allow biologists to limit the lab efforts to key structures carrying the maximum amount of information.

While approaches close to the one highlighted above have been reported by Morokuma, Yokoyama and coworkers,³²⁶⁻³²⁹ their applications focused on spectroscopy (color perception tuning) rather than photochemical reactivity. Furthermore, the report focused only on one or few QM/MM models. In order to overcome these limitations, hundreds, if not thousands, of QM/MM rhodopsin models can be generated automatically. Also, these models need to be based on *ab initio* multiconfigurational quantum chemistry, as they have to be able to also reproduce changes in the mechanism and dynamics of photochemical and thermal activation. A first attempt in such direction has been recently reported.³³⁰ Such technology, discussed in some more detail in Section A.I.2, is called Automatic Rhodopsin Model (ARM) and may be used to compute, for instance, the optical and photochemical properties of inferred ancestral sequences, then compared to modern visual pigments. The results of benchmark studies carried out using rhodopsins from different organisms (i.e. archaea, eubacteria, invertebrate and vertebrates) as well as eubacteria and vertebrate rhodopsin mutants, are displayed in Fig. 8.2A and 8.2B respectively.




Figure 8.2. Reported benchmark results of the ARM protocol. (A) Observed vs computed values for vertical excitation energies of wild-type rhodopsins. Rod rhodopsin Rh (Rh₁₈₁ indicate a model with an ionized 181 residue) and its bathoRh intermediate (orange), invertebrate and nonvisual vertebrate rhodopsins (blue), eubacterial sensory rhodopsin isomers (green), archaea rhodopsins (yellow), and a chimeric rhodopsin ChR_{C1C2} originating from a microbial eukaryote. A proton-pumping eubacterial rhodopsin PR is also reported. The colors indicate related rhodopsins (i.e. related by sequence homology as for SqRh and hMeOp, or because they are isomers as in both the bR and ASR cases, or because they are photocycle intermediates as for Rh₁₈₁ and bathoRh). (B) Comparison between computed and observed ΔE_{S1-S0} values (full circles) for a set of ASR_{AT}, ASR_{13C} (left) and Rh (right) mutants. Figures adapted and upgraded from ref.³³⁰. Copyright 2016 American Chemical Society.

In the future the community may be able to implement research projects that integrate currently distant disciplines such as *ab initio* quantum chemistry, computational molecular evolution and ancestral protein resurrection to achieve an unprecedented understanding of the functional divergence of light-responsive proteins. Starting these studies from proteins undergoing ultrafast double-bond isomerization will allow the

exploitation of existing knowledge on these systems. Accordingly, more effective computational protocols for the systematic construction of computer models of ancestral pigments need to be developed and benchmarked.

APPENDIX I

The acronyms encountered in several figure legends, such as 3-root-SA-CASPT2(IPEA=0)//2-root-SA-CASSCF/6-31G* and 2-root-SA-CASSCF/6-31G*/AMBER, refer to methods employed to calculate ground and excited-state potential energies and gradients. These acronyms are actually an extension of Pople's acronyms³³¹ used to indicate different "model chemistries".³³² The double slash "/" sign separates the level of theory used for computing the energy of the system (at the left) from the level of theory used for optimizing the geometry or computing the trajectory (at the right). Thus, 3-root CASPT2//2-root CASSCF means that energies were computed with 3-root CASPT2 as single point calculations on top of geometries that were obtained with 2-root CASSCF. The 3-root CASPT2 acronym implies that the corresponding zero-order wavefunction is the 3-root CASSCF wavefunction. The IPEA=0 label is a required CASPT2 parameter.²⁷¹

It is important to have a certain degree of understanding of these methods as the quality of the chemical information extracted from the calculations depends critically on their properties and limitations. More specifically, such acronyms refer to electronic structure methods that cannot be used in a black-box fashion and necessarily involve a number of approximations that need to be considered carefully to achieve a balanced description of spectroscopy and ground and excited-state reactivity.

A balanced description of the ground and excited state electronic structure appears to be necessary for chemical process that involves the breaking and making of π -bonds. In fact, as highlighted on several parts of the present review, double-bond isomerization may occur when a π -bond is broken homolytically, a process that involves the unpairing and re-pairing of electrons, or heterolytically, a process that involves a transfer of the bonding electron pair to one side of the broken bond. Therefore, the modeling of isomerization requires a suitable level of electronic structure theory that is capable of describing changes in electron spin-pairing and electron densities during isomerization. At the same time, the study of isomerization in photobiological systems requires an understanding of the effect of the protein environment on such a chemical reaction. These include the steric and electrostatic effects that influence the electronic structure of the reactive chromophore. However, due to the strong scaling of the cost of electronic structure methods with the size of the system, the treatment of an entire protein (i.e. a system containing thousands of atoms) is usually impractical if not impossible. Therefore, multi-scale methods, where the system is divided into two or more subsystems treated at different levels of theory, have become common tools for studying biological photoreceptors. When the protein and the chromophore are connected via a bond, a suitable approach to treat the frontier of these two subsystems is also required.

The present appendix comprises four parts reviewing the computational methods used in exploring double-bond isomerization in photoreceptor proteins. The first part (A.I.1) will focus on electronic structure methods commonly used for treating the breaking of π -bonds in the ground and excited states of biological chromophores. The second part (A.I.2) will describe two types of QM/MM multiscale methods. In the third part (A.I.3), the computational tools and the details regarding the calculation of non-adiabatic trajectories will be presented.

A.I.1. Electronic structure methods. The development and testing of electronic structure methods is a very active field of current theoretical chemistry research, with new methods being continuously developed by many groups around the world. Therefore, even if we limit ourselves to methods suitable for studying excited-state isomerization, a comprehensive review would require the presentation of a large arsenal of methods. For this reason, this part of Appendix I will only focus on a specific series of benchmark studies testing the ability of some of the more popular classes of electronic structure methods employed to study the energy changes occurring along the double-bond isomerization coordinate of a specific chromophore. We will not go into any detail regarding the formulation of these methods and their performance especially when it comes to studying other properties. We refer the readers to refs. ³³³ and ³³⁴ for a general background to excited-state electronic structure methods.

The benchmark studies that we will discuss employ a reduced model of rPSB11 that comprises three double-bonds (PSB3, see Fig. A.1A). Reduced models of rPSB11 have been used in many instances for benchmarking purposes or even as a model to understand the photochemistry of rPSB11 (for instance, see work by Olivucci, Garavelli, Robb and coworkers,^{16,20,191,335-339} Elstner and coworkers,^{340,341} Buss and coworkers,³³ Birge and coworkers,³⁴² Thiel and coworkers,³⁴³ Send and Sundholm and coworkers,^{344,345} Filippi and coworkers,^{346,347} Andrioniów and coworkers,^{348,349} and Wong and coworkers³⁵⁰).

PSB3 has many of the same features of the S_0 and S_1 potential energy surfaces as rPSB11. In both PSB3 and rPSB11, excitation from the S_0 to the S_1 electronic state corresponds to the 1Ag to 1Bu electronic transition described in Section 3.1 and results in a 30-50% positive charge transfer away from the Schiff base moiety.^{20,21,351} In the S_1 state, PSB3 and rPSB11 feature a very shallow barrier or barrierless path connecting the FC point to a peaked CI characterized by a ca. 90° twisted isomerizing bond, as shown in Fig. 3.5A.^{21,62,192,346} In both systems, the electronic character of the S_0 and S_1 potential energy surfaces varies between diradical and charge transfer along a loop encircling the CI and located along the BP.^{57,90} PSB3 also reproduces several important features of the rPSB11 S_0 potential energy surface; specifically, PSB3 features two transition states (TS_{CT} and TS_{DIR}), the former having charge-transfer character and the latter having diradical character, as shown in Fig. 3.5B.^{56,90} Therefore, PSB3 serves as a model system for rPSB11 and has been adopted to investigate the performance of different electronic structure methods. Its reduced size permits the use of methods not affordable for rPSB11.




Figure A.1. A. rPSB11 (left) and its reduced model PSB3 (right). B. The definitions of the bond-length alternation (BLA) mode, the hydrogen out-of-plane (HOOP) mode, and the torsion mode in PSB3. C. A schematic 2D representation of the S_0 and S_1 potential energy surfaces of PSB3 showing six of the seven pathways used to benchmark electronic structure methods (see text). The coordinate dominating the MEP_{CIS} and MEP_{TRANS} corresponds to a changing mixture of BLA, torsion, and HOOP, and therefore cannot be represented completely on the single 2D potential energy plot shown here. The MEP paths are mapped more fully in a 3D coordinate space plot as discussed in Appendix II. A specific torsion and HOOP mixture is also running along the CI intersection space (the IS curve already mentioned in Section 3.2 and reported in Fig. 2.3). The paths are labeled in yellow full (or dashed for hidden segments) lines, while key structures on the potential energy surfaces are labeled with blue (for minima and Franck-Condon points), orange (for transition states), and red (for conical intersections) full circles. The seventh benchmark path, the CI loop, is shown in Fig. A.2 because it lies on the BP orthogonal to the IS space.

Recently, we have mapped specific S_0 and S_1 paths of PSB3 and used them to determine the effect of the electron correlation on different regions of the corresponding potential energy surface. As explained below, the electron correlation energy is the electronic energy missed by the Hartree-Fock (HF) method where the interactions between electrons is accounted for only in a mean field way. This work was a collaborative effort involving several research groups and published over several installments.^{35,62,90,182,286,295,352-355} Overall, these studies employed seven different paths (see Fig. A.1). These paths have mechanistic relevance since they involve key structures on the S_0 and S_1 potential energy surfaces of PSB3. The first path, named MEP_{CT} , is a minimum energy path on the ground state that connects the *cis*-PSB3 equilibrium geometry, TS_{CT} , and *trans*-PSB3. The second path, MEP_{DIR} , connects *cis*-PSB3,

TS_{DIR}, and *trans*-PSB3. The third path is the BLA path, defined in Fig. A.1B, which connects the TS_{CT} and TS_{DIR} transition states but also intercepts a CI point (hereafter referred to as CI_{BLA} to distinguish it from other CI points located along the IS hyperline). Since the BLA mode removes the electronic degeneracy, it is one of the two modes belonging to the BP of the CI. The fourth path, MEP_{CIS}, is an S₁ path starting from the *cis*-PSB3 FC point and following a minimum energy path leading to a CI in the IS. This CI is geometrically and energetically distinct from CI_{BLA}, and so is labeled CI_{CIS}. Similarly, MEP_{TRANS} connects the *trans*-PSB3 FC point to another point on the IS hyperline, CI_{TRANS}. The sixth path, the CI "seam segment", belongs entirely to the IS and connects CI_{CIS}, CI_{BLA}, and CI_{TRANS}. MEP_{CIS} and MEP_{TRANS} span a mixed coordinate space dominated by BLA near the FC region, but then dominated by a mixture of torsion and HOOP. CI_{CIS}, CI_{BLA}, and CI_{TRANS} differ mainly along the torsion and HOOP coordinates. The final path, discussed in more detail below, is called the CI loop and is a circular path that lies in the BP and encircles CI_{BLA}.

The purpose of these benchmark studies is to go beyond understanding the performance of methods in reproducing one energy value (e.g., a vertical excitation). Instead, this allows us to look at the *differential* electron dynamical correlation effects on the potential energy surfaces involved in thermal and photochemical double-bond isomerization. Differential electron correlation energy arises due to changing electron density distribution along the molecular framework (e.g. when going from a delocalized to more localized electronic distribution), requiring different degrees of electron correlation to be accounted for. The seven pathways described above go through regions of the S₀ and S₁ potential energy surfaces with different electronic configurations and, therefore, their correct description requires a treatment of the changing electron correlation on an equal footing. Details regarding the construction of these paths and the results of the benchmarks using different methods are in refs. 35,62,90,182,286,295,352-355.




Figure A.2. A. A schematic representation of the PSB3 S₀ potential energy surface in the region of a S₀/S₁ CI and along a 2-dimensional plane defined by the BLA and the isomerization coordinates. Those two modes correspond to the BP vectors. The BLA path and the CI loop paths are indicated along the yellow and blue planes, respectively. B. The S₀ (opaque) and S₁ (translucent) CASSCF energy profiles along the BLA

coordinate. C. A more detailed scheme showing the CI loop path. Both the S₀ and S₁ surfaces are shown here. The loop is at the intersection of the circumference of the blue cylinder with the S₀ and S₁ potential energy surfaces (shown as red dashed lines) and lies in the BP defined by the isomerization coordinate and BLA. The isomerization coordinate here is a specific combination of C=C torsion and HOOP that results in the breaking of the π-bond. Along the loop, the wave function of each state is expected to change electronic character twice. Areas of the surface shaded brown have a predominantly charge-transfer character Φ_{CT}, while green regions have a predominantly diradical character Φ_{DIR}. D. The S₀ and S₁ CASSCF energies (top) and S₀-S₁ energy gap (bottom) along the CI loop as a function of α (as labeled in panel C). The energy profiles are shaded green or brown to show the dominant electronic character of the wave function for each state at that point (for the energy gap, only the character of the S₀ wave function is shown). Reproduced with permission from ref. 182.

We will focus on the results of two paths: the BLA path and the CI loop of Fig. A.2. The BLA path provides a stringent test for electronic structure theories because the states involved are dominated by open shell diradical (Φ_{DIR}) and charge-transfer (Φ_{CT}) characters respectively (see Fig. A.3). Φ_{CT} has a more localized π-electron density with respect to Φ_{DIR} and, therefore, the computation of the corresponding potential energies requires a method accounting for the variation in dynamic electron correlation energy. Along the BLA path, an unbalanced description of these electronic characters will generate an artificial shift in the S₀ and S₁ energies resulting in a change in the position of their crossing, giving a crossing geometry that depends on the accuracy of the method used.




Figure A.3. The dominant electronic configuration for the two electronic states observed for S₀ and S₁ in 90° twisted PSB3. Partially adapted with permission from ref. 286.

The CI loop is meant to probe whether an electronic structure method is capable of correctly describing the BP associated to a CI. Such loops encircling a CI point are often used to characterize the role of the CI in the photochemical process¹⁵ or to probe the topography of the CI.^{57,58,356-358} The BP has been introduced and discussed briefly in Section 3.2. At a crossing

between two electronic states of the same spin and symmetry, distorting the molecule along one of two specific degrees of freedom is the only way to lift the electronic degeneracy:

$$GDV_{IJ} = \frac{\partial}{\partial R} [E_I - E_J]$$

$$DCV_{IJ} = \left\langle \psi_I \left| \frac{\partial}{\partial R} \right| \psi_J \right\rangle$$

GDV is the gradient difference vector, DCV is the derivative coupling vector, R represents the nuclear coordinates, E_I and E_J are the potential energies of the intersecting states and ψ_I and ψ_J are the corresponding wave functions. These vectors are related to the \mathbf{g} and \mathbf{h} vectors;⁶⁰ GDV and \mathbf{g} are identical, while $DCV = \mathbf{h}/(E_I - E_J)$ and is therefore related to \mathbf{h} by an energy factor. GDV and DCV define the BP (or the \mathbf{g} - \mathbf{h}) plane.⁵⁸ Notice that, in practice, the BP vectors are not uniquely defined but the branching plane is. For instance, near a CI, the vectors may change as a function of geometry even when the change is infinitesimally small. This is equivalent to moving along an infinitesimal loop center on the true CI point.

The CI property arises naturally when considering the secular eigenvalue equation for the two states of interest that have been decoupled from other states in a block Hamiltonian:

$$\begin{vmatrix} H_{II} - E & H_{IJ} \\ H_{JI} & H_{JJ} - E \end{vmatrix} = 0$$

Where:

$$H_{II} = \langle \psi_I | \hat{H} | \psi_I \rangle$$

$$H_{IJ} = \langle \psi_I | \hat{H} | \psi_J \rangle$$

$$H_{JI} = \langle \psi_J | \hat{H} | \psi_I \rangle$$

$$H_{JJ} = \langle \psi_J | \hat{H} | \psi_J \rangle$$

The solution of the above secular equation gives the energies of states I and J:

$$E_{I,J} = \frac{1}{2} \left[H_{II} + H_{JJ} \pm \sqrt{(H_{JJ} - H_{II})^2 + 4H_{IJ}H_{JI}} \right]$$

To satisfy the condition of degeneracy ($E_I = E_J$), the radicand must be equal to zero, meaning that the off-diagonal coupling terms (H_{IJ} and H_{JI}) must vanish and that H_{II} and H_{JJ} must be equal. Meeting those two conditions requires two independent nuclear coordinates.^{359,360} These are the GDV and DCV. This gives the state crossing a conical shape, as shown in Fig. 3.4 and A.2C. The correct description of the BP can be compromised if an electronic structure method does not correctly describe H_{IJ} and H_{JI} . Some methods cannot account for coupling at all (i.e. the coupling terms are zero at any possible geometry). In such cases, the BP gets reduced to just one dimension, and the crossing becomes linear rather than conical.¹⁸¹ A similar problem occurs in electronic structure methods that apply energy-scaling corrections to H_{II} and H_{JJ} but that

do not account for the effect of the scaling on the coupling terms. In such cases, the intersection is displaced to another region of the potential energy surface but the S_0 and S_1 wave functions and the couplings are not correctly described. In the cases described above, the S_0 and S_1 energy profiles along the CI loop yield a qualitatively different shape when compared to those of Fig. A.2D. In fact, instead of remaining non-degenerate along the entire loop, the two states will cross twice along the loop due to the fact that one of the BP vectors no longer splits the degeneracy. The energy difference between the two states will therefore drop to zero at two points along the loop (see examples in Fig. A.5). In conclusion, the CI loop distinguishes between methods that yield an unphysical one-dimensional linear crossing from methods yielding a BP with the correct dimensionality.

It is worth mentioning that in non-Hermitian theories, H_{IJ} and H_{JI} are not necessarily equal, introducing an additional condition (and change along an additional geometrical coordinate) for reaching the degeneracy. In such cases, the BP may become a three dimensional "branching space".^{361,362} The CI loop test cannot distinguish between methods that have a BP with a dimensionality larger than 2, such as in non-Hermitian methods. This is also true for cases where crossings between the potential energy surfaces of more than two electronic states occur.

Finally, notice that in the case of crossings between two electronic states of different spin multiplicities (e.g., singlet-triplet crossings), H_{IJ} and H_{JI} are identically zero in the framework of non-relativistic quantum chemistry and so the corresponding crossings are also always linear. Spin-orbit coupling, however, can couple states of different multiplicity and therefore lifts the degeneracy formally yielding a conical crossing.^{363,364}




Figure A.4. Energy profiles for electronic structure methods along the BLA path. MRCISD+Q is shown in all three panels for reference. A. Results for CASSCF, MRCISD, MRCISD+Q, and standard (unmodified) MR-PT2 all using an active space comprising 6 electrons in 6 π orbitals. B. Results for CASPT2 and MS-CASPT2 with an IPEA shift of 0.25, QD-NEVPT2 with an expanded active space (6 electrons in 8 orbitals), and XMCQDPT2 with a modified operator that incorporates some terms arising due to the non-separable part Γ_{ns} of the CASSCF state-averaged second-order density matrix Γ . C. Results for EOM-SF-CCSD(dT), ADC(2), SORCI, and TD-DFT using the mPW2PLYP functional. In all three panels, the energy values are relative to those of the *cis*-PSB3 equilibrium structure. The position of the states crossing for each method is indicated with a filled circle (except for that of QD-NEVPT2 which lies outside of the BLA range used). The curves are labeled at the left margin to distinguish between states with predominantly charge-transfer (Φ_{CT}) and diradical (Φ_{DIR}) character for each method. Adapted with permission from refs. 90,182,286,352. Copyright 2012 - 2014. American Chemical Society.




Figure A.5. The BLA and CI loop energy profiles for CASSCF (red, same as Fig. A.2.D bottom), MRCISD (grey), MRCISD+Q (black), SS-CASPT2(IPEA=0) (dark blue), MS-CASPT2(IPEA=0) (violet), SS-CASPT2(IPEA=0.25) (light blue), MS-CASPT2(IPEA=0.25) (violet), XMCQDPT2/ Γ_{ns} (green), QD-NEVPT2 (brown), and TDDFT/mPW2PLYP (orange). Top panel shows S_0 and S_1 energy profiles along the BLA scan, relative to *cis*-PSB3. The positions of the crossings are shown as filled circles. Due to the different geometries of the crossings at different levels of theory, several CI loops were constructed, each centered at a different BLA value, in order to incorporate the crossings of all tested methods. One loop was centered at the CASSCF crossing (panel A), one around the MRCISD crossing (panel B), one around the CASPT2(IPEA=0) crossing (panel C), and one around the MRCISD+Q crossing (panel D). Note that MRCISD+Q, SS-CASPT2(IPEA=0.25), MS-CASPT2 (IPEA=0.25), XMCQDPT2, QD-NEVPT2, and EOM-SF-CCSD(dT) all have similar energy profiles along the BLA and a similar geometry for the crossing, and therefore only one loop is used to test all six methods. Along the loops, only the energy differences between S_0 and S_1 are shown for all methods. An energy difference close to 0 kcal/mol means that the two states cross at that point. Such crossing points are indicated by black circles. Adapted with permission from ref. 182. Copyright 2014 American Chemical Society.

The PSB3 potential energy profiles along the BLA scan and CI loop computed with several electronic structure methods are shown in Fig. A.4 and Fig. A.5, respectively. These

benchmarks have also been extended to approximate linear-response coupled-cluster (CC2),³⁵⁴ and a range of DFT methods, including spin-restricted ensemble-referenced Kohn Sham (REKS).^{90,182,286,355} Furthermore, the quantum Monte Carlo (QMC) method has been benchmarked using the same system.²⁹⁵ Although QMC was originally a family of ground state methods, its extension to excited states has started somewhat recently. In fact, QMC methods for electronic excited states have started to show promising results in photochemical applications due to its favorable computational scaling and ability to account for electron correlation.³⁶⁵⁻³⁶⁷

A large family of methods that is missing from these benchmark studies are semi-empirical methods. This includes popular methods used for photochemistry including ZINDO/S,^{368,369} MNDO/MNDOC,^{370,371} excited-state variants of PM3,³⁷² AM1,³⁷³ and orthogonalization-corrected OMx methods (OM1, OM2, and OM3).^{374,375} Such semi-empirical methods have been used to study the retinal protonated Schiff base chromophore and its models.^{341,376} Extensive benchmarking has already been performed by Thiel and coworkers for different semi-empirical methods using PSB3³⁴³ as well as for other organic chromophores.³⁷⁷ The attractiveness of semi-empirical methods comes from their low computational cost compared to *ab initio* quantum chemistry methods, which allows them to be used in larger systems. However, this is often at the cost of reduced accuracy or a requirement that the method be re-parameterized for a particular system.

In the benchmark studies reviewed below, the Davidson corrected-MRCISD (MRCISD+Q) method is taken as the reference method due to its superior treatment of both static and dynamic contributions to the total electron correlation energy. To be more specific we recall that, while the Hartree-Fock (HF) method does account for the electronic correlation due to the exchange interactions between electrons of parallel spins (Fermi correlation), the Coulomb correlation is completely missing. Therefore, the formal definition of electron (Coulomb) correlation energy is:³⁷⁸

$$E_{corr} = E_{exact} - E_{HF}$$

Where E_{exact} is the exact energy. The exact energy can in theory be obtained at the full configuration interaction (FCI) level of theory, which is characterized by an electronic wave function incorporating all electronic excitations of all orders with respect to the closed-shell electronic configuration. E_{HF} is the Hartree-Fock (HF) energy, which contains only a single closed-shell electronic configuration. Static electron correlation becomes very significant when a state has important contributions from more than one degenerate or nearly-degenerate electron configuration. In such cases, the HF wave function is qualitatively wrong because it only accounts for one configuration (since it is described by a single closed-shell Slater determinant). Static electron correlation can be accounted for by including the few specific electronic configurations that have major contributions to the wave function. For example, electron correlation would play an important role for Φ_{DIR} in PSB3, which involves two electronic configurations that need to be treated on an equal footing (Fig. A.3) as they become nearly degenerate during homolytic bond-breaking. Dynamical

electron correlation, on the other hand, is related to the instantaneous interaction between electron pairs, and can only be accounted for by including many configurations, even those with relatively small weights, in the electronic wave function. Dynamic electron correlation would play an important role in both Φ_{DIR} and Φ_{CT} , but it plays a bigger role in Φ_{CT} due to the fact that charge transfer involves a larger rearrangement of electronic density and therefore a larger perturbation with respect to the reference configuration. The different contribution of the dynamical electron correlation to the Φ_{DIR} -dominated and Φ_{CT} -dominated regions of the chromophore potential energy surface provides an important example of the differential electron correlation effect. This effect can modify both the shape and barriers on a single potential energy surface, or the energy gap between different potential energy surfaces (e.g., the excitation energies). This means that an unbalanced description of electron correlation could result in an incorrect description of barriers, potential energy surface shapes, gaps, and crossings. In the following points i-vii we briefly describe electronic structure methods benchmarked using PSB3, and the results of their benchmarks along the BLA and CI loop paths. Note that methods i and iii in particular are popular for the description of isomerization of π -conjugated systems, and are mentioned in several instances throughout the review.

(i) The Multiconfigurational complete-active-space self-consistent-field (CASSCF) method is an elegant solution to the missing static electron correlation. It allows the user to select certain electrons and orbitals, termed the "active space," and exactly accounts for their correlation with each other. However, the remaining electrons are left uncorrelated. The active space therefore must be selected carefully to include electrons and orbitals that are considered necessary for the description of the spectroscopic or reactivity properties under investigation. In other words, the selection of the active space is a function of the chemical/physical problem investigated. If a suitable active space is chosen, the CASSCF method should satisfactorily account for static electron correlation. However, the dynamic component is still missed (both for inactive electrons and partly for active space electrons because they are not excited to virtual orbitals).

In the BLA scan (Fig. A.4A), this missing dynamic correlation causes CASSCF to have an unbalanced description of the S_0 and S_1 energy surfaces yielding a destabilized charge-transfer state when compared to the MRCISD+Q result. On the other hand, CASSCF is able to correctly describe the shape of the potential energy surface along the BP, and passes the CI loop test (Fig. A.5A).

(ii) Multireference configuration interaction (MRCI) is a variational approach that uses the same principle as truncated configuration interaction methods, but instead of using HF as the reference zero-order wave function it uses a multi-configurational reference such as the CASSCF wave function. Thus, MRCISD (SD stands for single and double) would expand the reference CASSCF wave function by adding single and double excitations. Since the CASSCF wave function already includes the electron correlation within the active

space, MRCI effectively introduces correlation of the electrons outside of the active space. Note that MRCISD is not rigorously size extensive, meaning that its accuracy does not necessarily scale linearly with system size. A simple energy correction, the Davidson correction,³⁷⁹ can be applied on top of MRCISD to estimate the energy of configuration interaction up to quadruple (indicated with the letter Q) excitations with almost no additional computational cost. This correction also improves the size extensivity of MRCISD. This leads to the MRCISD+Q energy that we use as a reference above, which is considered to yield values close to exact energies obtained with the unaffordable FCI method. For the benchmark studies reviewed here, MRCISD+Q serves as the reference method to measure the performance of other methods discussed here. However, MRCISD+Q is not used to study photobiological chromophores because it is too expensive for systems larger than PSB3.

The inclusion of dynamical electron correlation causes significant changes in the S_0 and S_1 energies along the BLA scan. Compared to CASSCF, both curves are stabilized with respect to the reference *cis*-PSB3 energy, but the change is disproportionate with the Φ_{CT} -dominated state stabilized more than Φ_{DIR} -dominated state. This results in a shift in the geometry of the crossing (CI_{BLA}) along the BLA coordinate, and, in turn, to a change in the entire topography of the S_0 potential energy surface, as shown in Fig. A.6. By *topography* we refer to the relative tilt, pinch, and asymmetry of the surfaces forming the CI, which tells us about the shape of the intersection itself.³⁵⁸ The CASSCF CI_{BLA} is peaked, and therefore the ground-state potential energy surfaces hosts two different transition states mediating *cis-trans* isomerization. On the other hand, CI_{BLA} in MRCISD+Q has a sloped topography, and therefore its ground state only hosts one transition state that mediates isomerization (TS_{CT}). Here we distinguish between *topography* and *topology*, the latter of which refers to the dimensionality of the corresponding BP and IS (which determines if the crossing is a conical or a linear crossing). MRCISD+Q does not yield a correct topology of the BP as the S_0 and S_1 curves cross twice along the CI loop (Fig. A.5D). This is due to the Davidson correction, which is a simple energy correction term not correcting the electronic state coupling. MRCISD, on the other hand, is able to describe the BP correctly like CASSCF (Fig. A.5C).




Figure A.6. Topography of the CI region located on the S_0 potential energy surface. Schematic representation of the topography of the S_0 CASSCF potential energy surface derived from the results reported in Fig. A.4 and Fig. A.5. B. The same representations for the MRCISD+Q results. Adapted with permission from ref. ⁹⁰. Copyright 2012 American Chemical Society.

(iii) Similar to MRCI, multireference second order perturbation theory (MR-PT2) approaches employ a multiconfigurational reference zero-order wave function. However, they account for dynamic electron correlation perturbatively instead of using a variational scheme. This makes MR-PT2 methods computationally less expensive than MRCISD, although they may not be as reliable. The CASPT2,³⁸⁰ MS-CASPT2,³⁸¹ XMCQDPT2,¹¹⁸ and QD-NEVPT2^{277,382,383} are all examples of MR-PT2 methods based on a CASSCF zero-order wave function. However, due to their different formulations, they behave differently along the BLA coordinate. As shown in Fig. A.4A, CASPT2 and MS-CASPT2 overstabilize the Φ_{DIR} -dominated state, QD-NEVPT2 overstabilizes the Φ_{CT} -dominated state, while XMCQDPT2 slightly overstabilizes both states (the Φ_{DIR} state more than the Φ_{CT} state) with respect to MRCISD+Q. The performance of all four methods could be improved by introducing small modifications (see Fig. A.4B). For instance, the use of a shift that modifies the energies of the active orbitals in CASPT2 and MS-CASPT2 (the IPEA shift³⁸⁴) significantly improves the agreement with MRCISD+Q. For QD-NEVPT2, a similar improvement is obtained by enlarging the active space of the reference CASSCF wave function by including polarizing π^* -orbitals.^{90,385} Finally, the agreement of XMCQDPT2 could be improved by using a modified operator that incorporates terms arising due to the non-separable part Γ_{ms} of the CASSCF state-averaged second-order density matrix Γ .⁹⁰ Overall, MR-PT2 methods show a significant improvement over CASSCF. However, it is difficult to know if the modifications mentioned above are generally applicable. For instance, the use of the IPEA shift in CASPT2 is a topic of active debate in the computational chemistry community.^{386,387} Also note that while CASSCF is size-extensive (if certain conditions are met), and perturbation theory is formally also size-extensive, MR-PT2 is not rigorously size-extensive.³⁸⁸

Despite their drawbacks, MR-PT2 methods are widely used in computational photochemistry and photobiology because of their affordable cost coupled with the ability to capture a large portion of electron correlation.³⁸⁹ Unfortunately, MR-PT2 geometry optimizations are still not practical for large systems (e.g., for biological chromophores discussed in this review). For this reason a common protocol has been to obtain CASSCF equilibrium structures and then compute the potential energies via MR-PT2 calculations. Such MR-PT2//CASSCF protocols have been widely employed for evaluating the energies of QM/MM models, as discussed in the next section of this appendix.

The CI loop results show that both the regular CASPT2 (SS-CASPT2) and MS-CASPT2 display artifacts near a CI point. CASPT2 suffers from its inability to describe the derivative coupling, and therefore yields a linear crossing. MS-CASPT2 corrects for this, but behaves erratically when the underlying CASSCF wave function has degeneracies.^{90,118,182} MCQDPT2 and NEVPT2 had similar problems, but were remedied using the extended version of MCQDPT2 (XMCQDPT2¹¹⁸) and the quasi-degenerate implementation of NEVPT2 (QD-NEVPT2) respectively.^{277,382,383} An extended implementation of MS-

CASPT2, termed XMS-CASPT2 been developed²⁷⁹ and would be expected to behave correctly in the vicinity of intersections.

(iv) Spectroscopy Oriented Configuration Interaction (SORCI) is effectively a multireference method that combines elements of MRCI and MRPT2 to achieve a balance between accuracy and affordability in a black-box framework.³⁹⁰ SORCI uses several mathematical approximations such as the resolution of identity and different levels of user-controlled truncations. While SORCI has many of the benefits of multireference methods and can calculate ground and excited states on an equal footing, it is aimed at vertical energy difference calculations (hence the “spectroscopically oriented” title) and not for potential energy surface mapping, particularly because the use of cutoffs results in an unsmooth surface.

Nonetheless, SORCI was tested along the BLA path (Fig. A.4C) and found to be capable of producing relatively smooth profiles, with the exception of regions in close proximity to the CI. These artifacts near the CI occur due to the Davidson correction³⁷⁹ used within SORCI.³⁹⁰ SORCI captures the relative energies of the Φ_{DIR} and Φ_{CT} states correctly with respect to each other, and therefore gives a correct geometry for CI_{BLA} . However, both states are over-stabilized relative to the *cis*-PSB3 geometry.³⁵³ While SORCI was not tested along the CI loop, it would not be expected to yield a conical crossing due to its use of the Davidson correction similar to MRCISD+Q.

(v) Equation-of-motion coupled-cluster (EOM-CC) is a multi-state extension of coupled-cluster that allows the description of a variety of target states. Coupled-cluster methods are currently among the most robust approaches for solving electronic structure problems. The exponential ansatz used for coupled-cluster wave functions ensures size extensivity (i.e., ensures that the quality of the calculation does not deteriorate with increasing size of the system, which is a common problem encountered in some electronic structure methods). Furthermore, these methods are relatively black box and can be systematically improved by increasing the level of excitations. The EOM-CC extension to coupled-cluster allows the description of a variety of target states in a single-reference formalism (unlike multireference methods such as MRCI and MR-PT2 that employ a multiconfigurational zero-order wave function and then introduce dynamical electron correlation in a second step). The choice of excitation operator used in EOM-CC allows this method to be used to describe different kinds of target states (see Fig. A.7).³⁹¹ For instance, different flavors of EOM-CC allow the calculation of excited states (EOM-EE),³⁹² ionized states (EOM-IP),³⁹³ electron-attached states (EOM-EA),³⁹⁴ or spin-flip states (EOM-SF). The spin-flip approach is a powerful way to treat closed-shell and open-shell, as well as doubly-excited configurations, on an equal footing if they can all be described as a single excitation from a reference triplet state.³⁹⁵⁻³⁹⁸




Figure A.7. Some of the target determinants generated by the singles and doubles EOM approach using an ionization potential (IP) operator, electron attachment (EA) operator, excitation energy (EE) operator, and a spin-flipping (SF) excitation operator acting on different reference configurations. *i, j* refer to occupied orbitals in the reference, *a, b* refer to unoccupied orbitals, and Ψ_0 shows the reference configuration for each operator. Adapted with permission from ref. ³⁵². Copyright 2013 American Chemical Society.

In ref. ³⁵² a number of EOM-CC approaches were tested, including both EOM-EE and EOM-SF. In Fig. A.4C we show the most successful, which was able to reproduce MRCISD+Q potential energy profile closely. This required using the spin-flip variant of EOM-CC, and the inclusion of perturbative triples (dT) corrections on top of single and double excitations. However, the EOM-SF-CCSD(dT) approach was unable to describe the topology of the BP near CI_{BLA} correctly (Fig. A.5D). This is most likely due to the triples correction, which employs EOM-SF-CCSD as a reference and corrects the energy to account for triple excitations and therefore behaves similarly to any other energy correction term (e.g., the Davidson correction in MRCISD+Q), yielding an incorrect crossing topology. Unperturbed EOM-SF-CCSD, on the other hand, would be expected to yield a BP spanning three dimensions, due to the non-hermiticity of the coupled-cluster ansatz.^{361,362}

(vi) Algebraic-Diagrammatic-Construction (ADC). The ADC method is a polarization propagator approach based on many-body Green’s function theory.³⁹⁹⁻⁴⁰¹ It describes the effect of an external perturbation on electronic structure using Møller-Plesset perturbation theory. The ground state therefore is described at the corresponding level of Møller-Plesset theory (i.e., MP2 for ADC(2) and MP3 for ADC(3)). Fig. A.4C shows that ADC(2) gives energies along the BLA that are in very good agreement with MRCISD+Q. However, because of the formulation of ADC methods, no coupling exists between the ground MP and excited ADC states, yielding a linear crossing.³⁵⁴

(vii) Time-dependent density function theory (TD-DFT). TD-DFT is one of the most widely used approaches in spectroscopy. Its most attractive feature is its ability to account for electron correlation and its relatively low computational cost scaling, which allows it to be used in much larger systems than would be possible with any of the electronic structure methods discussed above. However, there are several drawbacks for the use of TD-DFT in photochemistry. There are a number of well-documented problems with traditional TD-DFT methods,^{333,402} and particularly in the description of charge-transfer states,⁴⁰³ bond-breaking,⁴⁰⁴ and in being able to describe in a

balanced way states with double-excitation character.⁴⁰⁵ Some of these problems are mitigated by using hybrid or long-range separated functionals (to address problems with describing charge-transfer states) or double-hybrid functionals (to better describe states with double excitation). However, due to the large number of different DFT functionals being developed, it is not clear from the outset which functional is best for a given problem. A growing number of benchmarks have helped to identify the functional more suitable for a given class of systems or applications.^{405,406}

Huix Rotllant et al. have also benchmarked a number of different DFT functionals along the BLA path of PSB3, including hybrid functionals, hybrid meta-GGA functionals, range-separated hybrid functionals, and double-hybrid functionals.²⁸⁶

It was found that the relative energies of the Φ_{CT} and Φ_{DIR} states are sensitive to the type of functional, and the choice of functional is not always obvious. In Fig. A.4C we show only one example corresponding to the double-hybrid mPW2PLYP functional. Similarly, Casida, Filippi, and co-workers have benchmarked TD-DFT along torsional coordinates (both double-bond and single-bond isomerization) in reduced rPSB11 models and found that different functionals may yield a qualitatively different potential energy surface along these coordinates.³⁴⁷

A second problem with TD-DFT is that it is effectively a single-reference method, which may result in the calculation not converging in regions of degeneracy or near-degeneracy. In the case of the BLA scan, linear-response TD-DFT would often break down near the intersection of the Φ_{CT} and Φ_{DIR} states. This is only somewhat mitigated by using a double-hybrid functional.

A third problem with traditional TD-DFT approaches is their inability to account for coupling between the S_0 and S_1 state, resulting in a linear crossing (as revealed by the CI loop in Fig. A.5A) instead of a conical crossing.¹⁸² Such artifacts in the case of TD-DFT has been explored in detail by Levine, Martínez and co-workers.¹⁸¹

Despite these drawbacks, TD-DFT remains a powerful approach in photochemistry and photobiology, particularly for calculating spectroscopic properties and even excited-state processes. Attempts are being made to extend DFT methods to be able to treat non-adiabatic regions of the potential energy surface correctly. Examples of this include configuration-interaction corrected Tamm-Dancoff approximation (CIC-TDA) TD-DFT,²⁸³ constrained density functional theory – configuration interaction (CDFT-CI), and the state-interaction state-averaged spin-restricted ensemble-referenced Kohn – Sham (SI-SA-REKS) approach based on ensemble DFT.^{286,407} Another very promising approach is the application of the spin-flip method to TD-DFT.⁴⁰⁸ SF-TD-DFT is capable of yielding the correct 2D topology of the branching space around the conical intersection^{182,287} However, SF-TD-DFT often suffers from severe problems with spin-contamination, especially far from the Franck-Condon region. A spin-adapted variant of SF-TD-DFT (SA-SF-TD-DFT) has recently been

developed that resolves the issue with spin-contamination. Development of analytical gradients may allow this method to be used affordably to run *ab initio* excited-state or multi-state molecular dynamics.²⁸⁸

As interest in photochemical and photobiological problems continues to grow, there is increasing demand for accurate yet affordable electronic structure methods that can treat extended regions of the ground and excited-state potential energy surfaces. Electronic structure methods have made considerable progress in recent years. However, remarkably, classical methods that have existed for over 20 years, particularly those building on multi-configurational methods and coupled-cluster approaches, remain among the most widely used *ab initio* methods for photobiology today. While coupled cluster methods are mainly used for highly accurate studies of small molecules, it has been possible to employ multiconfigurational and multireference approaches to larger systems such as biological chromophores, particularly through the use of the MR-PT2//CASSCF protocol where geometries are optimized at the CASSCF level of theory and energies obtained at the MR-PT2 level of theory.³⁸⁹ Many of the studies we discuss in the present review employ the MR-PT2//CASSCF method. The reason is that CASSCF can capture multi-configurational character, which is often a necessity for describing photochemical processes such as bond-breaking and isomerization, and can also capture the correct topology of conical intersections, allowing it to be used for non-adiabatic trajectory calculations. At the same time, to capture energetics correctly it is necessary to account for dynamical electron correlation, which can be achieved relatively affordably with MR-PT2.

On the other hand, while the CASPT2//CASSCF protocol is used routinely for systems such as rPSB11 and chromophores of similar size, it requires the user to be knowledgeable about the drawbacks of each of CASSCF and CASPT2, and therefore is certainly not a black-box approach. This protocol may also become too expensive when looking at larger chromophores, in particular those with more electrons that need to be included in the active space. In particular, this might apply for chromophores with a larger π -conjugated framework, or with heteroatoms with lone pairs that may be involved in the photochemistry. Since the computational cost of CASSCF scales factorially with the size of active space, large active spaces become rapidly intractable. It is worth mentioning here two families of methods that are being developed to overcome this shortcoming. The first approach is the restricted active space (RAS) method²⁹⁹ that divides the total active space into three subspaces. One subspace is similar to CAS and allows all possible excitations, while the other two are typically restricted to single and double excitations from one subspace to another. A more general implementation was achieved by developing the general active space (GAS) method where the number of such spaces is arbitrary.³⁰⁰ The second strategy is based on the Density Matrix Renormalization Group (DMRG) in combination with CASSCF.⁴⁰⁹ This method allows an active space of up to 50 electrons and orbitals. Several programs are already available to do such calculations, e.g. BLOCK by the Chan group,^{410,411} QCMAQUIS by the Reiher group,^{412,413} and CheMPS2 by Wouters et al.⁴¹⁴ Furthermore, DMRG can be combined with MR-PT2 methods such as NEVPT2 or

CASPT2 to yield more quantitative results.^{301,302,415} However, while a large size active space is becoming feasible in a foreseeable future this also means that the selection of a relevant active space will be more difficult. For this purpose, the Reiher group has developed an algorithm that takes over the task for the user.⁴¹⁶

A.1.2. Hybrid Quantum Mechanics/Molecular Mechanics Approach.

The computational investigation of light-responsive proteins necessitates an approach that can treat the chromophore in the excited state and account for the protein and/or solvent environment as well. This is most commonly achieved by using a QM/MM model.⁴¹⁷ The QM/MM approach belongs to a family of multi-scale methods, which are suitable for the investigation of the properties of a large molecular system. Multi-scale methods employ two or more levels of theory with distinct accuracy and computational cost. Specifically, in QM/MM, the use of an accurate, but relatively expensive, quantum mechanical (QM) level of theory for treating the group involved in electronic excitation and less accurate, but affordable, molecular mechanical (MM) approach for the remaining larger part makes it feasible to calculate protein-chromophore complexes. The key steps in constructing a basic two level QM/MM model are: (i) partitioning the protein into QM and MM subsystems and (ii) choosing the scheme that describes the interaction (i.e., coupling) between the two subsystems. Moreover, a special situation arises in cases where the boundary between the QM and MM subsystems is located at one or more covalent bonds, as in the example of rhodopsin mentioned in the introduction section.⁴¹⁸ In such cases, a suitable scheme describing the covalent link between the two subsystems needs to be adopted as well.

QM/MM models of photoreceptor proteins have reached an evident level of maturity, and are capable of treating the absorption and emission spectra of different types of proteins (e.g. with backbones dominated by α -helices and β -sheet structures) incorporating different chromophores (e.g. cationic, anionic and neutral chromophores).⁴⁰⁷ The CASPT2//CASSCF/MM approach in particular has been used to study a number of different photoactive systems as shown in Fig. A.8.




Figure A.8. Comparison between calculated and measured λ_{\max} values for various photoreceptor proteins (including fluorescent proteins). Calculated

values were obtained by employing QM/MM models based on *ab initio* multireference QM methods. The displayed data are collected in ref. ⁴⁰⁷ and in ref. ^{419,420} for GFP S85T/H148D. The vertical axis shows experimentally observed wavelengths and the horizontal axis shows the CASPT2//CASSCF/MM computed λ_{\max}^a ($S_0 \rightarrow S_1$, $S_0 \rightarrow S_2$) and λ_{\max}^f ($S_1 \rightarrow S_0$) values. The employed force fields vary from AMBER to CHARMM. The abbreviation of the proteins are bovine rhodopsin (Rh), bathorhodopsin (bathoRh), isorhodopsin (iso-Rh), the Rh mutants E113D and G121L, squid rhodopsin (sqRh), bacteriorhodopsin (bR), *Anabaena* sensory rhodopsin (ASR), human Melanopsin (hMeOp), Channel Rhodopsin (ChR2), green fluorescent protein (GFP), the GFP double mutant S85T/H148D, photoactive yellow protein (PYP), the azurin mutant (azurin48W), blue-light using FAD (BLUF), the tryptophan containing proteins parvalbumin (Parv) and monellin (Mone), firefly luciferase, the GFP-like mutant Dronpa, and phytochrome. Figure adapted and upgraded from ref. ⁴⁰⁷. Copyright 2014 The Royal Society of Chemistry.




Figure A.9. QM/MM scheme illustrated for the bovine rhodopsin. A. Bovine rhodopsin model incorporating a rhodopsin trimer embedded in a membrane (see for instance figure reproduced from ref. ⁴²¹. Copyright 2013 American Chemical Society). Only one rhodopsin monomer is a QM/MM model that contains QM and MM subsystems, while the other two rhodopsins are treated exclusively at the MM level. B. A constrained monomer rhodopsin model where only the QM chromophore, MM covalently bonded lysine side chain, and nearby side-chains of the protein binding pocket are free to move. In the example shown here, cavity side chains in red are defined as any residue that has any of its atoms within 4 Å from any atom of the chromophore at equilibrium geometry. Such QM/MM models are often used in cases where the membrane environment is not available and the whole protein cannot be left to relax (see for instance figure adapted from ref. ³³⁰. Copyright 2016 American Chemical Society).

A.1.2.1 QM/MM embedding schemes. Besides the accuracy of the electronic structure method used to describe the QM subsystem (see Section A.1.), the QM and MM subsystems coupling is of central importance. For a detailed discussion of different QM/MM embedding schemes, we refer the reader to ref. ⁴²²). There are three main coupling schemes:

- 1) Mechanical embedding: The QM subsystem does not “feel” the electrostatic field generated by the point charges of the MM subsystem. The only non-bonding interaction coupling the two subsystems is the short-range van der Waals interaction.
- 2) Electrostatic embedding: In addition to the mechanical embedding, the QM subsystem is also perturbed by the point charges of the MM subsystem. This interaction can lead to differential effect on different electronic states that might result in stabilization or destabilization of the ground and excited states as shown in Fig. 3.2. However, only the QM subsystem experiences an electrostatic effect. The MM subsystem is described by point charges

whose values are not affected by the QM subsystem electron density.

- 3) Polarizable embedding: For a mutual polarization between the QM and MM subsystems, the MM environment should also respond to the changes in the electron density of the QM region. This requires a MM subsystem that features a polarizable charge model.

Although the polarizable embedding scheme is, in principle, the most accurate, it is rarely used. The reason is that the few available polarizable force fields are still being benchmarked and developed.^{423,424} In practice, presently, the electrostatic embedding is the most commonly used QM/MM coupling scheme. Although recent attempts to use polarizable embedding have been reported, their application in the simulation of ultrafast dynamics is limited.⁴²⁵ The same holds also for QM/DFT embedding, which is only available for single point calculations at the moment.⁴²⁶ The inherent intricacy of excited-state dynamics studies is also the reason why complex QM/MM models (e.g., large membrane embedded photoreceptor models) are mostly used for calculations of vertical excitation energies.⁴²¹ In some cases, the order of the electronic states is perturbed so much by the electrostatic embedding that a subsequent molecular dynamics or reaction path mapping is not possible due to the presence of several intruder electronic states between the spectroscopic state and the ground state. If these intruder states are physical (i.e. they would be experimentally observed) one can, in principle, correctly account for them in a non-adiabatic trajectory computations (see Section A.I.3) or, when using an ad hoc surface hopping strategy,⁴²⁷ in a MEP computation. However, certain intruder electronic states are unphysical as they would be removed at a higher levels of theory (e.g. CASSCF generated intruder states may be removed at the CASPT2 level). In these cases, certain authors have switched to a simpler mechanical embedding that avoids these technical problems and allows to get a qualitative description of the system reactivity.²⁹ This is of course acceptable only when energy gradients computed at the higher level of theory are unaffordable.

A.I.2.2 QM/MM energy expression. In the first formulations of the QM/MM method^{417,428,429} the energy was calculated according to the formula:

$$E_{QM/MM}(S) = E_{QM}(I) + E_{MM}(O) + E_{QM-MM}(I, O)$$

where the energy of the “inner” subsystem (I) is evaluated at the QM level (E_{QM}) and the energy of the “outer” subsystem (O) is computed separately using an MM force field (E_{MM}). The coupling term between the inner and the outer systems E_{QM-MM} is added to the total energy. Therefore this energy expression is called *additive*. The terms included in the coupling are determined by the embedding scheme, as reviewed in A.I.2.1 above.

An alternative QM/MM energy expression was proposed and implemented by Morokuma and coworkers:⁴³⁰⁻⁴³³

$$E_{QM/MM}(S) = E_{QM}(I) + E_{MM}(S) - E_{MM}(I)$$

The three energy terms arise from one QM calculation of the inner system and two MM calculations of the total (inner+outer) system and the inner system, respectively. The last term is subtracted from the total MM energy thus eliminating

double counting. This QM/MM energy expression was coined *subtractive*. The main advantage of the subtractive scheme is the avoidance of explicitly calculating the QM-MM coupling term and, therefore, regular QM and MM programs can be used without any modifications. The first realization of the subtractive scheme was named Integrated Molecular Orbital/Molecular Mechanics (IMOMM).⁴³⁴ Further development extended this method to combine a high level QM method with a lower-level one (IMOMO),⁴³² eventually leading to a fully generalized multi-layer scheme called ONIOM.^{431,435}

A.I.2.3 Link atom scheme. Partitioning a molecular system into two subsystems is trivial for a solute surrounded by solvent molecules. In such a scenario there is no covalent bond that connects the QM and MM subsystems. However, in many photoreceptor proteins the chromophore is covalently bound to the apoprotein. Hence a QM/MM model has to deal with bonds at the boundary between the subsystems. In the following we discuss the case of Rh, where the rPSB11 chromophore is covalently bound to the Lys296 residue (see Fig. A.10). Since we are concerned with the photochemistry of rhodopsin, the QM subsystem needs to include the whole conjugated π -electron system. Hence, the boundary must be located on a bond of the Lys296 side-chain beyond the Schiff base functional group. The $C_{\delta}-C_{\epsilon}$ was selected to minimize the computational cost without altering significantly the electronic structure of the QM subsystem. However, the bond has to be cleaved homolytically or heterolytically leading to undesirable changes in electronic structure. Three different approaches have been proposed to overcome this problem: 1) link atom schemes⁴²⁹, 2) boundary atoms⁴³⁶ and 3) localized orbital schemes⁴³⁷. The studies presented in this review make use of the link-atom scheme exclusively, so we will limit our discussion to this particular scheme and refer the interested reader to the review by Senn and Thiel⁴²² for a description of the other two schemes.

In the link-atom scheme, a new atom is introduced to cap the cleaved bond. In most case it is a hydrogen atom, and the bond that is truncated is selected to be a single bond of small or negligible polarity (like the Lys296 $C_{\delta}-C_{\epsilon}$ bond in bovine rhodopsin). The introduction of a new atom that is not present in the real system introduces a few problems, however. One problem is the addition of three degrees of freedom that an atom has and that needs to be optimized in a geometry optimization or that can introduce some artifacts in trajectory computations. This disadvantage was resolved by constraining the bond between the QM frontier atom and the link atom to be aligned along the MM bond (see Fig. A.7) connecting the QM and MM subsystems.⁴³⁸ Such a constraint removes all additional degrees of freedom except the link atom bond length. Another caveat that results from the link atom scheme is the overpolarization of the link atom by the adjacent MM frontier atom. If not addressed, this could lead to large artifacts that even lead to an incorrect ordering of excited states. The most common solution to this problem is the charge-shifting scheme that was put forward by Sherwood and coworkers.⁴³⁹ In this approach, the charge of the closest MM frontier atom is distributed onto neighboring MM atoms bound to it. For instance, the C_{δ} charge (e.g., the AMBER point charge) of the Lys296 in Rh is set to zero and its value is evenly added to the remain-

ing MM atom of the Lys residue (leaving the total residue charge unchanged).




Figure A.10. A link atom scheme for Rh. The rPSB11 chromophore is covalently bound to the lysine residue Lys296 (in blue) embedded in the protein cavity (see also Fig. A.9B as a reference). The common partitioning puts the boundary on the C δ -C ϵ single bond. The hydrogen link atom (shown in orange) is placed along this bond.

A.I.2.4 Protocols for QM/MM model building. There are many ways to construct QM/MM models, so here we will discuss one example to illustrate how such a model construction could be done. A photoreceptor QM/MM model is often derived from a crystallographic or Nuclear Magnetic Resonance (NMR) structure deposited in the Protein Data Bank,⁴⁴⁰ or from a homology model. Usually, these structures do not include protons, may be missing a number of residues, and do not contain information on the state of ionizable residues or on the distribution of nearby counterions. Therefore, such missing parts must be incorporated into the model using some QM/MM model-building workflow (see Fig. A.11). This may require structural manipulations including the relaxation of the system after addition of the hydrogen to the crystal structure. This relaxation is usually carried out at the MM level of theory via geometry optimization and/or molecular dynamics (MD) simulations. There is no universal way to do this, and the result depends on the protocol, the software, and the parameterization of the chromophore used. This often results in different labs having slightly different models. Furthermore, to have a more complete model the entire protein may be placed in a solvent (or a membrane patch if it is a membrane protein). Finally, it may be important to add ions in the simulation to neutralize the charges of the protein. Once the classical model is completed, a minimization and a subsequent MD trajectory are often necessary. Upon equilibration, such a trajectory could be used to generate one single QM/MM model, or different snapshots taken at different times can be used as starting points to generate a series of QM/MM models. The QM/MM model(s) can then be optimized at the QM/MM level of theory and used for subsequent excitation energy, reaction path, or trajectory calculations.




Figure A.11. A sample workflow for the construction of light-responsive protein QM/MM models for the simulation of the S₁ population dynamics. The workflow is divided into four phases, with each phase broken down into necessary tasks. Figure adapted from ref. ⁴⁴¹. Copyright 2016 Elsevier.

Another option for generating a set of QM/MM models that can be used to run an ensemble of trajectory calculations is to use a sampling protocol to generate initial geometries and velocities according to a specific distribution such as the Wigner or the Boltzmann distribution. In this way, one has the opportunity to investigate, by running a group of trajectories in parallel, the dynamics of an entire population and even compute photoisomerization QYs. As discussed below, the accuracy of these computations will depend on the number of trajectories run. Due to the computational cost limitations, QM/MM trajectories computed at the CASSCF/MM level are usually limited to lifetimes that do not exceed a few picoseconds. However, this is enough for investigating the light-driven primary event in the ultrafast activation of biological photoreceptors.

Recently, there have been attempts to automate and, partially, standardize the building of QM/MM models for the purpose of comparison of mutants or of photoreceptors of the same family.³³⁰ Presently, these are monomer QM/MM models of retinal proteins whose construction only requires an experimental structure or homology model and information about the protonation states of the ionizable residues and ions that counterbalance the charges. The so called Automatic Rhodopsin Protocol (ARM) generates a QM/MM model consisting of a frozen protein environment with a flexible protein binding pocket formed by the side-chains of the chromophore cavity (i.e. consistent with the model in Fig. A.9B and A.10). The quality of the models has been assessed by comparing calculated and observed excitation energies for a series of rhodopsins from different organisms. As detailed in Section 8.2.4, it is shown that ARM models are capable of providing information on excitation energy trends.

A.I.3 Non-adiabatic molecular dynamics simulations and trajectory surface hopping. In the course of photoisomerization, a molecule progresses on the excited state potential energy surface(s) and ultimately decays to the S₀ state before completing the isomerization event or re-generating the reactant. The simulation of photoisomerization dynamics therefore requires the description of transitions between the potential energy surfaces associated with different electronic states, also referred to as hopping events. In the context of photoisomerization, such a transition occurs in the proximity of CIs, where

the adiabatic (Born-Oppenheimer) approximation breaks down.

Such non-adiabatic transitions can be described in several different ways, ranging from full quantum dynamics (where the nuclear motion is treated quantum mechanically) to semi-classical methods (where nuclei are treated classically but the surface hop is treated in a special way to account for non-adiabatic events).⁴⁴² Subotnik et al. have recently examined popular surface hopping algorithms and presented modern advances in such algorithms.⁴⁴³ Here, in the context of photo-biology, we will focus our discussion on three trajectory surface hopping (TSH) algorithms commonly used for systems the size of biological chromophores.

One approximate algorithm implemented by Robb and coworkers^{136,335,338,444} describes the coupling between two states, k and j , using the numerical approximation:

$$\langle \Psi_k(t_0) | \Psi_j(t_0 + \Delta t) \rangle = \langle \Psi_k | \frac{d}{dt} \Psi_j \rangle$$

where Ψ_k and Ψ_j are the time-dependent electronic wave function of the states, t_0 the initial time and Δt the time-step. The coupling is used to estimate the interaction of the two states and to decide if a hop between the states occurs (Fig. A.12). A transition between Ψ_k and Ψ_j occurs if a specified threshold is reached.




Figure A.12. Change in the adiabatic electronic wave functions for two consecutive steps of a semi-classical surface-hop trajectory calculation. The first time-point is marked by t_0 and the time-step by Δt . Ψ_k and Ψ_j are time dependent electronic wave function of the k and j states, while ϕ_k and ϕ_j denote stationary diabatic states (i.e. one can think of ϕ_k and ϕ_j as reference, time-independent electronic configurations whose mixing provide a description of the adiabatic states). The change in mixing coefficients at each time step, makes possible the description of the time evolution of the adiabatic states). Adopted from ref. ²⁷¹. Copyright 2015 American Chemical Society.

After a hopping event, the energy is conserved by correcting the momentum in the direction parallel to the derivative coupling vector.^{338,445,446}

The second is the Tully's fewest switches algorithm.⁴⁴⁷ This approach requires the coupled propagation of both the nuclei and the electrons motion. As for the previous method, the former is performed by integrating the Newton's equations of motion for the nuclei moving on a given potential energy surface on the femtosecond timescale, while the latter is carried out by integrating the time-dependent Schrödinger equation:

$$i\hbar\dot{c}_k = \sum_j c_j (V_{kj} - i\hbar\dot{\mathbf{R}}\mathbf{d}_{kj})$$

on the attosecond timescale. \mathbf{R} stands for the nuclear position (representing the molecule geometry), \mathbf{r} stands for the electronic coordinates, $\mathbf{d}_{kj} = \langle \Psi_k(\mathbf{r}; \mathbf{R}) | \nabla_{\mathbf{R}} \Psi_j(\mathbf{r}; \mathbf{R}) \rangle$ are the derivative coupling terms, and $V_{kj} = \langle \Psi_k(\mathbf{r}; \mathbf{R}) | \mathbf{H}(\mathbf{r}; \mathbf{R}) \Psi_j(\mathbf{r}; \mathbf{R}) \rangle$ are the electronic Hamiltonian elements for a given geometry \mathbf{R} . The wave function $\Psi(\mathbf{r}, \mathbf{R}, t)$, is expanded as a linear combination of stationary states $\phi_j(\mathbf{r}; \mathbf{R})$ yielding the time dependent expression $\Psi(\mathbf{r}, \mathbf{R}, t) = \sum_j c_j(t) \phi_j(\mathbf{r}; \mathbf{R})$. The \mathbf{d}_{kj} terms are key parameters for inducing the surface hopping as they promote the mixing of the electronic states.

The equation above is integrated by computing the $\dot{\mathbf{R}}\mathbf{d}_{kj}$ terms. By applying the chain rule, the gradient of the wave function is transformed into a wave function time-derivative yielding $\dot{\mathbf{R}} \cdot \mathbf{d}_{kj}(\mathbf{R}) = \langle \Psi_k | \frac{\partial \Psi_j}{\partial t} \rangle \approx \langle \Psi_k | \frac{\Delta \Psi_j}{\Delta t} \rangle$. These terms can be calculated analytically, but there is also a possibility to determine them numerically along the trajectory. It is required that at each step of the nuclear motion the integration step defined by Δt coincides with the exact calculation of the wave function. However, the time-step of the electronic integration is typically at least one or two orders of magnitude smaller than Δt . Therefore, to get accurate results in the integration of the time-dependent Schrödinger equation, an interpolation between consecutive values of $\dot{\mathbf{R}} \cdot \mathbf{d}_{kj}(\mathbf{R})$ is required. An interpolation/extrapolation scheme for the determination of time-derivative coupling terms has been put forward by Hammes-Schiffer and Tully.⁴⁴⁸ Additionally, in order to preserve the consistency of the population density matrix a decoherence correction proposed by Granucci and Persico⁴⁴⁹ is required.

By integrating the equation above one obtains, for each time step, the populations of each electronic state $|c_i|^2$. Depending on the time variation of these populations, the fewest switches algorithm⁴⁴⁷ stochastically decides whether the trajectory should continue to propagate on the same potential energy surface (e.g. the S_1 potential energy surface) or should jump onto another one (e.g. to the S_0 potential energy surface). After the hop, the next energy gradient for the nuclear motion is calculated using the potential energy surface associated to the new state while the nuclear velocities are scaled to preserve the total energy.

The third algorithm is based on the Zhu-Nakamura theory.⁴⁵⁰ The surface hopping approach based on the Zhu-Nakamura theory provides a way to compute the probability for a non-adiabatic transition for a wide spectrum of energy gaps.⁴⁵¹ This approach is computational feasible and straightforward to implement because it can be also applied if the derivative coupling terms are not available.⁴⁵² However, it has several important advantages compared to some of the other surface hopping approaches, such as the treatment of surface hops that are energetically forbidden. The Zhu-Nakamura surface hopping scheme has been applied to several one-dimensional problems before it was recently extended for multi-dimensional problems.⁴⁵³ The key step in the Zhu-Nakamura approach is based on the assumption that hops can take place in the close vicinity to a CI and that the derivative coupling direction can be approximated by the eigenvector of the Hes-

sian of the potential energy surface driving the dynamics $\partial^2 \Delta V / \partial R_i \partial R_j$ corresponding to its maximum eigenvalue.

As mentioned earlier, the surface-hopping methods presented above belong to the group of semi-classical methods where the electrons are treated quantum mechanically and the nuclei are described classically. However, due to the breakdown of the Born-Oppenheimer approximation in the vicinity of conical intersections, the quantum nature of the nuclear motion can become important. For example, quantum effects such as quantum decoherence, zero-point energy effects, tunneling, geometric phase effects, and superexchange cannot be described classically. As a result, there have been numerous attempts at extending classical (e.g., surface-hopping) approaches to account for some quantum effects (e.g., see refs. ⁴⁵⁴⁻⁴⁵⁸).

While semi-classical approaches are affordable and work well for many types of problems, their extension to treating quantum effects is not without difficulty. Therefore, full-dimensional quantum wavepacket propagation methods are still widely applied to accurately study dynamics of small molecules with relatively few degrees of freedom. Examples of such approaches includes multi-configuration time-dependent Hartree (MCTDH) and variational multiconfiguration Gaussian wavepackets (vMCG).^{459,460} (see ref. ⁴⁶¹ for a discussion of these methods). However, such methods are not usually applied for systems the size of biological chromophores reviewed in this work due to the exponential growth of their computational cost with increasing number of degrees of nuclear freedom. However, progress in this direction, such as the development of multi-layer techniques,⁴⁶² is allowing MCTDH to be applicable to larger systems. For example, multi-layer MCTDH was recently applied to several hundreds of degrees of freedom to study the FMO complex.⁴⁶³ Similar advancements have been reported for the vMCG method.⁴⁶⁴

One approach that combines elements of semi-classical and quantum mechanics is the *ab-initio* Multiple-Spawning (AIMS) approach developed by Martínez and co-workers.^{465,466} In this method, a set of frozen Gaussians, simulating a nuclear vibrational wavepacket, follow classical trajectories while their expansion coefficients are propagated according to the time-dependent Schrödinger equation. Non-adiabatic transitions are then described by expanding the basis set of the Gaussian functions after the transition leading to "spawning" of new populations that may either continue to evolve on the excited state or decay to the ground state. The AIMS approach improves on semi-classical methods by describing non-adiabatic transitions quantum mechanically, including coherence and decoherence effects. However, the AIMS computational cost may increase in cases of many non-adiabatic transitions (i.e., depending on the topology of the potential energy surface) and can become intractable for extended molecular systems. One workaround for the rapidly increasing number of spawned function is stochastic selection⁴⁶⁷ where the number of basis function can be reduced for efficiency. In general, the AIMS method has been constantly extended over the last decade by introducing full *ab initio* spawning that is free from any empirical parameters⁴⁶⁶ and recently including the treatment of intersystem crossings.⁴⁶⁸

APPENDIX II

A.II.1 Relationship between reaction coordinate and intersection space. The MEP coordinate described in Section A.I.1 is expected to operate in all rPSB chromophores. Since the MEP is dominated by three modes (i.e. BLA, HOOP and torsional coordinate), it cannot be realistically visualized in 2D. This has forced us to illustrate the relationship between S_1 MEPs, S_0 MEPs and the CI only schematically as seen in Fig. A.1C. In such 2D representation, the relationship between an IS segment (containing the CI_{CIS} , CI_{BLA} and CI_{TRANS}) is puzzling. In fact, the IS appears parallel to the mixed torsion and HOOP mode usually associated to the isomerization coordinate. In order to clarify the relationship between S_1 MEPs and the IS, in this section we present such a detailed mapping for the case of the minimal three double-bond model of rPSB, PSB3 introduced in Section A.I.1.

It has been possible to map the evolution of PSB3 along its S_1 MEPs in a 3D coordinate space.⁶² The 3D schematic representation of the relationship between the BLA, HOOP and torsional coordinate change in PSB3 is given in In Fig. A.13 for both *cis* to *trans* (starting from $PSB3_{CIS}$) and *trans* to *cis* (starting from $PSB3_{TRANS}$) isomerization of the central $C2=C3$ double bond. In the same figure we show that the segments of the S_1 MEPs hitting the CIs is a combination of skeletal double-bond torsion and HOOP while the BLA value remains substantially constant in this region of the MEP. On the other hand, a change in the BLA value would lift the S_1/S_0 energy degeneracy at the CI. Therefore, the MEP and BLA coordinates have components belonging to the two-dimensional BP seen in Fig. A.2. In contrast, a combination of skeletal double-bond torsion and HOOP opposite to the MEP hitting the CI would span the IS curve. As anticipated in Fig. 2.3 and related text of the Introduction section, the IS a N-2 dimensional space (i.e. a hyperline with respect to the N-dimensional space of the full set of independent vibrational degrees of freedom of the system). The IS is orthogonal to the BP and consists entirely of CI points. S_1 MEPs launched from different stereoisomers reach different CIs along the IS, each mediating different relaxations. This is illustrated in Fig. A.13 where excited-state MEP calculations launched from $PSB3_{CIS}$ and $PSB3_{TRANS}$ reach the distinct CI_{TRANS} and CI_{CIS} located along the IS. CI_{TRANS} and CI_{CIS} are not minimum energy CI points (i.e. energy minima along the IS). In fact, the search for a minimum energy CI⁶² leads to CI_{BLA} , which is located between CI_{TRANS} and CI_{CIS} and mainly characterized by a different value of the HOOP coordinate. More specifically, CI_{BLA} features the carbon atoms of the isomerizing double bond in a substantially sp^2 -hybridized state and, therefore, has a nearly zero HOOP value. Note that Fig. A.13 is a representation of the MEP_{CIS} , MEP_{TRANS} , and CI seam pathways displayed in Fig. A.1, although in coordinate space.




Figure A.13. Schematic representation of the geometric progression of PSB3 driven by the S_1 force field. Full circles indicate the position of S_0 energy minima and CIs. Full black lines represent S_1 MEPs while the red line represents the S_1/S_0 IS. Dashed black lines are just projections of MEPcis on the different places. The arrows indicate the direction of the force along the MEPs. The progression mainly occurs along the 3D configuration space defined by the BLA, HOOP and C2=C3 double-bond torsion. Such space contains a cross-section of the S_1/S_0 intersection space (IS) running along the HOOP mode coupled with smaller components of the other modes. The geometry and π -overlap change in the region of the isomerizing double bond is indicated using Lewis formulas and p-orbital diagrams. With permission from ref. ⁶². Copyright 2013 American Chemical Society.

The geometrical progression described by the MEP is consistent with that of the S_1 trajectory of Fig. 2.3. More specifically (follow, for instance, the PSB3_{CIS} progression towards CI_{CIS}) the geometrical changes are driven by the need to rapidly decrease the π -overlap across the isomerizing double bond. This happens because, in the FC region, S_1 has a Φ_{CT} character setting the orbital interaction across the bond into an anti-bonding state. Therefore, the fastest and most effective way to decrease the π -overlap is the expansion of the C2-C3 bond length. This is immediately followed by the HOOP deformation and finally by the skeletal twisting. The initial change in HOOP has the effect of changing the hybridization status of the two carbon atoms of the isomerizing double bond. This is the main reason why at both CI_{TRANS} and CI_{CIS} such carbons are largely sp^3 hybridized. The hybridization then changes as one moves to CI_{BLA} where the same atoms have sp^2 hybridization again.

AUTHOR INFORMATION

Corresponding Author

* massimo.olivucci@unisi.it, molivuc@bgsu.edu

Present Addresses

††Nanoscience Center and Department of Chemistry, University of Jyväskylä, P. O. Box 35, 40014 Jyväskylä, Finland

Author Contributions

The manuscript was written through contributions of all authors who have given approval to the final version of the manuscript.

Funding Sources

This work has been in part supported by NSF (CHE-1551416, CHE-1710191), MIUR (PRIN 2015), HFSP (RGP0049/2012) and the USIAS fellowship 2015 of the University of Strasbourg, France.

ACKNOWLEDGMENT

We thank Gerrit Groenhof for many useful suggestions.

REFERENCES

- (1) Ernst, O. P.; Lodowski, D. T.; Elstner, M.; Hegemann, P.; Brown, L. S.; Kandori, H. Microbial and Animal Rhodopsins: Structures, Functions, and Molecular Mechanisms. *Chem. Rev.* **2014**, *114*, 126-163.
- (2) Christie, J. M.; Blackwood, L.; Petersen, J.; Sullivan, S. Plant Flavoprotein Photoreceptors. *Plant. Cell. Physiol.* **2014**, *56*, 401-413.
- (3) Bibikov, S. I.; Grishanin, R. N.; Kaulen, A. D.; Marwan, W.; Oesterhelt, D.; Skulachev, V. P. Bacteriorhodopsin Is Involved in Halobacterial Photoreception. *Proc. Natl. Acad. Sci. U. S. A.* **1993**, *90*, 9446-9450.
- (4) Jung, K. H.; Trivedi, V. D.; Spudich, J. L. Demonstration of a Sensory Rhodopsin in Eubacteria. *Mol. Microbiol.* **2003**, *47*, 1513-1522.
- (5) Kandori, H. Ion-pumping Microbial Rhodopsins. *Front. Mol. Biosci.* **2015**, *2*, 52.
- (6) Nagel, G.; Szellas, T.; Huhn, W.; Kateriya, S.; Adeishvili, N.; Berthold, P.; Ollig, D.; Hegemann, P.; Bamberg, E. Channelrhodopsin-2, a Directly Light-gated Cation-selective Membrane Channel. *Proc. Natl. Acad. Sci. U. S. A.* **2003**, *100*, 13940-13945.
- (7) Heintzen, C. Plant and Fungal Photopigments. *Wiley. Interdiscip. Rev. Membr. Transp. Sign.* **2012**, *1*, 411-432.
- (8) Menon, S. T.; Han, M.; Sakmar, T. P. Rhodopsin: Structural Basis of Molecular Physiology. *Physiol. Rev.* **2001**, *81*, 1659-1688.
- (9) van der Horst, M. A.; Hellingwerf, K. J. Photoreceptor Proteins, "star Actors of Modern Times": A Review of the Functional Dynamics in the Structure of Representative Members of Six Different Photoreceptor Families. *Acc. Chem. Res.* **2004**, *37*, 13-20.
- (10) Schoenlein, R. W.; Peteanu, L. A.; Mathies, R. A.; Shank, C. V. The First Step in Vision: Femtosecond Isomerization of Rhodopsin. *Science.* **1991**, *254*, 412-415.
- (11) Stehle, J.; Silvers, R.; Werner, K.; Chatterjee, D.; Gande, S.; Scholz, F.; Dutta, A.; Wachtveitl, J.; Klein-Seetharaman, J.; Schwalbe, H. Characterization of the Simultaneous Decay Kinetics of Metarhodopsin States II and III in Rhodopsin by Solution-state NMR Spectroscopy. *Angew. Chem. Int. Ed. Engl.* **2014**, *53*, 2078-2084.
- (12) Mathies, R. A. Photochemistry: A Coherent Picture of Vision. *Nat. Chem.* **2015**, *7*, 945-947.
- (13) Wang, Q.; Schoenlein, R. W.; Peteanu, L. A.; Mathies, R. A.; Shank, C. V. Vibrationally Coherent Photochemistry in the Femtosecond Primary Event of Vision. *Science.* **1994**, *266*, 422-424.
- (14) Johnson, P. J. M.; Halpin, A.; Morizumi, T.; Prokhorenko, V. I.; Ernst, O. P.; Miller, R. J. D. Local Vibrational Coherences Drive the Primary Photochemistry of Vision. *Nat. Chem.* **2015**, *7*, 980-986.

- (15) Bernardi, F.; Olivucci, M.; Robb, M. A. Potential Energy Surface Crossings in Organic Photochemistry. *Chem. Soc. Rev.* **1996**, *25*, 321.
- (16) González-Luque, R.; Garavelli, M.; Bernardi, F.; Merchán, M.; Robb, M. A.; Olivucci, M. Computational Evidence in Favor of a Two-state, Two-mode Model of the Retinal Chromophore Photoisomerization. *Proc. Natl. Acad. Sci. U. S. A.* **2000**, *97*, 9379-9384.
- (17) Schapiro, I.; Melaccio, F.; Laricheva, E. N.; Olivucci, M. Using the Computer to Understand the Chemistry of Conical Intersections. *Photochem. Photobiol. Sci.* **2011**, *10*, 867-886.
- (18) Rivalta, I.; Nenov, A.; Garavelli, M. Modelling Retinal Chromophores Photoisomerization: From Minimal Models in Vacuo to Ultimate Bidimensional Spectroscopy in Rhodopsins. *Phys. Chem. Chem. Phys.* **2014**, *16*, 16865-16879.
- (19) Bonačić-Koutecký, V.; Schöffel, K.; Michl, J. Critically Heterosymmetric Biradicaloid Geometries of Protonated Schiff Bases. *Theor. Chem. Acc.* **1987**, *72*, 459-474.
- (20) Garavelli, M.; Celani, P.; Bernardi, F.; Robb, M. A.; Olivucci, M. The C5H6NH²⁺ Protonated Schiff Base: An Ab Initio Minimal Model for Retinal Photoisomerization. *J. Am. Chem. Soc.* **1997**, *119*, 6891-6901.
- (21) Frutos, L. M.; Andruniów, T.; Santoro, F.; Ferré, N.; Olivucci, M. Tracking the Excited-state Time Evolution of the Visual Pigment with Multiconfigurational Quantum Chemistry. *Proc. Natl. Acad. Sci. U. S. A.* **2007**, *104*, 7764-7769.
- (22) Warshel, A. Bicycle-pedal Model for the First Step in the Vision Process. *Nature.* **1976**, *260*, 679-683.
- (23) Luk, H. L.; Melaccio, F.; Rinaldi, S.; Gozem, S.; Olivucci, M. Molecular Bases for the Selection of the Chromophore of Animal Rhodopsins. *Proc. Natl. Acad. Sci. U. S. A.* **2015**, *112*, 15297-15302.
- (24) Rinaldi, S.; Melaccio, F.; Gozem, S.; Fanelli, F.; Olivucci, M. Comparison of the Isomerization Mechanisms of Human Melanopsin and Invertebrate and Vertebrate Rhodopsins. *Proc. Natl. Acad. Sci. U. S. A.* **2014**, *111*, 1714-1719.
- (25) Ferré, N.; Olivucci, M. Probing the Rhodopsin Cavity with Reduced Retinal Models at the CASPT2//CASSCF/AMBER Level of Theory. *J. Am. Chem. Soc.* **2003**, *125*, 6868-6869.
- (26) Schapiro, I.; Weingart, O.; Buss, V. Bicycle-pedal Isomerization in a Rhodopsin Chromophore Model. *J. Am. Chem. Soc.* **2009**, *131*, 16-17.
- (27) Schapiro, I.; Ryazantsev, M. N.; Frutos, L. M.; Ferré, N.; Lindh, R.; Olivucci, M. The Ultrafast Photoisomerizations of Rhodopsin and Bathorhodopsin Are Modulated by Bond Alternation and HOOP Driven Electronic Effects. *J. Am. Chem. Soc.* **2011**, *133*, 3354-3364.
- (28) Weingart, O. The Twisted C11=C12 Bond of the Rhodopsin Chromophore--a Photochemical Hot Spot. *J. Am. Chem. Soc.* **2007**, *129*, 10618-10619.
- (29) Hayashi, S.; Taikhorshid, E.; Schulten, K. Photochemical Reaction Dynamics of the Primary Event of Vision Studied by Means of a Hybrid Molecular Simulation. *Biophys. J.* **2009**, *96*, 403-416.
- (30) Röhrig, U. F.; Guidoni, L.; Laió, A.; Frank, I.; Rothlisberger, U. A Molecular Spring for Vision. *J. Am. Chem. Soc.* **2004**, *126*, 15328-15329.
- (31) Sekharan, S.; Sugihara, M.; Weingart, O.; Okada, T.; Buss, V. Protein Assistance in the Photoisomerization of Rhodopsin and 9-cis-rhodopsin--insights From Experiment and Theory. *J. Am. Chem. Soc.* **2007**, *129*, 1052-1054.
- (32) Polli, D.; Altoè, P.; Weingart, O.; Spillane, K. M.; Manzoni, C.; Brida, D.; Tomasello, G.; Orlandi, G.; Kukura, P.; Mathies, R. A.; et al. Conical Intersection Dynamics of the Primary Photoisomerization Event in Vision. *Nature.* **2010**, *467*, 440-443.
- (33) Weingart, O.; Schapiro, I.; Buss, V. Photochemistry of Visual Pigment Chromophore Models by Ab Initio Molecular Dynamics. *J. Phys. Chem. B.* **2007**, *111*, 3782-3788.
- (34) Virshup, A. M.; Punwong, C.; Pogorelov, T. V.; Lindquist, B. A.; Ko, C.; Martínez, T. J. Photodynamics in Complex Environments: Ab Initio Multiple Spawning Quantum Mechanical/molecular Mechanical Dynamics. *J. Phys. Chem. B.* **2009**, *113*, 3280-3291.
- (35) Gozem, S.; Melaccio, F.; Lindh, R.; Krylov, A.; Granovsky, A.; Angeli, C.; Olivucci, M. Mapping the Excited State Potential Energy Surface of a Retinal Chromophore Model with Multireference and EOM-CC Methods. *J. Chem. Theory. Comput.* **2013**, *9*, 4495-4506.
- (36) Weiss, R. M.; Warshel, A. A New View of the Dynamics of Singlet Cis-trans Photoisomerization. *J. Am. Chem. Soc.* **1979**, *101*, 6131-6133.
- (37) Kim, J. E.; Tauber, M. J.; Mathies, R. A. Analysis of the Mode-specific Excited-state Energy Distribution and Wavelength-dependent Photoreaction Quantum Yield in Rhodopsin. *Biophys. J.* **2003**, *84*, 2492-2501.
- (38) Zener, C. Non-adiabatic Crossing of Energy Levels. *Proc. R. Soc. London., Ser. A.* **1932**, *137*, 696-702.
- (39) Weingart, O.; Altoè, P.; Stenta, M.; Bottoni, A.; Orlandi, G.; Garavelli, M. Product Formation in Rhodopsin by Fast Hydrogen Motions. *Phys. Chem. Chem. Phys.* **2011**, *13*, 3645-3648.
- (40) Polli, D.; Weingart, O.; Brida, D.; Poli, E.; Maiuri, M.; Spillane, K. M.; Bottoni, A.; Kukura, P.; Mathies, R. A.; Cerullo, G.; et al. Wavepacket Splitting and Two-pathway Deactivation in the Photoexcited Visual Pigment Isorhodopsin. *Angew. Chem. Int. Ed. Engl.* **2014**, *53*, 2504-2507.
- (41) Palczewski, K.; Kumasaka, T.; Hori, T.; Behnke, C. A.; Moto-shima, H.; Fox, B. A.; Trong, I. L.; Teller, D. C.; Okada, T.; Stenkamp, R. E.; et al. Crystal Structure of Rhodopsin: A G Protein-Coupled Receptor. *Science.* **2000**, *289*, 739-745.
- (42) Okada, T.; Sugihara, M.; Bondar, A. -N.; Elstner, M.; Entel, P.; Buss, V. The Retinal Conformation and Its Environment in Rhodopsin in Light of a New 2.2 Å Crystal Structure. *J. Mol. Biol.* **2004**, *342*, 571-583.
- (43) De Vico, L.; Page, C. S.; Garavelli, M.; Bernardi, F.; Basosi, R.; Olivucci, M. Reaction Path Analysis of the "Tunable" Photoisomerization Selectivity of Free and Locked Retinal Chromophores. *J. Am. Chem. Soc.* **2002**, *124*, 4124-4134.
- (44) Celani, P.; Garavelli, M.; Ottani, S.; Bernardi, F.; Robb, M. A.; Olivucci, M. Molecular "Trigger" for the Radiationless Deactivation of Photoexcited Conjugated Hydrocarbons. *J. Am. Chem. Soc.* **1995**, *117*, 11584-11585.
- (45) Ioffe, I. N.; Granovsky, A. A. Photoisomerization of Stilbene: The Detailed XMCQDPT2 Treatment. *J. Chem. Theory. Comput.* **2013**, *9*, 4973-4990.
- (46) Andersen, L. H.; Nielsen, I. B.; Kristensen, M. B.; El Ghazaly, M. O.; Haacke, S.; Nielsen, M. B.; Petersen, M. A. Absorption of Schiff-base Retinal Chromophores in Vacuo. *J. Am. Chem. Soc.* **2005**, *127*, 12347-12350.
- (47) Mizukami, T.; Kandori, H.; Shichida, Y.; Chen, A. -H.; Derguini, F.; Caldwell, C. G.; Biffe, C. F.; Nakanishi, K.; Yoshizawa, T. Photoisomerization Mechanism of the Rhodopsin Chromophore: Picosecond Photolysis of Pigment Containing 11-cis-locked Eight-membered Ring Retinal. *Proc. Natl. Acad. Sci. U. S. A.* **1993**, *90*, 4072-4076.
- (48) Becker, R. S. The Visual Process: Photophysics and Photoisomerization of Model Visual Pigments and the Primary Reaction. *Photochem. Photobiol.* **1988**, *48*, 369-399.
- (49) Munoz-Losa, A.; Ignacio, I.; Aguilar, M. A.; Martín, M. E. Simultaneous Solvent and Counterion Effects on the Absorption Properties of a Model of the Rhodopsin Chromophore. *J. Chem. Theory. Comput.* **2013**, *9*, 1548-1556.
- (50) Fujimoto, K.; Hayashi, S.; Hasegawa, J. Y.; Nakatsuji, H. Theoretical Studies on the Color-Tuning Mechanism in Retinal Proteins. *J. Chem. Theory. Comput.* **2007**, *3*, 605-618.
- (51) Altun, A.; Yokoyama, S.; Morokuma, K. Spectral Tuning in Visual Pigments: An ONIOM(QM:MM) Study on Bovine Rhodopsin and Its Mutants. *J. Phys. Chem. B.* **2008**, *112*, 6814-6827.
- (52) Sekharan, S.; Altun, A.; Morokuma, K. in *Annual Reports in Computational Chemistry* Wheeler, R., Ed.; Elsevier: Amsterdam, 2011; *7*, pp 215-233.
- (53) Melaccio, F.; Ferré, N.; Olivucci, M. Quantum Chemical Modeling of Rhodopsin Mutants Displaying Switchable Colors. *Phys. Chem. Chem. Phys.* **2012**, *14*, 12485-12495.

- (54) Wang, W.; Nossoni, Z.; Berbasova, T.; Watson, C. T.; Yapici, I.; Lee, K. S. S.; Vasileiou, C.; Geiger, J. H.; Borhan, B. Tuning the Electronic Absorption of Protein-Embedded All-trans-Retinal. *Science*. **2012**, *338*, 1340-1343.
- (55) Cheng, C.; Kamiya, M.; Uchida, Y.; Hayashi, S. Molecular Mechanism of Wide Photoabsorption Spectral Shifts of Color Variants of Human Cellular Retinol Binding Protein II. *J. Am. Chem. Soc.* **2015**, *137*, 13362-13370.
- (56) Gozem, S.; Schapiro, I.; Ferré, N.; Olivucci, M. The Molecular Mechanism of Dark Noise in Rod Photoreceptors. *Science*. **2012**, *337*, 1225-1228.
- (57) Coto, P. B.; Sinicropi, A.; De Vico, L.; Ferre, N.; Olivucci, M. Characterization of the Conical Intersection of the Visual Pigment Rhodopsin at the CASPT2//CASSCF/AMBER Level of Theory. *Mol. Phys.* **2006**, *104*, 983-991.
- (58) Atchity, G. J.; Xantheas, S. S.; Ruedenberg, K. Potential Energy Surfaces Near Intersections. *J. Phys. Chem.* **1991**, *95*, 1862-1876.
- (59) Levine, B. G.; Martínez, T. J. Isomerization Through Conical Intersections. *Annu. Rev. Phys. Chem.* **2007**, *58*, 613-634.
- (60) Yarkony, D. R. Nonadiabatic Quantum Chemistry--past, Present, and Future. *Chem. Rev.* **2012**, *112*, 481-498.
- (61) Berry, M. V. Quantal Phase Factors Accompanying Adiabatic Changes. *Proc. R. Soc. London, Ser. A*. **1984**, *392*, 45-57.
- (62) Gozem, S.; Melaccio, F.; Lindh, R.; Krylov, A. I.; Granovsky, A. A.; Angeli, C.; Olivucci, M. Mapping the Excited State Potential Energy Surface of a Retinal Chromophore Model with Multireference and Equation-of-Motion Coupled-Cluster Methods. *J. Chem. Theory. Comput.* **2013**, *9*, 4495-4506.
- (63) Wigner, E. On the Quantum Correction for Thermodynamic Equilibrium. *Phys. Rev.* **1932**, *40*, 749.
- (64) Bonacić-Koutecký, V.; Mitrić, R. Theoretical Exploration of Ultrafast Dynamics in Atomic Clusters: Analysis and Control. *Chem. Rev.* **2005**, *105*, 11-66.
- (65) Zgrablić, G.; Novello, A. M.; Parmigiani, F. Population Branching in the Conical Intersection of the Retinal Chromophore Revealed by Multipulse Ultrafast Optical Spectroscopy. *J. Am. Chem. Soc.* **2012**, *134*, 955-961.
- (66) Landau, L. D. Zur Theorie Der Energieübertragung. II. *Physikalische Zeitschrift der Sowjetunion*. **1932**, *2*, 46-51.
- (67) Stueckelberg, E. K. G. Theorie Der Unelastischen Stöße Zwischen Atomen. *Helv. Phys. Acta*. **1932**, *5*, 369-322.
- (68) Majorana, E. Atomi Orientati in Campo Magnetico Variabile. II. *Nuovo Cimento. (1924-1942)*. **1932**, *9*, 43-50.
- (69) Zgrablić, G.; Novello, A. M.; Parmigiani, F. Population Branching in the Conical Intersection of the Retinal Chromophore Revealed by Multipulse Ultrafast Optical Spectroscopy. *J. Am. Chem. Soc.* **2012**, *134*, 955-961.
- (70) Cheminal, A.; Léonard, J.; Kim, S. -Y.; Jung, K. -H.; Kandori, H.; Haacke, S. 100 Fs Photo-isomerization with Vibrational Coherences but Low Quantum Yield in Anabaena Sensory Rhodopsin. *Phys. Chem. Chem. Phys.* **2015**, *17*, 25429-25439.
- (71) Brumer, P.; Shapiro, M. Quantum Interference in the Control of Molecular Processes. *Philos. Trans. R. Soc. London, Ser. A*. **1997**, *355*, 2409-2412.
- (72) Klaffki, N.; Weingart, O.; Garavelli, M.; Spohr, E. Sampling Excited State Dynamics: Influence of HOOP Mode Excitations in a Retinal Model. *Phys. Chem. Chem. Phys.* **2012**, *14*, 14299-14305.
- (73) Manathunga, M.; Yang, X.; Luk, H. L.; Gozem, S.; Frutos, L. M.; Valentini, A.; Ferré, N.; Olivucci, M. Probing the Photodynamics of Rhodopsins with Reduced Retinal Chromophores. *J. Chem. Theory. Comput.* **2016**, *12*, 839-850.
- (74) Davis, D. A.; Hamilton, A.; Yang, J.; Cremer, L. D.; Van Gough, D.; Potisek, S. L.; Ong, M. T.; Braun, P. V.; Martínez, T. J.; White, S. R.; et al. Force-induced Activation of Covalent Bonds in Mechanoresponsive Polymeric Materials. *Nature*. **2009**, *459*, 68-72.
- (75) Valentini, A.; Rivero, D.; Zapata, F.; García-Iriepa, C.; Marazzi, M.; Palmeiro, R.; Fdez Galván, I.; Sampedro, D.; Olivucci, M.; Frutos, L. M. Optomechanical Control of Quantum Yield in Trans-Cis Ultrafast Photoisomerization of a Retinal Chromophore Model. *Angew. Chem. Int. Ed. Engl.* **2017**, *56*, 3842-3846.
- (76) Spudich, J. L. The Multitalented Microbial Sensory Rhodopsins. *Trends Microbiol.* **2006**, *14*, 480-487.
- (77) Cajochen, C.; Chellappa, S. L. Commentary: Melanopsin Regulates Both Sleep-Promoting and Arousal-Promoting Responses to Light. *Frontiers in neural circuits*. **2016**, *10*, 94.
- (78) Altoè, P.; Cembran, A.; Olivucci, M.; Garavelli, M. Aborted Double Bicycle-pedal Isomerization with Hydrogen Bond Breaking Is the Primary Event of Bacteriorhodopsin Proton Pumping. *Proc. Natl. Acad. Sci. U. S. A.* **2010**, *107*, 20172-20177.
- (79) Strambi, A.; Durbeij, B.; Ferré, N.; Olivucci, M. Anabaena Sensory Rhodopsin Is a Light-driven Unidirectional Rotor. *Proc. Natl. Acad. Sci. U. S. A.* **2010**, *107*, 21322-21326.
- (80) Kawanabe, A.; Furutani, Y.; Jung, K. H.; Kandori, H. Photochromism of Anabaena Sensory Rhodopsin. *J. Am. Chem. Soc.* **2007**, *129*, 8644-8649.
- (81) Vogeley, L.; Sineshchekov, O. A.; Trivedi, V. D.; Sasaki, J.; Spudich, J. L.; Luecke, H. Anabaena Sensory Rhodopsin: A Photochromic Color Sensor at 2.0 Å. *Science*. **2004**, *306*, 1390-1393.
- (82) Suzuki, T.; Callender, R. H. Primary Photochemistry and Photoisomerization of Retinal at 77 Degrees K in Cattle and Squid Rhodopsins. *Biophys. J.* **1981**, *34*, 261-270.
- (83) Nadochenko, V. A.; Smitienko, O. A.; Feldman, T. B.; Mozgovaya, M. N.; Shelaev, I. V.; Gostev, F. E.; Sarkisov, O. M.; Ostrovsky, M. A. Conical Intersection Participation in Femtosecond Dynamics of Visual Pigment Rhodopsin Chromophore Cis-trans Photoisomerization. *Dokl. Biochem. Biophys.* **2012**, *446*, 242-246.
- (84) Smitienko, O. A.; Mozgovaya, M. N.; Shelaev, I. V.; Gostev, F. E.; Feldman, T. B.; Nadochenko, V. A.; Sarkisov, O. M.; Ostrovsky, M. A. Femtosecond Formation Dynamics of Primary Photoproducts of Visual Pigment Rhodopsin. *Biochemistry. (Moscow)*. **2010**, *75*, 25-35.
- (85) Birge, R. R.; Hubbard, L. M. Molecular Dynamics of Trans-cis Isomerization in Bathorhodopsin. *Biophys. J.* **1981**, *34*, 517-534.
- (86) Altoè, P.; Cembran, A.; Olivucci, M.; Garavelli, M. Aborted Double Bicycle-pedal Isomerization with Hydrogen Bond Breaking Is the Primary Event of Bacteriorhodopsin Proton Pumping. *Proc. Natl. Acad. Sci. U. S. A.* **2010**, *107*, 20172-20177.
- (87) Koumura, N.; Zijlstra, R. W. J.; van Delden, R. W.; Harada, N.; Feringa, B. L. Light-driven Monodirectional Molecular Rotor. *Nature*. **1999**, *401*, 152-155.
- (88) Dokukina, I.; Weingart, O. Spectral Properties and Isomerisation Path of Retinal in C1C2 Channelrhodopsin. *Phys. Chem. Chem. Phys.* **2015**, *17*, 25142-25150.
- (89) Hontani, Y.; Marazzi, M.; Stehfest, K.; Mathes, T.; van Stokkum, I. H. M.; Elstner, M.; Hegemann, P.; Kennis, J. T. M. Reaction Dynamics of the Chimeric Channelrhodopsin C1C2. *Sci. Rep.* **2017**, *7*, 7217.
- (90) Gozem, S.; Huntress, M.; Schapiro, I.; Lindh, R.; Granovsky, A. A.; Angeli, C.; Olivucci, M. Dynamic Electron Correlation Effects on the Ground State Potential Energy Surface of a Retinal Chromophore Model. *J. Chem. Theory. Comput.* **2012**, *8*, 4069-4080.
- (91) Khrenova, M. G.; Bochenkova, A. V.; Nemukhin, A. V. Modeling Reaction Routes From Rhodopsin to Bathorhodopsin. *Proteins. Struct. Funct. Bioinf.* **2010**, *78*, 614-622.
- (92) Barlow, H. B. Purkinje Shift and Retinal Noise. *Nature*. **1957**, *179*, 255-256.
- (93) Ala-Laurila, P.; Pahlberg, J.; Koskelainen, A.; Donner, K. On the Relation Between the Photoactivation Energy and the Absorbance Spectrum of Visual Pigments. *Vision. Res.* **2004**, *44*, 2153-2158.
- (94) Ala-Laurila, P.; Donner, K.; Koskelainen, A. Thermal Activation and Photoactivation of Visual Pigments. *Biophys. J.* **2004**, *86*, 3653-3662.
- (95) Luo, D. G.; Yue, W. W.; Ala-Laurila, P.; Yau, K. W. Activation of Visual Pigments by Light and Heat. *Science*. **2011**, *332*, 1307-1312.

- (96) Luk, H. L.; Bhattacharyya, N.; Montisci, F.; Morrow, J. M.; Melaccio, F.; Wada, A.; Sheves, M.; Fanelli, F.; Chang, B. S. W.; Olivucci, M. Modulation of Thermal Noise and Spectral Sensitivity in Lake Baikal Cottoid Fish Rhodopsins. *Sci. Rep.* **2016**, *6*, 38425.
- (97) Warshel, A.; Karplus, M. Calculation of Ground and Excited State Potential Surfaces of Conjugated Molecules. I. Formulation and Parametrization. *J. Am. Chem. Soc.* **1972**, *94*, 5612-5625.
- (98) Warshel, A. Quantum Mechanical Consistent Force Field (QCFF/PI) Method: Calculations of Energies, Conformations and Vibronic Interactions of Ground and Excited States of Conjugated Molecules. *Isr. J. Chem.* **1973**, *11*, 709-717.
- (99) Birge, R. R.; Hubbard, L. M. Molecular Dynamics of Cis-trans Isomerization in Rhodopsin. *J. Am. Chem. Soc.* **1980**, *102*, 2195-2205.
- (100) Warshel, A.; Barboy, N. Energy Storage and Reaction Pathways in the First Step of the Vision Process. *J. Am. Chem. Soc.* **1982**, *104*, 1469-1476.
- (101) Liu, R. S.; Asato, A. E. The Primary Process of Vision and the Structure of Bathorhodopsin: A Mechanism for Photoisomerization of Polyenes. *Proc. Natl. Acad. Sci. U. S. A.* **1985**, *82*, 259-263.
- (102) Liu, R. S.; Hammond, G. S. The Case of Medium-dependent Dual Mechanisms for Photoisomerization: One-bond-flip and Hula-twist. *Proc. Natl. Acad. Sci. U. S. A.* **2000**, *97*, 11153-11158.
- (103) Liu, R. S. Photoisomerization by Hula-twist: A Fundamental Supramolecular Photochemical Reaction. *Acc. Chem. Res.* **2001**, *34*, 555-562.
- (104) Szymczak, J. J.; Barbatti, M.; Lischka, H. Is the Photoinduced Isomerization in Retinal Protonated Schiff Bases a Single- or Double-torsional Process?. *J. Phys. Chem. A.* **2009**, *113*, 11907-11918.
- (105) Granucci, G.; Toniolo, A. Molecular Gradients for Semiempirical CI Wavefunctions with Floating Occupation Molecular Orbitals. *Chem. Phys. Lett.* **2000**, *325*, 79-85.
- (106) Chung, W. C.; Nanbu, S.; Ishida, T. QM/MM Trajectory Surface Hopping Approach to Photoisomerization of Rhodopsin and Isorhodopsin: The Origin of Faster and More Efficient Isomerization for Rhodopsin. *J. Phys. Chem. B.* **2012**, *116*, 8009-8023.
- (107) Snyder, J. W.; Parrish, R. M.; Martínez, T. J. A-CASSCF: An Efficient, Empirical Correction for SA-CASSCF to Closely Approximate MS-CASPT2 Potential Energy Surfaces. *J. Phys. Chem. Lett.* **2017**, *8*, 2432-2437.
- (108) Birge, R. R.; Findsen, L. A.; Pierce, B. M. Molecular Dynamics of the Primary Photochemical Event in Bacteriorhodopsin. Theoretical Evidence for An Excited Singlet State Assignment for the J Intermediate. *J. Am. Chem. Soc.* **1987**, *109*, 5041-5043.
- (109) Warshel, A.; Chu, Z. T. Nature of the Surface Crossing Process in Bacteriorhodopsin: Computer Simulations of the Quantum Dynamics of the Primary Photochemical Event. *J. Phys. Chem. B.* **2001**, *105*, 9857-9871.
- (110) Warshel, A. Molecular Dynamics Simulations of Biological Reactions. *Acc. Chem. Res.* **2002**, *35*, 385-395.
- (111) Hayashi, S.; Tajkhorshid, E.; Schulten, K. Ab Initio QM/MM Molecular Dynamics Study of Retinal Photoisomerization in Bacteriorhodopsin. *Biophys. J.* **2003**, *84*, 269A-269A.
- (112) Li, X.; Chung, L. W.; Morokuma, K. Photodynamics of All-trans Retinal Protonated Schiff Base in Bacteriorhodopsin and Methanol Solution. *J. Chem. Theory. Comput.* **2011**, *7*, 2694-2698.
- (113) Punwong, C.; Owens, J.; Martínez, T. J. Direct QM/MM Excited-state Dynamics of Retinal Protonated Schiff Base in Isolation and Methanol Solution. *J. Phys. Chem. B.* **2014**, *119*, 704-714.
- (114) Ben-Nun, M.; Molnar, F.; Schulten, K.; Martinez, T. J. The Role of Intersection Topography in Bond Selectivity of Cis-trans Photoisomerization. *Proc. Natl. Acad. Sci. U. S. A.* **2002**, *99*, 1769-1773.
- (115) Punwong, C.; Martínez, T. J.; Hannongbua, S. Direct QM/MM Simulation of Photoexcitation Dynamics in Bacteriorhodopsin and Halorhodopsin. *Chem. Phys. Lett.* **2014**, *610*, 213-218.
- (116) Wand, A.; Rozin, R.; Eliash, T.; Jung, K. H.; Sheves, M.; Ruhman, S. Asymmetric Toggling of a Natural Photoswitch: Ultrafast Spectroscopy of Anabaena Sensory Rhodopsin. *J. Am. Chem. Soc.* **2011**, *133*, 20922-20932.
- (117) Schapiro, I.; Ruhman, S. Ultrafast Photochemistry of Anabaena Sensory Rhodopsin: Experiment and Theory. *Biochim. Biophys. Acta.* **2014**, *1837*, 589-597.
- (118) Granovsky, A. A. Extended Multi-configuration Quasi-degenerate Perturbation Theory: The New Approach to Multi-state Multi-reference Perturbation Theory. *J. Phys. Chem.* **2011**, *134*, 214113.
- (119) Yabushita, A.; Kobayashi, T.; Tsuda, M. Time-resolved Spectroscopy of Ultrafast Photoisomerization of Octopus Rhodopsin Under Photoexcitation. *J. Phys. Chem. B.* **2012**, *116*, 1920-1926.
- (120) Gai, F.; Hasson, K. C.; McDonald, J. C.; Anfinsenrud, P. A. Chemical Dynamics in Proteins: The Photoisomerization of Retinal in Bacteriorhodopsin. *Science.* **1998**, *279*, 1886.
- (121) Garavelli, M.; Celani, P.; Bernardi, F.; Robb, M. A.; Olivucci, M. Force Fields for ultrafast Photochemistry: The S2 (1Bu) S1 (2Ag) S0 (1Ag) Reaction Path for All-trans-Hexa-1, 3, 5-triene. *J. Am. Chem. Soc.* **1997**, *119*, 11487-11494.
- (122) Cembran, A.; González-Luque, R.; Serrano-Andrés, L.; Merchán, M.; Garavelli, M. About the Intrinsic Photochemical Properties of the 11-cis Retinal Chromophore: Computational Clues for a Trap State and a Lever Effect in Rhodopsin Catalysis. *Theor. Chem. Acc.* **2007**, *118*, 173-183.
- (123) Briand, J.; Léonard, J.; Haacke, S. Ultrafast Photo-induced Reaction Dynamics in Bacteriorhodopsin and Its Trp Mutants. *J. Opt.* **2010**, *12*, 084004.
- (124) Bassolino, G.; Sovdat, T.; Liebel, M.; Schnedermann, C.; Odell, B.; Claridge, T. D.; Kukura, P.; Fletcher, S. P. Synthetic Control of Retinal Photochemistry and Photophysics in Solution. *J. Am. Chem. Soc.* **2014**, *136*, 2650-2658.
- (125) Cembran, A.; Bernardi, F.; Olivucci, M.; Garavelli, M. The Retinal Chromophore/chloride Ion Pair: Structure of the Photoisomerization Path and Interplay of Charge Transfer and Covalent States. *Proc. Natl. Acad. Sci. U. S. A.* **2005**, *102*, 6255-6260.
- (126) Zgrablić, G.; Voitchovsky, K.; Kindermann, M.; Haacke, S.; Chergui, M. Ultrafast Excited State Dynamics of the Protonated Schiff Base of All-trans Retinal in Solvents. *Biophys. J.* **2005**, *88*, 2779-2788.
- (127) Muñoz-Losa, A.; Martín, M. E.; Fernández Galván, I.; Sánchez, M. L.; Aguilar, M. Solvent Effects on the Radiative and Non-radiative Decay of a Model of the Rhodopsin Chromophore. *J. Chem. Theory. Comput.* **2011**, *7*, 4050-4059.
- (128) Sprenger, W. W.; Hoff, W. D.; Armitage, J. P.; Hellingwerf, K. J. The Eubacterium Ectothiorhodospira Halophila Is Negatively Phototactic, with a Wavelength Dependence That Fits the Absorption Spectrum of the Photoactive Yellow Protein. *J. Bacteriol.* **1993**, *175*, 3096-3104.
- (129) Kort, R.; Hoff, W. D.; Van West, M.; Kroon, A. R.; Hoffer, S. M.; Vlieg, K. H.; Crielaand, W.; Van Beeumen, J. J.; Hellingwerf, K. J. The Xanthopsins: A New Family of Eubacterial Blue-light Photoreceptors. *EMBO. J.* **1996**, *15*, 3209-3218.
- (130) Meyer, T. E. Isolation and Characterization of Soluble Cytochromes, Ferredoxins and Other Chromophoric Proteins From the Halophilic Phototrophic Bacterium Ectothiorhodospira Halophila. *Biochim. Biophys. Acta.* **1985**, *806*, 175-183.
- (131) Zhao, J. M.; Lee, H.; Nome, R. A.; Majid, S.; Scherer, N. F.; Hoff, W. D. Single-molecule Detection of Structural Changes During Per-Arnt-Sim (PAS) Domain Activation. *Proc. Natl. Acad. Sci. U. S. A.* **2006**, *103*, 11561-11566.
- (132) Cusanovich, M. A.; Meyer, T. E. Photoactive Yellow Protein: A Prototypic PAS Domain Sensory Protein and Development of a Common Signaling Mechanism. *Biochemistry.* **2003**, *42*, 4759-4770.
- (133) Lee, I. R.; Lee, W.; Zewail, A. H. Primary Steps of the Photoactive Yellow Protein: Isolated Chromophore Dynamics and Protein Directed Function. *Proc. Natl. Acad. Sci. U. S. A.* **2006**, *103*, 258-262.

- (134) El-Gezawy, H.; Rettig, W.; Danel, A.; Jonusauskas, G. Probing the Photochemical Mechanism in Photoactive Yellow Protein. *J. Phys. Chem. B.* **2005**, *109*, 18699-18705.
- (135) Espagne, A.; Paik, D. H.; Changuenet-Barret, P.; Martin, M. M.; Zewail, A. H. Ultrafast Photoisomerization of Photoactive Yellow Protein Chromophore Analogues in Solution: Influence of the Protonation State. *ChemPhysChem.* **2006**, *7*, 1717-1726.
- (136) Groenhof, G.; Bouxin-Cademartory, M.; Hess, B.; De Visser, S. P.; Berendsen, H. J.; Olivucci, M.; Mark, A. E.; Robb, M. A. Photoactivation of the Photoactive Yellow Protein: Why Photon Absorption Triggers a Trans-to-cis Isomerization of the Chromophore in the Protein. *J. Am. Chem. Soc.* **2004**, *126*, 4228-4233.
- (137) Gromov, E. V.; Burghardt, I.; Hynes, J. T.; Köppel, H.; Cederbaum, L. S. Electronic Structure of the Photoactive Yellow Protein Chromophore: Ab Initio Study of the Low-lying Excited Singlet States. *J. Photochem. Photobiol. A.* **2007**, *190*, 241-257.
- (138) Gromov, E. V.; Burghardt, I.; Köppel, H.; Cederbaum, L. S. Photoinduced Isomerization of the Photoactive Yellow Protein (PYP) Chromophore: Interplay of Two Torsions, a HOOP Mode and Hydrogen Bonding. *J. Phys. Chem. A.* **2011**, *115*, 9237-9248.
- (139) Boggio-Pasqua, M.; Robb, M. A.; Groenhof, G. Hydrogen Bonding Controls Excited-state Decay of the Photoactive Yellow Protein Chromophore. *J. Am. Chem. Soc.* **2009**, *131*, 13580-13581.
- (140) Wei, L.; Wang, H.; Chen, X.; Fang, W.; Wang, H. A Comprehensive Study of Isomerization and Protonation Reactions in the Photocycle of the Photoactive Yellow Protein. *Phys. Chem. Chem. Phys.* **2014**, *16*, 25263-25272.
- (141) Pande, K.; Hutchison, C. D.; Groenhof, G.; Aquila, A.; Robinson, J. S.; Tenboer, J.; Basu, S.; Boutet, S.; DePonte, D. P.; Liang, M.; et al. Femtosecond Structural Dynamics Drives the Trans/cis Isomerization in Photoactive Yellow Protein. *Science.* **2016**, *352*, 725-729.
- (142) Ko, C.; Virshup, A. M.; Martínez, T. J. Electrostatic Control of Photoisomerization in the Photoactive Yellow Protein Chromophore: Ab Initio Multiple Spawning Dynamics. *Chem. Phys. Lett.* **2008**, *460*, 272-277.
- (143) Jung, Y. O.; Lee, J. H.; Kim, J.; Schmidt, M.; Moffat, K.; Srajer, V.; Ihee, H. Volume-conserving Trans-cis Isomerization Pathways in Photoactive Yellow Protein Visualized by Picosecond X-ray Crystallography. *Nat. Chem.* **2013**, *5*, 212-220.
- (144) Kaila, V. R.; Schotte, F.; Cho, H. S.; Hummer, G.; Anfinrud, P. A. Contradictions in X-ray Structures of Intermediates in the Photocycle of Photoactive Yellow Protein. *Nat. Chem.* **2014**, *6*, 258-259.
- (145) Jung, Y. O.; Lee, J. H.; Kim, J.; Schmidt, M.; Moffat, K.; Srajer, V.; Ihee, H. Reply to 'Contradictions in X-ray Structures of Intermediates in the Photocycle of Photoactive Yellow Protein'. *Nat. Chem.* **2014**, *6*, 259-260.
- (146) Schotte, F.; Cho, H. S.; Kaila, V. R.; Kamikubo, H.; Dashdorj, N.; Henry, E. R.; Graber, T. J.; Henning, R.; Wulff, M.; Hummer, G.; et al. Watching a Signaling Protein Function in Real Time Via 100-ps Time-resolved Laue Crystallography. *Proc. Natl. Acad. Sci. U. S. A.* **2012**, *109*, 19256-19261.
- (147) Gromov, E. V. Unveiling the Mechanism of Photoinduced Isomerization of the Photoactive Yellow Protein (PYP) Chromophore. *J. Phys. Chem.* **2014**, *141*, 224308.
- (148) Ihee, H.; Rajagopal, S.; Srajer, V.; Pahl, R.; Anderson, S.; Schmidt, M.; Schotte, F.; Anfinrud, P. A.; Wulff, M.; Moffat, K. Visualizing Reaction Pathways in Photoactive Yellow Protein From Nanoseconds to Seconds. *Proc. Natl. Acad. Sci. U. S. A.* **2005**, *102*, 7145-7150.
- (149) Liu, R. S.; Asato, A. E. The Primary Process of Vision and the Structure of Bathorhodopsin: A Mechanism for Photoisomerization of Polyenes. *Proc. Natl. Acad. Sci. U. S. A.* **1985**, *82*, 259-263.
- (150) Meyer, T. E.; Tollin, G.; Causgrove, T. P.; Cheng, P.; Blankenship, R. E. Picosecond Decay Kinetics and Quantum Yield of Fluorescence of the Photoactive Yellow Protein From the Halophilic Purple Phototrophic Bacterium, *Ectothiorhodospira Halophila*. *Biophys. J.* **1991**, *59*, 988-991.
- (151) Hellingwerf, K. J.; Hendriks, J.; Gensch, T. Photoactive Yellow Protein, A New Type of Photoreceptor Protein: Will This Yellow Lab Bring Us Where We Want to Go?. *J. Phys. Chem. A.* **2003**, *107*, 1082-1094.
- (152) Naseem, S.; Laurent, A. D.; Carroll, E. C.; Vengris, M.; Kumauchi, M.; Hoff, W. D.; Krylov, A. I.; Larsen, D. S. Photoisomerization Upshifts the PK a of the Photoactive Yellow Protein Chromophore to Contribute to Photocycle Propagation. *J. Photochem. Photobiol. A.* **2013**, *270*, 43-52.
- (153) Changuenet-Barret, P.; Plaza, P.; Martin, M. M.; Chosrowjan, H.; Taniguchi, S.; Mataga, N.; Imamoto, Y.; Kataoka, M. Role of Arginine 52 on the Primary Photoinduced Events in the PYP Photocycle. *Chem. Phys. Lett.* **2007**, *434*, 320-325.
- (154) Groenhof, G.; Schäfer, L. V.; Boggio-Pasqua, M.; Grubmüller, H.; Robb, M. A. Arginine52 Controls the Photoisomerization Process in Photoactive Yellow Protein. *J. Am. Chem. Soc.* **2008**, *130*, 3250-3251.
- (155) Takeshita, K.; Imamoto, Y.; Kataoka, M.; Mihara, K.; Tokunaga, F.; Terazima, M. Structural Change of Site-directed Mutants of PYP: New Dynamics During PR State. *Biophys. J.* **2002**, *83*, 1567-1577.
- (156) Cleland, W. W.; Kreevoy, M. M. Low-barrier Hydrogen Bonds and Enzymic Catalysis. *Science.* **1994**, *264*, 1887-1890.
- (157) Yamaguchi, S.; Kamikubo, H.; Kurihara, K.; Kuroki, R.; Nii-mura, N.; Shimizu, N.; Yamazaki, Y.; Kataoka, M. Low-barrier Hydrogen Bond in Photoactive Yellow Protein. *Proc. Natl. Acad. Sci. U. S. A.* **2009**, *106*, 440-444.
- (158) Saito, K.; Ishikita, H. Energetics of Short Hydrogen Bonds in Photoactive Yellow Protein. *Proc. Natl. Acad. Sci. U. S. A.* **2012**, *109*, 167-172.
- (159) Nadal-Ferret, M.; Gelabert, R.; Moreno, M.; Lluch, J. M. Are There Really Low-barrier Hydrogen Bonds in Proteins? The Case of Photoactive Yellow Protein. *J. Am. Chem. Soc.* **2014**, *136*, 3542-3552.
- (160) Graen, T.; Inhester, L.; Clemens, M.; Grubmüller, H.; Groenhof, G. The Low Barrier Hydrogen Bond in the Photoactive Yellow Protein: A Vacuum Artifact Absent in the Crystal and Solution. *J. Am. Chem. Soc.* **2016**, *138*, 16620-16631.
- (161) Kuramochi, H.; Takeuchi, S.; Tahara, T. Ultrafast Structural Evolution of Photoactive Yellow Protein Chromophore Revealed by Ultraviolet Resonance Femtosecond Stimulated Raman Spectroscopy. *J. Phys. Chem. Lett.* **2012**, *3*, 2025-2029.
- (162) Nakamura, R.; Hamada, N.; Abe, K.; Yoshizawa, M. Ultrafast Hydrogen-bonding Dynamics in the Electronic Excited State of Photoactive Yellow Protein Revealed by Femtosecond Stimulated Raman Spectroscopy. *J. Phys. Chem. B.* **2012**, *116*, 14768-14775.
- (163) Mix, L. T.; Kirpich, J.; Kumauchi, M.; Ren, J.; Vengris, M.; Hoff, W. D.; Larsen, D. S. Bifurcation in the Ultrafast Dynamics of the Photoactive Yellow Proteins From *Leptospira Biflexa* and *Halorhodospira Halophila*. *Biochemistry.* **2016**, *55*, 6138-6149.
- (164) Karniol, B.; Wagner, J. R.; Walker, J. M.; Vierstra, R. D. Phylogenetic Analysis of the Phytochrome Superfamily Reveals Distinct Microbial Subfamilies of Photoreceptors. *Biochem. J.* **2005**, *392*, 103-116.
- (165) Rockwell, N. C.; Su, Y. -S.; Lagarias, J. C. Phytochrome Structure and Signaling Mechanisms. *Annu. Rev. Plant. Biol.* **2006**, *57*, 837-858.
- (166) Kneip, C.; Hildebrandt, P.; Schlamann, W.; Braslavsky, S. E.; Mark, F.; Schaffner, K. Protonation State and Structural Changes of the Tetrapyrrole Chromophore During the Pr --> Pfr Phototransformation of Phytochrome: A Resonance Raman Spectroscopic Study. *Biochemistry.* **1999**, *38*, 15185-15192.
- (167) Andel, F.; Murphy, J. T.; Haas, J. A.; McDowell, M. T.; van der Hoef, I.; Lugtenburg, J.; Lagarias, J. C.; Mathies, R. A. Probing the Photoreaction Mechanism of Phytochrome Through Analysis of Resonance Raman Vibrational Spectra of Recombinant Analogues. *Biochemistry.* **2000**, *39*, 2667-2676.
- (168) Burgie, E. S.; Bussell, A. N.; Walker, J. M.; Dubiel, K.; Vierstra, R. D. Crystal Structure of the Photosensing Module From a

- Red/far-red Light-absorbing Plant Phytochrome. *Proc. Natl. Acad. Sci. U. S. A.* **2014**, *111*, 10179-10184.
- (169) Schumann, C.; Gross, R.; Michael, N.; Lamparter, T.; Diller, R. Sub-picosecond Mid-infrared Spectroscopy of Phytochrome Agp1 From *Agrobacterium Tumefaciens*. *ChemPhysChem*. **2007**, *8*, 1657-1663.
- (170) Fodor, S. P.; Lagarias, J. C.; Mathies, R. A. Resonance Raman Analysis of the Pr and Pfr Forms of Phytochrome. *Biochemistry*. **1990**, *29*, 11141-11146.
- (171) Andel, F.; Lagarias, J. C.; Mathies, R. A. Resonance Raman Analysis of Chromophore Structure in the Lumi-R Photoproduct of Phytochrome. *Biochemistry*. **1996**, *35*, 15997-16008.
- (172) Murgida, D. H.; von Stetten, D.; Hildebrandt, P.; Schwinté, P.; Siebert, F.; Sharda, S.; Gärtner, W.; Mroginski, M. A. The Chromophore Structures of the Pr States in Plant and Bacterial Phytochromes. *Biophys. J.* **2007**, *93*, 2410-2417.
- (173) Mroginski, M. A.; Von Stetten, D.; Kaminski, S.; Escobar, F. V.; Michael, N.; Daminelli-Widany, G.; Hildebrandt, P. Elucidating Photoinduced Structural Changes in Phytochromes by the Combined Application of Resonance Raman Spectroscopy and Theoretical Methods. *J. Mol. Struct.* **2011**, *993*, 15-25.
- (174) Song, C.; Essen, L. O.; Gärtner, W.; Hughes, J.; Matysik, J. Solid-state NMR Spectroscopic Study of Chromophore-protein Interactions in the Pr Ground State of Plant Phytochrome A. *Mol. Plant*. **2012**, *5*, 698-715.
- (175) Song, C.; Rohmer, T.; Tiersch, M.; Zaanen, J.; Hughes, J.; Matysik, J. Solid-state NMR Spectroscopy to Probe Photoactivation in Canonical Phytochromes. *Photochem. Photobiol.* **2013**, *89*, 259-273.
- (176) Wagner, J. R.; Brunzelle, J. S.; Forest, K. T.; Vierstra, R. D. A Light-sensing Knot Revealed by the Structure of the Chromophore-binding Domain of Phytochrome. *Nature*. **2005**, *438*, 325-331.
- (177) Wagner, J. R.; Zhang, J.; Brunzelle, J. S.; Vierstra, R. D.; Forest, K. T. High Resolution Structure of *Deinococcus* Bacteriophytochrome Yields New Insights Into Phytochrome Architecture and Evolution. *J. Biol. Chem.* **2007**, *282*, 12298-12309.
- (178) Takala, H.; Björling, A.; Berntsson, O.; Lehtivuori, H.; Niebling, S.; Hoernke, M.; Kosheleva, I.; Henning, R.; Menzel, A.; Ihalainen, J. A.; et al. Signal Amplification and Transduction in Phytochrome Photosensors. *Nature*. **2014**, *509*, 245-248.
- (179) Yang, Y.; Linke, M.; von Haimberger, T.; Hahn, J.; Matute, R.; González, L.; Schmieder, P.; Heyne, K. Real-time Tracking of Phytochrome's Orientational Changes During Pr Photoisomerization. *J. Am. Chem. Soc.* **2012**, *134*, 1408-1411.
- (180) Yang, Y.; Linke, M.; von Haimberger, T.; Matute, R.; González, L.; Schmieder, P.; Heyne, K. Active and Silent Chromophore Isoforms for Phytochrome Pr Photoisomerization: An Alternative Evolutionary Strategy to Optimize Photoreaction Quantum Yields. *Struct. Dyn.* **2014**, *1*, 014701.
- (181) Levine, B. G.; Ko, C.; Quenneville, J.; Martínez, T. J. Conical Intersections and Double Excitations in Time-dependent Density Functional Theory. *Mol. Phys.* **2006**, *104*, 1039-1051.
- (182) Gozem, S.; Melaccio, F.; Valentini, A.; Filatov, M.; Huix-Rotllant, M.; Ferre, N.; Frutos, L. M.; Angeli, C.; Krylov, A. I.; Granovsky, A. A. Shape of Multireference, Equation-of-motion Coupled-cluster, and Density Functional Theory Potential Energy Surfaces at a Conical Intersection. *J. Chem. Theory. Comput.* **2014**, *10*, 3074-3084.
- (183) Tomasi, J.; Persico, M. Molecular Interactions in Solution: An Overview of Methods Based on Continuous Distributions of the Solvent. *Chem. Rev.* **1994**, *94*, 2027-2094.
- (184) Cramer, C. J.; Truhlar, D. G. Implicit Solvation Models: Equilibria, Structure, Spectra, and Dynamics. *Chem. Rev.* **1999**, *99*, 2161-2200.
- (185) Tomasi, J.; Mennucci, B.; Cammi, R. Quantum Mechanical Continuum Solvation Models. *Chem. Rev.* **2005**, *105*, 2999-3094.
- (186) Hasegawa, J. -Y.; Isshiki, M.; Fujimoto, K.; Nakatsuji, H. Structure of Phytochromobilin in the P R and P Fr Forms: SAC-CI Theoretical Study. *Chem. Phys. Lett.* **2005**, *410*, 90-93.
- (187) Nakatsuji, H. Cluster Expansion of the Wavefunction. Electron Correlations in Ground and Excited States by SAC (symmetry-adapted-cluster) and SAC CI Theories. *Chem. Phys. Lett.* **1979**, *67*, 329-333.
- (188) Nakatsuji, H. Cluster Expansion of the Wavefunction. Calculation of Electron Correlations in Ground and Excited States by SAC and SAC CI Theories. *Chem. Phys. Lett.* **1979**, *67*, 334-342.
- (189) Nakatsuji, H. Description of Two-and Many-electron Processes by the SAC-CI Method. *Chem. Phys. Lett.* **1991**, *177*, 331-337.
- (190) Velazquez Escobar, F.; von Stetten, D.; Günther-Lütken, M.; Keidel, A.; Michael, N.; Lamparter, T.; Essen, L. O.; Hughes, J.; Gärtner, W.; Yang, Y.; et al. Conformational Heterogeneity of the Pfr Chromophore in Plant and Cyanobacterial Phytochromes. *Front. Mol. Biosci.* **2015**, *2*, 37.
- (191) Garavelli, M.; Vreven, T.; Celani, P.; Bernardi, F.; Robb, M. A.; Olivucci, M. Photoisomerization Path for a Realistic Retinal Chromophore Model: The Nonatetraeniminium Cation. *J. Am. Chem. Soc.* **1998**, *120*, 1285-1288.
- (192) Garavelli, M.; Bernardi, F.; Robb, M. A.; Olivucci, M. The Short-chain Acroleiniminium and Pentadieniminium Cations: Towards a Model for Retinal Photoisomerization. A CASSCF/PT2 Study. *J. Mol. Struct. THEOCHEM.* **1999**, *463*, 59-64.
- (193) Andruniów, T.; Ferré, N.; Olivucci, M. Structure, Initial Excited-state Relaxation, and Energy Storage of Rhodopsin Resolved at the Multiconfigurational Perturbation Theory Level. *Proc. Natl. Acad. Sci. U. S. A.* **2004**, *101*, 17908-17913.
- (194) Durbeej, B.; Borg, O. A.; Eriksson, L. A. Phytochromobilin C15-Z, Syn C15-E, Anti Isomerization: Concerted or Stepwise?. *Phys. Chem. Chem. Phys.* **2004**, *6*, 5066-5073.
- (195) Durbeej, B.; Borg, O. A.; Eriksson, L. A. Computational Evidence in Favor of a Protonated Chromophore in the Photoactivation of Phytochrome. *Chem. Phys. Lett.* **2005**, *416*, 83-88.
- (196) Durbeej, B.; Eriksson, L. A. Protein-bound Chromophores Astaxanthin and Phytochromobilin: Excited State Quantum Chemical Studies. *Phys. Chem. Chem. Phys.* **2006**, *8*, 4053-4071.
- (197) Durbeej, B. On the Primary Event of Phytochrome: Quantum Chemical Comparison of Photoreactions at C4, C10 and C15. *Phys. Chem. Chem. Phys.* **2009**, *11*, 1354-1361.
- (198) Strambi, A.; Durbeej, B. Initial Excited-state Relaxation of the Bilin Chromophores of Phytochromes: A Computational Study. *Photochem. Photobiol. Sci.* **2011**, *10*, 569-579.
- (199) Zhuang, X.; Wang, J.; Lan, Z. Tracking of the Molecular Motion in the Primary Event of Photoinduced Reactions of a Phytochromobilin Model. *J. Phys. Chem. B.* **2013**, *117*, 15976-15986.
- (200) Falklöf, O.; Durbeej, B. Steric Effects Govern the Photoactivation of Phytochromes. *ChemPhysChem*. **2016**, *17*, 954-957.
- (201) Altoè, P.; Climent, T.; De Fusco, G. C.; Stenta, M.; Bottoni, A.; Serrano-Andrés, L.; Merchán, M.; Orlandi, G.; Garavelli, M. Deciphering Intrinsic Deactivation/isomerization Routes in a Phytochrome Chromophore Model. *J. Phys. Chem. B.* **2009**, *113*, 15067-15073.
- (202) Shcherbakova, D. M.; Baloban, M.; Pletnev, S.; Malashkevich, V. N.; Xiao, H.; Dauter, Z.; Verkhusha, V. V. Molecular Basis of Spectral Diversity in Near-Infrared Phytochrome-Based Fluorescent Proteins. *Chem. Biol.* **2015**, *22*, 1540-1551.
- (203) Shu, X.; Royant, A.; Lin, M. Z.; Aguilera, T. A.; Lev-Ram, V.; Steinbach, P. A.; Tsien, R. Y. Mammalian Expression of Infrared Fluorescent Proteins Engineered From a Bacterial Phytochrome. *Science*. **2009**, *324*, 804-807.
- (204) Rodriguez, E. A.; Tran, G. N.; Gross, L. A.; Crisp, J. L.; Shu, X.; Lin, J. Y.; Tsien, R. Y. A Far-red Fluorescent Protein Evolved From a Cyanobacterial Phycobiliprotein. *Nat. Methods.* **2016**, *13*, 763-769.
- (205) Li, E.; Norris, A. W. Structure/function of Cytoplasmic Vitamin A-binding Proteins. *Annu. Rev. Nutr.* **1996**, *16*, 205-234.

- (206) Crist, R. M.; Vasileiou, C.; Rabago-Smith, M.; Geiger, J. H.; Borhan, B. Engineering a Rhodopsin Protein Mimic. *J. Am. Chem. Soc.* **2006**, *128*, 4522-4523.
- (207) Vasileiou, C.; Vaezslami, S.; Crist, R. M.; Rabago-Smith, M.; Geiger, J. H.; Borhan, B. Protein Design: Reengineering Cellular Retinoic Acid Binding Protein II Into a Rhodopsin Protein Mimic. *J. Am. Chem. Soc.* **2007**, *129*, 6140-6148.
- (208) Gunasekaran, K.; Hagler, A. T.; Gierasch, L. M. Sequence and Structural Analysis of Cellular Retinoic Acid-binding Proteins Reveals a Network of Conserved Hydrophobic Interactions. *Proteins. Struct. Funct. Bioinf.* **2004**, *54*, 179-194.
- (209) Nosrati, M.; Berbasova, T.; Vasileiou, C.; Borhan, B.; Geiger, J. H. A Photoisomerizing Rhodopsin Mimic Observed at Atomic Resolution. *J. Am. Chem. Soc.* **2016**, *138*, 8802-8808.
- (210) Suomivuori, C. M.; Lang, L.; Sundholm, D.; Gamiz-Hernandez, A. P.; Kaila, V. R. Tuning the Protein-Induced Absorption Shifts of Retinal in Engineered Rhodopsin Mimics. *Chem. Eu. J.* **2016**
- (211) Huntress, M.; Gozem, S.; Malley, K. R.; Jailaubekov, A. E.; Vasileiou, C.; Vengris, M.; Geiger, J. H.; Borhan, B.; Schapiro, I.; Larsen, D.; et al. Towards An Understanding of the Retinal Chromophore in Rhodopsin Mimics. *J. Phys. Chem. B.* **2013**, *117*, 10053-10070.
- (212) Acharya, A.; Bogdanov, A. M.; Grigorenko, B. L.; Bravaya, K. B.; Nemukhin, A. V.; Lukyanov, K. A.; Krylov, A. I. Photoinduced Chemistry in Fluorescent Proteins: Curse or Blessing?. *Chem. Rev.* **2016**
- (213) Tsien, R. Y. The Green Fluorescent Protein. *Annu. Rev. Biochem.* **1998**, *67*, 509-544.
- (214) Chalfie, M. Green Fluorescent Protein As a Marker for Gene Expression. *Trends. Genet.* **1994**, *10*, 151.
- (215) Shaner, N. C.; Campbell, R. E.; Steinbach, P. A.; Giepmans, B. N. G.; Palmer, A. E.; Tsien, R. Y. Improved Monomeric Red, Orange and Yellow Fluorescent Proteins Derived From *Discosoma* Sp. Red Fluorescent Protein. *Nat. Biotech.* **2004**, *22*, 1567-1572.
- (216) Mizuno, H.; Mal, T. K.; Wälchli, M.; Kikuchi, A.; Fukano, T.; Ando, R.; Jayakanthan, J.; Taka, J.; Shiro, Y.; Ikura, M. Light-dependent Regulation of Structural Flexibility in a Photochromic Fluorescent Protein. *Proc. Natl. Acad. Sci. U. S. A.* **2008**, *105*, 9227-9232.
- (217) Shcherbakova, D. M.; Verkhusha, V. V. Chromophore Chemistry of Fluorescent Proteins Controlled by Light. *Curr. Opin. Chem. Biol.* **2014**, *20*, 60-68.
- (218) Martin, M. E.; Negri, F.; Olivucci, M. Origin, Nature, and Fate of the Fluorescent State of the Green Fluorescent Protein Chromophore at the CASPT2//CASCF Resolution. *J. Am. Chem. Soc.* **2004**, *126*, 5452-5464.
- (219) Altoè, P.; Bernardi, F.; Garavelli, M.; Orlandi, G.; Negri, F. Solvent Effects on the Vibrational Activity and Photodynamics of the Green Fluorescent Protein Chromophore: A Quantum-chemical Study. *J. Am. Chem. Soc.* **2005**, *127*, 3952-3963.
- (220) Olsen, S.; Smith, S. C. Bond Selection in the Photoisomerization Reaction of Anionic Green Fluorescent Protein and Kindling Fluorescent Protein Chromophore Models. *J. Am. Chem. Soc.* **2008**, *130*, 8677-8689.
- (221) Polyakov, I. V.; Grigorenko, B. L.; Epifanovsky, E. M.; Krylov, A. I.; Nemukhin, A. V. Potential Energy Landscape of the Electronic States of the GFP Chromophore in Different Protonation Forms: Electronic Transition Energies and Conical Intersections. *J. Chem. Theory. Comput.* **2010**, *6*, 2377-2387.
- (222) Toniolo, A.; Granucci, G.; Martínez, T. J. Conical Intersections in Solution: A QM/MM Study Using Floating Occupation Semiempirical Configuration Interaction Wave Functions. *J. Phys. Chem. A.* **2003**, *107*, 3822-3830.
- (223) Toniolo, A.; Olsen, S.; Manohar, L.; Martínez, T. J. Conical Intersection Dynamics in Solution: The Chromophore of Green Fluorescent Protein. *Faraday. Discuss.* **2004**, *127*, 149-163.
- (224) Pierce, D. W.; Hom-Booher, N.; Vale, R. D. Imaging Individual Green Fluorescent Proteins. *Nature.* **1997**, *388*, 338-338.
- (225) Dickson, R. M.; Cubitt, A. B.; Tsien, R. Y.; Moerner, W. E. On/off Blinking and Switching Behaviour of Single Molecules of Green Fluorescent Protein. *Nature.* **1997**, *388*, 355-358.
- (226) Haupts, U.; Maiti, S.; Schwill, P.; Webb, W. W. Dynamics of Fluorescence Fluctuations in Green Fluorescent Protein Observed by Fluorescence Correlation Spectroscopy. *Proc. Natl. Acad. Sci. U. S. A.* **1998**, *95*, 13573-13578.
- (227) Schwill, P.; Kummer, S.; Heikal, A. A.; Moerner, W. E.; Webb, W. W. Fluorescence Correlation Spectroscopy Reveals Fast Optical Excitation-driven Intramolecular Dynamics of Yellow Fluorescent Proteins. *Proc. Natl. Acad. Sci. U. S. A.* **2000**, *97*, 151-156.
- (228) Niwa, H.; Inouye, S.; Hirano, T.; Matsuno, T.; Kojima, S.; Kubota, M.; Ohashi, M.; Tsuji, F. I. Chemical Nature of the Light Emitter of the Aequorea Green Fluorescent Protein. *Proc. Natl. Acad. Sci. U. S. A.* **1996**, *93*, 13617-13622.
- (229) Meech, S. R. Excited State Reactions in Fluorescent Proteins. *Chem. Soc. Rev.* **2009**, *38*, 2922-2934.
- (230) Andersen, L. H.; Lapiere, A.; Nielsen, S. B.; Nielsen, I. B.; Pedersen, S. U.; Pedersen, U. V.; Tomita, S. Chromophores of the Green Fluorescent Protein Studied in the Gas Phase. *Eu. Phys. J. D.* **2002**, *20*, 597-600.
- (231) Baldrige, A.; Samanta, S. R.; Jayaraj, N.; Ramamurthy, V.; Tolbert, L. M. Steric and Electronic Effects in Capsule-confined Green Fluorescent Protein Chromophores. *J. Am. Chem. Soc.* **2011**, *133*, 712-715.
- (232) Baldrige, A.; Feng, S.; Chang, Y. -T.; Tolbert, L. M. Recapture of GFP Chromophore Fluorescence in a Protein Host. *ACS. Comb. Sci.* **2011**, *13*, 214-217.
- (233) Dedecker, P.; De Schryver, F. C.; Hofkens, J. Fluorescent Proteins: Shine On, You Crazy Diamond. *J. Am. Chem. Soc.* **2013**, *135*, 2387-2402.
- (234) Henderson, J. N.; Remington, S. J. The Kindling Fluorescent Protein: A Transient Photoswitchable Marker. *Physiology. (Bethesda).* **2006**, *21*, 162-170.
- (235) Stennett, E. M.; Ciuba, M. A.; Levitus, M. Photophysical Processes in Single Molecule Organic Fluorescent Probes. *Chem. Soc. Rev.* **2014**, *43*, 1057-1075.
- (236) Habuchi, S.; Ando, R.; Dedecker, P.; Verheijen, W.; Mizuno, H.; Miyawaki, A.; Hofkens, J. Reversible Single-molecule Photoswitching in the GFP-like Fluorescent Protein Dronpa. *Proc. Natl. Acad. Sci. U. S. A.* **2005**, *102*, 9511-9516.
- (237) Andresen, M.; Stiel, A. C.; Fölling, J.; Wenzel, D.; Schönle, A.; Egner, A.; Eggeling, C.; Hell, S. W.; Jakobs, S. Photoswitchable Fluorescent Proteins Enable Monochromatic Multilabel Imaging and Dual Color Fluorescence Nanoscopy. *Nat. Biotechnol.* **2008**, *26*, 1035-1040.
- (238) Chudakov, D. M.; Belousov, V. V.; Zaraisky, A. G.; Novoselov, V. V.; Staroverov, D. B.; Zorov, D. B.; Lukyanov, S.; Lukyanov, K. A. Kindling Fluorescent Proteins for Precise in Vivo Photolabeling. *Nat. Biotech.* **2003**, *21*, 191-194.
- (239) Li, X.; Chung, L. W.; Mizuno, H.; Miyawaki, A.; Morokuma, K. Primary Events of Photodynamics in Reversible Photoswitching Fluorescent Protein Dronpa. *J. Phys. Chem. Lett.* **2010**, *1*, 3328-3333.
- (240) Morozov, D.; Groenhof, G. Hydrogen Bond Fluctuations Control Photochromism in a Reversibly Photo-Switchable Fluorescent Protein. *Angew. Chem. Int. Ed. Engl.* **2016**, *55*, 576-578.
- (241) Cope, A. C.; Bach, R. D. Trans-Cyclooctene. *Org. Synth.* **1973**39-39.
- (242) Squillacote, M. E.; DeFellipis, J.; Shu, Q. How Stable Is Trans-cycloheptene?. *J. Am. Chem. Soc.* **2005**, *127*, 15983-15988.
- (243) Kandori, H.; Matuoka, S.; Shichida, Y.; Yoshizawa, T.; Ito, M.; Tsukida, K.; Balogh-Nair, V.; Nakanishi, K. Mechanism of Isomerization of Rhodopsin Studied by Use of 11-cis-locked Rhodopsin Analogues Excited with a Picosecond Laser Pulse. *Biochemistry.* **1989**, *28*, 6460-6467.
- (244) Kandori, H.; Furutani, Y.; Nishimura, S.; Shichida, Y.; Chosrowjan, H.; Shibata, Y.; Mataga, N. Excited-state Dynamics of

- Rhodopsin Probed by Femtosecond Fluorescence Spectroscopy. *Chem. Phys. Lett.* **2001**, *334*, 271-276.
- (245) Laricheva, E. N.; Gozem, S.; Rinaldi, S.; Melaccio, F.; Valentini, A.; Olivucci, M. Origin of Fluorescence in 11-cis Locked Bovine Rhodopsin. *J. Chem. Theory. Comput.* **2012**, *8*, 2559-2563.
- (246) Gozem, S.; Mirzakulova, E.; Schapiro, I.; Melaccio, F.; Glusac, K. D.; Olivucci, M. A Conical Intersection Controls the Deactivation of the Bacterial Luciferase Fluorophore. *Angew. Chem. Int. Ed. Engl.* **2014**, *53*, 9870-9875.
- (247) Lumento, F.; Zanirato, V.; Fusi, S.; Busi, E.; Latterini, L.; Elisei, F.; Sinicropi, A.; Andruniów, T.; Ferré, N.; Basosi, R.; et al. Quantum Chemical Modeling and Preparation of a Biomimetic Photochemical Switch. *Angew. Chem. Int. Ed. Engl.* **2007**, *46*, 414-420.
- (248) Sinicropi, A.; Martin, E.; Ryazantsev, M.; Helbing, J.; Briand, J.; Sharma, D.; Léonard, J.; Haacke, S.; Cannizzo, A.; Chergui, M.; et al. An Artificial Molecular Switch That Mimics the Visual Pigment and Completes Its Photocycle in Picoseconds. *Proc. Natl. Acad. Sci. U. S. A.* **2008**, *105*, 17642-17647.
- (249) Melloni, A.; Rossi Paccani, R.; Donati, D.; Zanirato, V.; Sinicropi, A.; Parisi, M. L.; Martin, E.; Ryazantsev, M.; Ding, W. J.; Frutos, L. M.; et al. Modeling, Preparation, and Characterization of a Dipole Moment Switch Driven by Z/E Photoisomerization. *J. Am. Chem. Soc.* **2010**, *132*, 9310-9319.
- (250) Rivado-Casas, L.; Sampedro, D.; Campos, P. J.; Fusi, S.; Zanirato, V.; Olivucci, M. Fluorenylidene-pyrroline Biomimetic Light-driven Molecular Switches. *J. Org. Chem.* **2009**, *74*, 4666-4674.
- (251) Blanco-Lomas, M.; Samanta, S.; Campos, P. J.; Woolley, G. A.; Sampedro, D. Reversible Photocontrol of Peptide Conformation with a Rhodopsin-like Photoswitch. *J. Am. Chem. Soc.* **2012**, *134*, 6960-6963.
- (252) Paolino, M.; Gueye, M.; Pieri, E.; Manathunga, M.; Fusi, S.; Cappelli, A.; Latterini, L.; Pannacci, D.; Filatov, M.; Léonard, J.; et al. Design, Synthesis and Dynamics of a GFP Fluorophore Mimic with An Ultrafast Switching Function. *J. Am. Chem. Soc.* **2016**, *138*, 9807-9825.
- (253) Merchán, M.; González-Luque, R.; Climent, T.; Serrano-Andrés, L.; Rodríguez, E.; Reguero, M.; Peláez, D. Unified Model for the Ultrafast Decay of Pyrimidine Nucleobases. *J. Phys. Chem. B.* **2006**, *110*, 26471-26476.
- (254) Lei, B.; Ding, Q.; Tu, S. C. Identity of the Emitter in the Bacterial Luciferase Luminescence Reaction: Binding and Fluorescence Quantum Yield Studies of 5-decyl-4a-hydroxy-4a,5-dihydro-10,10-dimethyl-5H-benzofluoren-5-yl-phosphate As a Model. *Biochemistry.* **2004**, *43*, 15975-15982.
- (255) Matsika, S. Radiationless Decay of Excited States of Uracil Through Conical Intersections. *J. Phys. Chem. A.* **2004**, *108*, 7584-7590.
- (256) Delchev, V. B.; Sobolewski, A. L.; Domcke, W. Comparison of the Non-radiative Decay Mechanisms of 4-pyrimidinone and Uracil: An Ab Initio Study. *Phys. Chem. Chem. Phys.* **2010**, *12*, 5007-5015.
- (257) Bernardi, F.; Olivucci, M.; Schegel, H. B. Michael Alfred Robb: A Short Biography. *Mol. Phys.* **2006**, *104*, 655-659.
- (258) Michl J., Bonačić-Koutecký V. *Electronic aspects of organic photochemistry* Wiley: New York, **1990**.
- (259) Bearpark, M. J.; Robb, M. A.; Schlegel, H. B. A Direct Method for the Location of the Lowest Energy Point on a Potential Surface Crossing. *Chem. Phys. Lett.* **1994**, *223*, 269-274.
- (260) Yarkony, D. R. Diabolical Conical Intersections. *Rev. Mod. Phys.* **1996**, *68*, 985.
- (261) Matsika, S.; Yarkony, D. R. Conical Intersections of Three Electronic States Affect the Ground State of Radical Species with Little or No Symmetry: Pyrazolyl. *J. Am. Chem. Soc.* **2003**, *125*, 12428-12429.
- (262) Maeda, S.; Ohno, K.; Morokuma, K. Updated Branching Plane for Finding Conical Intersections Without Coupling Derivative Vectors. *J. Chem. Theory. Comput.* **2010**, *6*, 1538-1545.
- (263) Birge, R. R. Photophysics and Molecular Electronic Applications of the Rhodopsins. *Annu. Rev. Phys. Chem.* **1990**, *41*, 683-733.
- (264) Birge, R. R. Nature of the Primary Photochemical Events in Rhodopsin and Bacteriorhodopsin. *Biochim. Biophys. Acta.* **1990**, *1016*, 293-327.
- (265) Sobolewski, A. L.; Domcke, W. Ab Initio Investigations on the Photophysics of Indole. *Chem. Phys. Lett.* **1999**, *315*, 293-298.
- (266) Manthe, U.; Köppel, H. Dynamics on Potential Energy Surfaces with a Conical Intersection: Adiabatic, Intermediate, and Diabatic Behavior. *J. Chem. Phys.* **1990**, *93*, 1658-1669.
- (267) Mahapatra, S.; Köppel, H.; Cederbaum, L. S. Impact of Nonadiabatic Coupling Between the Conically Intersecting X 2 A 1 and Å 2 B 2 States of NO 2 on the Negative Ion Photoelectron Spectra of NO 2-. *J. Chem. Phys.* **1999**, *110*, 5691-5701.
- (268) Martínez, T. J. Ab Initio Molecular Dynamics Around a Conical Intersection: Li (2p)+ H 2. *Chem. Phys. Lett.* **1997**, *272*, 139-147.
- (269) Olsen, J. The CASSCF Method: A Perspective and Commentary. *Int. J. Quantum. Chem.* **2011**, *111*, 3267-3272.
- (270) Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Scalmani, G.; Barone, V.; Petersson, G. A.; Nakatsuji, H.; et al. Gaussian 16 Revision A.03; 2016.
- (271) Aquilante, F.; Autschbach, J.; Carlson, R. K.; Chibotaru, L. F.; Delcey, M. G.; De Vico, L.; Fernandez Galvan, I.; Ferré, N.; Frutos, L. M.; Gagliardi, L. M.; et al. Molcas8: New Capabilities for Multi-configurational Quantum Chemical Calculations Across the Periodic Table. *J. Comp. Chem.* **2016**, *37*, 506-541.
- (272) Pulay, P. A Perspective on the CASPT2 Method. *Int. J. Quantum. Chem.* **2011**, *111*, 3273-3279.
- (273) Ruckebauer, M.; Barbatti, M.; Müller, T.; Lischka, H. Nonadiabatic Excited-state Dynamics with Hybrid Ab Initio Quantum-mechanical/molecular-mechanical Methods: Solvation of the Pentadieniminium Cation in Apolar Media. *J. Phys. Chem. A.* **2010**, *114*, 6757-6765.
- (274) Lischka, H.; Dallos, M.; Szalay, P. G.; Yarkony, D. R.; Shepard, R. Analytic Evaluation of Nonadiabatic Coupling Terms at the MR-CI Level. I. Formalism. *J. Chem. Phys.* **2004**, *120*, 7322-7329.
- (275) Dallos, M.; Lischka, H.; Shepard, R.; Yarkony, D. R.; Szalay, P. G. Analytic Evaluation of Nonadiabatic Coupling Terms at the MR-CI Level. II. Minima on the Crossing Seam: Formaldehyde and the Photodimerization of Ethylene. *J. Chem. Phys.* **2004**, *120*, 7330-7339.
- (276) Tao, H.; Levine, B. G.; Martínez, T. J. Ab Initio Multiple Spawning Dynamics Using Multi-state Second-order Perturbation Theory. *J. Phys. Chem. A.* **2009**, *113*, 13656-13662.
- (277) Angeli, C.; Borini, S.; Cestari, M.; Cimiriaglia, R. A Quasidegenerate Formulation of the Second Order N-electron Valence State Perturbation Theory Approach. *J. Chem. Phys.* **2004**, *121*, 4043-4049.
- (278) Freitag, L.; Knecht, S.; Angeli, C.; Reiher, M. Multireference Perturbation Theory with Cholesky Decomposition for the Density Matrix Renormalization Group. *J. Chem. Theory. Comput.* **2017**, *13*, 451-459.
- (279) Shiozaki, T.; Gyorffy, W.; Celani, P.; Werner, H. J. Communication: Extended Multi-state Complete Active Space Second-order Perturbation Theory: Energy and Nuclear Gradients. *J. Phys. Chem.* **2011**, *135*, 081106.
- (280) Vlaisavljevich, B.; Shiozaki, T. Nuclear Energy Gradients for Internally Contracted Complete Active Space Second-Order Perturbation Theory: Multistate Extensions. *J. Chem. Theory. Comput.* **2016**, *12*, 3781-3787.
- (281) Park, J. W.; Shiozaki, T. Analytical Derivative Coupling for Multistate CASPT2 Theory. *J. Chem. Theory. Comput.* **2017**, *13*, 2561-2570.
- (282) Park, J. W.; Shiozaki, T. On-the-fly CASPT2 Surface Hopping Dynamics. *J. Chem. Theory. Comput.* **2017**, *13*, 3676-3683.
- (283) Li, S. L.; Marenich, A. V.; Xu, X.; Truhlar, D. G. Configuration Interaction-Corrected Tamm-Dancoff Approximation: A Time-Dependent Density Functional Method with the Correct Dimensionality of Conical Intersections. *J. Phys. Chem. Lett.* **2014**, *5*, 322-328.

- (284) Wu, Q.; Cheng, C. L.; Van Voorhis, T. Configuration Interaction Based on Constrained Density Functional Theory: A Multireference Method. *J. Chem. Phys.* **2007**, *127*, 164119.
- (285) Wu, Q.; Kaduk, B.; Van Voorhis, T. Constrained Density Functional Theory Based Configuration Interaction Improves the Prediction of Reaction Barrier Heights. *J. Phys. Chem.* **2009**, *130*, 034109.
- (286) Huix-Rotllant, M.; Filatov, M.; Gozem, S.; Schapiro, I.; Olivucci, M.; Ferré, N. Assessment of Density Functional Theory for Describing the Correlation Effects on the Ground and Excited State Potential Energy Surfaces of a Retinal Chromophore Model. *J. Chem. Theory. Comput.* **2013**, *9*, 3917-3932.
- (287) Zhang, X.; Herbert, J. M. Analytic Derivative Couplings for Spin-flip Configuration Interaction Singles and Spin-flip Time-dependent Density Functional Theory. *J. Phys. Chem.* **2014**, *141*, 064104.
- (288) Herbert, J. M.; Zhang, X.; Morrison, A. F.; Liu, J. Beyond Time-Dependent Density Functional Theory Using Only Single Excitations: Methods for Computational Studies of Excited States in Complex Systems. *Acc. Chem. Res.* **2016**, *49*, 931-941.
- (289) Li Manni, G.; Carlson, R. K.; Luo, S.; Ma, D.; Olsen, J.; Truhlar, D. G.; Gagliardi, L. Multiconfiguration Pair-Density Functional Theory. *J. Chem. Theory. Comput.* **2014**, *10*, 3669-3680.
- (290) Epifanovsky, E.; Wormit, M.; Kuš, T.; Landau, A.; Zuev, D.; Khistyayev, K.; Manohar, P.; Kaliman, I.; Dreuw, A.; Krylov, A. I. New Implementation of High-level Correlated Methods Using a General Block Tensor Library for High-performance Electronic Structure Calculations. *J. Comput. Chem.* **2013**, *34*, 2293-2309.
- (291) Kaliman, I. A.; Krylov, A. I. New Algorithm for Tensor Contractions on Multi-core CPUs, GPUs, and Accelerators Enables CCSD and EOM-CCSD Calculations with Over 1000 Basis Functions on a Single Compute Node. *J. Comp. Chem.* **2017**, *38*, 842-853.
- (292) Krauter, C. M.; Pernpointner, M.; Dreuw, A. Application of the Scaled-opposite-spin Approximation to Algebraic Diagrammatic Construction Schemes of Second Order. *J. Phys. Chem.* **2013**, *138*, 044107.
- (293) Wormit, M.; Rehn, D. R.; Harbach, P. H.; Wenzel, J.; Krauter, C. M.; Epifanovsky, E.; Dreuw, A. Investigating Excited Electronic States Using the Algebraic Diagrammatic Construction (ADC) Approach of the Polarisation Propagator. *Mol. Phys.* **2014**, *112*, 774-784.
- (294) Lefrancois, D.; Wormit, M.; Dreuw, A. Adapting Algebraic Diagrammatic Construction Schemes for the Polarization Propagator to Problems with Multi-reference Electronic Ground States Exploiting the Spin-flip Ansatz. *J. Phys. Chem.* **2015**, *143*, 124107.
- (295) Zen, A.; Coccia, E.; Gozem, S.; Olivucci, M.; Guidoni, L. Quantum Monte Carlo Treatment of the Charge Transfer and Diradical Electronic Character in a Retinal Chromophore Minimal Model. *J. Chem. Theory. Comput.* **2015**, *11*, 992-1005.
- (296) Coccia, E.; Varsano, D.; Guidoni, L. Ab Initio Geometry and Bright Excitation of Carotenoids: Quantum Monte Carlo and Many Body Green's Function Theory Calculations on Peridinin. *J. Chem. Theory. Comput.* **2014**, *10*, 501-506.
- (297) Aquilante, F.; Malmqvist, P. Å.; Pedersen, T. B.; Ghosh, A.; Roos, B. O. Cholesky Decomposition-Based Multiconfiguration Second-Order Perturbation Theory (CD-CASPT2): Application to the Spin-State Energetics of Co(III)(diiminato)(NPh). *J. Chem. Theory. Comput.* **2008**, *4*, 694-702.
- (298) Delcey, M. G.; Pedersen, T. B.; Aquilante, F.; Lindh, R. Analytical Gradients of the State-average Complete Active Space Self-consistent Field Method with Density Fitting. *J. Phys. Chem.* **2015**, *143*, 044110.
- (299) Malmqvist, P. Å.; Rendell, A.; Roos, B. O. The Restricted Active Space Self-consistent-field Method, Implemented with a Split Graph Unitary Group Approach. *J. Phys. Chem.* **1990**, *94*, 5477-5482.
- (300) Vogiatzis, K. D.; Li Manni, G.; Stoneburner, S. J.; Ma, D.; Gagliardi, L. Systematic Expansion of Active Spaces Beyond the CASSCF Limit: A GASSCF/SplitGAS Benchmark Study. *J. Chem. Theory. Comput.* **2015**, *11*, 3010-3021.
- (301) Wouters, S.; Van Speybroeck, V.; Van Neck, D. DMRG-CASPT2 Study of the Longitudinal Static Second Hyperpolarizability of All-trans Polyenes. *J. Phys. Chem.* **2016**, *145*, 054120.
- (302) Roemelt, M.; Guo, S.; Chan, G. K. A Projected Approximation to Strongly Contracted N-electron Valence Perturbation Theory for DMRG Wavefunctions. *J. Phys. Chem.* **2016**, *144*, 204113.
- (303) Hedegård, E. D.; Reiher, M. Polarizable Embedding Density Matrix Renormalization Group. *J. Chem. Theory. Comput.* **2016**, *12*, 4242-4253.
- (304) Sjulstok, E.; Olsen, J. M.; Solov'yov, I. A. Quantifying Electron Transfer Reactions in Biological Systems: What Interactions Play the Major Role?. *Sci. Rep.* **2015**, *5*, 18446.
- (305) Guareschi, R.; Valsson, O.; Curutchet, C.; Mennucci, B.; Filippi, C. Electrostatic Versus Resonance Interactions in Photoreceptor Proteins: The Case of Rhodopsin. *J. Phys. Chem. Lett.* **2016**, *7*, 4547-4553.
- (306) Gordon, M. S.; Slipchenko, L.; Li, H.; Jensen, J. H. in *Annual reports in computational chemistry* Wheeler, R. and Spellmeyer, D. C., Eds.; Elsevier: Amsterdam, 2007; 3, pp 177-193.
- (307) Grigorenko, B. L.; Nemukhin, A. V.; Topol, I. A.; Burt, S. K. Modeling of Biomolecular Systems with the Quantum Mechanical and Molecular Mechanical Method Based on the Effective Fragment Potential Technique: Proposal of Flexible Fragments. *J. Phys. Chem. A.* **2002**, *106*, 10663-10672.
- (308) Gurunathan, P. K.; Acharya, A.; Ghosh, D.; Kosenkov, D.; Kaliman, I.; Shao, Y.; Krylov, A. I.; Slipchenko, L. V. Extension of the Effective Fragment Potential Method to Macromolecules. *J. Phys. Chem. B.* **2016**, *120*, 6562-6574.
- (309) Ufimtsev, I. S.; Martinez, T. J. Quantum Chemistry on Graphical Processing Units. 3. Analytical Energy Gradients, Geometry Optimization, and First Principles Molecular Dynamics. *J. Chem. Theory. Comput.* **2009**, *5*, 2619-2628.
- (310) Titov, A. V.; Ufimtsev, I. S.; Luehr, N.; Martinez, T. J. Generating Efficient Quantum Chemistry Codes for Novel Architectures. *J. Chem. Theory. Comput.* **2013**, *9*, 213-221.
- (311) Isborn, C. M.; Luehr, N.; Ufimtsev, I. S.; Martínez, T. J. Excited-State Electronic Structure with Configuration Interaction Singles and Tamm-Dancoff Time-Dependent Density Functional Theory on Graphical Processing Units. *J. Chem. Theory. Comput.* **2011**, *7*, 1814-1823.
- (312) Snyder Jr, J. W.; Curchod, B. F.; Martínez, T. J. GPU-Accelerated State-Averaged Complete Active Space Self-Consistent Field Interfaced with Ab Initio Multiple Spawning Unravels the Photodynamics of Provitamin D3. *J. Phys. Chem. Lett.* **2016**, *7*, 2444-2449.
- (313) Vacher, M.; Bearpark, M. J.; Robb, M. A.; Malhado, J. P. Electron Dynamics Upon Ionization of Polyatomic Molecules: Coupling to Quantum Nuclear Motion and Decoherence. *Phys. Rev. Lett.* **2017**, *118*, 083001.
- (314) Malhado, J. P.; Bearpark, M. J.; Hynes, J. T. Non-adiabatic Dynamics Close to Conical Intersections and the Surface Hopping Perspective. *Front. Chem.* **2014**, *2*, 97.
- (315) Calegari, F.; Ayuso, D.; Trabattini, A.; Belshaw, L.; De Camillis, S.; Anumula, S.; Frassetto, F.; Poletto, L.; Palacios, A.; Decleva, P.; et al. Ultrafast Electron Dynamics in Phenylalanine Initiated by Attosecond Pulses. *Science.* **2014**, *346*, 336-339.
- (316) Calegari, F.; Sansone, G.; Stagira, S.; Vozzi, C.; Nisoli, M. Advances in Attosecond Science. *J. Phys. Pt. B. Atom. M. P.* **2016**, *49*, 062001.
- (317) Fang, C.; Frontiera, R. R.; Tran, R.; Mathies, R. A. Mapping GFP Structure Evolution During Proton Transfer with Femtosecond Raman Spectroscopy. *Nature.* **2009**, *462*, 200-204.
- (318) Warren, M. M.; Kaucikas, M.; Fitzpatrick, A.; Champion, P.; Sage, J. T.; van Thor, J. J. Ground-state Proton Transfer in the Photoswitching Reactions of the Fluorescent Protein Dronpa. *Nat. Commun.* **2013**, *4*, 1461.
- (319) Luecke, H.; Schobert, B.; Cartailier, J. P.; Richter, H. T.; Rosengarth, A.; Needleman, R.; Lanyi, J. K. Coupling Photoisomeri-

- zation of Retinal to Directional Transport in Bacteriorhodopsin. *J. Mol. Biol.* **2000**, *300*, 1237-1255.
- (320) Melaccio, F.; Calimet, N.; Schapiro, I.; Valentini, A.; Cecchini, M.; Olivucci, M. Space and Time Evolution of the Electrostatic Potential During the Activation of a Visual Pigment. *J. Phys. Chem. Lett.* **2016**, *7*, 2563-2567.
- (321) van Keulen, S. C.; Solano, A.; Rothlisberger, U. How Rhodopsin Tunes the Equilibrium Between Protonated and Deprotonated Forms of the Retinal Chromophore. *J. Chem. Theory. Comput.* **2017**
- (322) Goyal, P.; Hammes-Schiffer, S. Role of Active Site Conformational Changes in Photocycle Activation of the AppA BLUF Photoreceptor. *Proc. Natl. Acad. Sci. U. S. A.* **2017**, *114*, 1480-1485.
- (323) Shalabney, A.; George, J.; Hutchison, J.; Pupillo, G.; Genet, C.; Ebbesen, T. W. Coherent Coupling of Molecular Resonators with a Microcavity Mode. *Nat. Commun.* **2015**, *6*, 5981.
- (324) Törmä, P.; Barnes, W. L. Strong Coupling Between Surface Plasmon Polaritons and Emitters: A Review. *Rep. Prog. Phys.* **2015**, *78*, 013901.
- (325) Chang, B. S. W.; Jonsson, K.; Kazmi, M. A.; Donoghue, M. J. Recreating a Functional Ancestral Archosaur Visual Pigment. *Mol. Biol. Evol.* **2002**, *19*, 1483-1489.
- (326) Sekharan, S.; Yokoyama, S.; Morokuma, K. Quantum Mechanical/molecular Mechanical Structure, Enantioselectivity, and Spectroscopy of Hydroxyretinals and Insights Into the Evolution of Color Vision in Small White Butterflies. *J. Phys. Chem. B.* **2011**, *115*, 15380-15388.
- (327) Altun, A.; Morokuma, K.; Yokoyama, S. H-bond Network Around Retinal Regulates the Evolution of Ultraviolet and Violet Vision. *ACS. Chem. Biol.* **2011**, *6*, 775-780.
- (328) Watanabe, H. C.; Mori, Y.; Tada, T.; Yokoyama, S.; Yamato, T. Molecular Mechanism of Long-range Synergetic Color Tuning Between Multiple Amino Acid Residues in Conger Rhodopsin. *Biophys. (Oxford)*. **2010**, *6*, 67-68.
- (329) Altun, A.; Yokoyama, S.; Morokuma, K. Color Tuning in Short Wavelength-sensitive Human and Mouse Visual Pigments: Ab Initio Quantum Mechanics/molecular Mechanics Studies. *J. Phys. Chem. A.* **2009**, *113*, 11685-11692.
- (330) Melaccio, F.; del Carmen Marín, M.; Valentini, A.; Montisci, F.; Rinaldi, S.; Cherubini, M.; Yang, X.; Kato, Y.; Stenrup, M.; Orozco-Gonzalez, Y.; et al. Towards Automatic Rhodopsin Modeling As a Tool for High-throughput Computational Photobiology. *J. Chem. Theory. Comput.* **2016**, *12*, 6020-6034.
- (331) Ross, B. in *Chemistry, 1996-2000* Grenthe, I., Ed.; World Scientific: Singapore, 2003; pp 197-213.
- (332) Ochterski, J. W.; Petersson, G. A.; Wiberg, K. B. A Comparison of Model Chemistries. *J. Am. Chem. Soc.* **1995**, *117*, 11299-11308.
- (333) Dreuw, A.; Head-Gordon, M. Single-reference Ab Initio Methods for the Calculation of Excited States of Large Molecules. *Chem. Rev.* **2005**, *105*, 4009-4037.
- (334) Serrano-Andrés, L.; Merchán, M. Quantum Chemistry of the Excited State: 2005 Overview. *J. Mol. Struct.* **2005**, *729*, 99-108.
- (335) Vreven, T.; Bernardi, F.; Garavelli, M.; Olivucci, M.; Robb, M. A.; Schlegel, H. B. Ab Initio Photoisomerization Dynamics of a Simple Retinal Chromophore Model. *J. Am. Chem. Soc.* **1997**, *119*, 12687-12688.
- (336) Migani, A.; Robb, M. A.; Olivucci, M. Relationship Between Photoisomerization Path and Intersection Space in a Retinal Chromophore Model. *J. Am. Chem. Soc.* **2003**, *125*, 2804-2808.
- (337) Fantacci, S.; Migani, A.; Olivucci, M. CASPT2//CASSCF and TDDFT//CASSCF Mapping of the Excited State Isomerization Path of a Minimal Model of the Retinal Chromophore. *J. Phys. Chem. A.* **2004**, *108*, 1208-1213.
- (338) Weingart, O.; Migani, A.; Olivucci, M.; Robb, M. A.; Buss, V.; Hunt, P. Probing the Photochemical Funnel of a Retinal Chromophore Model Via Zero-point Energy Sampling Semiclassical Dynamics. *J. Phys. Chem. A.* **2004**, *108*, 4685-4693.
- (339) Weingart, O.; Buss, V.; Robb, M. A. Excited State Molecular Dynamics of Retinal Model Chromophores. *Phase. Transitions.* **2005**, *78*, 17-24.
- (340) Wanko, M.; Garavelli, M.; Bernardi, F.; Niehaus, T. A.; Frauenheim, T.; Elstner, M. A Global Investigation of Excited State Surfaces Within Time-dependent Density-functional Response Theory. *J. Chem. Phys.* **2004**, *120*, 1674-1692.
- (341) Wanko, M.; Hoffmann, M.; Strodel, P.; Koslowski, A.; Thiel, W.; Neese, F.; Frauenheim, T.; Elstner, M. Calculating Absorption Shifts for Retinal Proteins: Computational Challenges. *J. Phys. Chem. B.* **2005**, *109*, 3606-3615.
- (342) Martin, C. H.; Birge, R. R. Reparametrizing MNDO for Excited-state Calculations by Using Ab Initio Effective Hamiltonian Theory: Application to the 2, 4-pentadien-1-iminium Cation. *J. Phys. Chem. A.* **1998**, *102*, 852-860.
- (343) Keal, T. W.; Wanko, M.; Thiel, W. Assessment of Semiempirical Methods for the Photoisomerisation of a Protonated Schiff Base. *Theor. Chem. Acc.* **2009**, *123*, 145-156.
- (344) Send, R.; Sundholm, D.; Johansson, M. P.; Pawłowski, F. Excited State Potential Energy Surfaces of Polyenes and Protonated Schiff Bases. *J. Chem. Theory. Comput.* **2009**, *5*, 2401-2414.
- (345) Lehtonen, O.; Sundholm, D.; Send, R.; Johansson, M. P. Coupled-cluster and Density Functional Theory Studies of the Electronic Excitation Spectra of Trans-1,3-butadiene and Trans-2-propeniminium. *J. Phys. Chem.* **2009**, *131*, 024301.
- (346) Valsson, O.; Filippi, C. Photoisomerization of Model Retinal Chromophores: Insight From Quantum Monte Carlo and Multiconfigurational Perturbation Theory. *J. Chem. Theory. Comput.* **2010**, *6*, 1275-1292.
- (347) Valsson, O.; Filippi, C.; Casida, M. E. Regarding the Use and Misuse of Retinal Protonated Schiff Base Photochemistry As a Test Case for Time-dependent Density-functional Theory. *J. Phys. Chem.* **2015**, *142*, 144104.
- (348) Walczak, E.; Szeferczyk, B.; Andruniów, T. Geometries and Vertical Excitation Energies in Retinal Analogues Resolved at the CASPT2 Level of Theory: Critical Assessment of the Performance of CASSCF, CC2, and DFT Methods. *J. Chem. Theory. Comput.* **2013**, *9*, 4915-4927.
- (349) Grabarek, D.; Walczak, E.; Andruniów, T. Assessing the Accuracy of Various Ab Initio Methods for Geometries and Excitation Energies of Retinal Chromophore Minimal Model by Comparison with CASPT3 Results. *J. Chem. Theory. Comput.* **2016**, *12*, 2346-2356.
- (350) Zaari, R. R.; Wong, S. Y. Photoexcitation of 11-Z-cis-7, 8-dihydro Retinal and 11-Z-cis Retinal: A Comparative Computational Study. *Chem. Phys. Lett.* **2009**, *469*, 224-228.
- (351) Strambi, A.; Coto, P. B.; Ferré, N.; Olivucci, M. Effects of Water Re-location and Cavity Trimming on the CASPT2//CASSCF/AMBER Excitation Energy of Rhodopsin. *Theor. Chem. Acc.* **2007**, *118*, 185-191.
- (352) Gozem, S.; Krylov, A. I.; Olivucci, M. Conical Intersection and Potential Energy Surface Features of a Model Retinal Chromophore: Comparison of EOM-CC and Multireference Methods. *J. Chem. Theory. Comput.* **2013**, *9*, 284-292.
- (353) Schapiro, I.; Neese, F. SORCI for Photochemical and Thermal Reaction Paths: A Benchmark Study. *Comput. Theor. Chem.* **2014**, *1040*, 84-98.
- (354) Tuna, D.; Lefrancois, D.; Wolański, Ł.; Gozem, S.; Schapiro, I.; Andruniów, T.; Dreuw, A.; Olivucci, M. Assessment of Approximate Coupled-Cluster and Algebraic-Diagrammatic-Construction Methods for Ground- and Excited-State Reaction Paths and the Conical-Intersection Seam of a Retinal-Chromophore Model. *J. Chem. Theory. Comput.* **2015**, *11*, 5758-5781.
- (355) Xu, X.; Gozem, S.; Olivucci, M.; Truhlar, D. G. Combined Self-Consistent-Field and Spin-Flip Tamm-Dancoff Density Functional Approach to Potential Energy Surfaces for Photochemistry. *J. Phys. Chem. Lett.* **2013**, *4*, 253-258.

- (356) Robb, M. A.; Garavelli, M.; Olivucci, M.; Bernardi, F. in *Reviews in computational chemistry* Lipkowitz, K. B. and Boyd, D. B., Eds.; Wiley: New York, 2000; 15, pp 87-146.
- (357) Han, S.; Yarkony, D. R. Nonadiabatic Processes Involving Three Electronic States. I. Branch Cuts and Linked Pairs of Conical Intersections. *J. Chem. Phys.* **2003**, *119*, 5058-5068.
- (358) Fdez Galván, I.; Delcey, M. G.; Pedersen, T. B.; Aquilante, F.; Lindh, R. Analytical State-Average Complete-Active-Space Self-Consistent Field Nonadiabatic Coupling Vectors: Implementation with Density-Fitted Two-Electron Integrals and Application to Conical Intersections. *J. Chem. Theory. Comput.* **2016**, *12*, 3636-3653.
- (359) Wigner E. P., Wightman A. S. *The Collected Works of Eugene Paul Wigner Springer*: Berlin, **1993**.
- (360) Teller, E. The Crossing of Potential Surfaces. *J. Phys. Chem.* **1937**, *41*, 109-116.
- (361) Hättig, C. in *Advances in quantum chemistry* Jensen, H., Ed.; Elsevier: Amsterdam, 2005; 50, pp 37-60.
- (362) Köhn, A.; Tajiri, A. Can Coupled-cluster Theory Treat Conical Intersections?. *J. Chem. Phys.* **2007**, *127*, 044105.
- (363) Blancafort, L.; Ogliaro, F.; Olivucci, M.; Robb, M. A.; Bearpark, M. J.; Sinicropi, A. Computational Investigation of Photochemical Reaction Mechanisms. *Mol. Supramol. Photochem.* **2005**, *13*, 31.
- (364) Marian, C. M. Spin-orbit Coupling and Intersystem Crossing in Molecules. *Wiley Interdiscip. Rev. Comp. Mol. Sci.* **2012**, *2*, 187-203.
- (365) Schautz, F.; Filippi, C. Optimized Jastrow-Slater Wave Functions for Ground and Excited States: Application to the Lowest States of Ethene. *J. Chem. Phys.* **2004**, *120*, 10931-10941.
- (366) Schautz, F.; Buda, F.; Filippi, C. Excitations in Photoactive Molecules From Quantum Monte Carlo. *J. Chem. Phys.* **2004**, *121*, 5836-5844.
- (367) Varsano, D.; Caprasecca, S.; Coccia, E. Theoretical Description of Protein Field Effects on Electronic Excitations of Biological Chromophores. *J. Phys. Condens. Matter.* **2017**, *29*, 013002.
- (368) Ridley, J.; Zerner, M. An Intermediate Neglect of Differential Overlap Technique for Spectroscopy: Pyrrole and the Azines. *Theo. Chim. Acta.* **1973**, *32*, 111-134.
- (369) Ridley, J. E.; Zerner, M. C. Triplet States Via Intermediate Neglect of Differential Overlap: Benzene, Pyridine and the Diazines. *Theo. Chim. Acta.* **1976**, *42*, 223-236.
- (370) Dewar, M. J.; Thiel, W. Ground States of Molecules. 38. The MNDO Method. Approximations and Parameters. *J. Am. Chem. Soc.* **1977**, *99*, 4899-4907.
- (371) Schweig, A.; Thiel, W. MNDOC Study of Excited States. *J. Am. Chem. Soc.* **1981**, *103*, 1425-1431.
- (372) Stewart, J. J. Optimization of Parameters for Semiempirical Methods I. Method. *J. Comp. Chem.* **1989**, *10*, 209-220.
- (373) Dewar, M. J.; Zoebisch, E. G.; Healy, E. F.; Stewart, J. J. Development and Use of Quantum Mechanical Molecular Models. 76. AM1: A New General Purpose Quantum Mechanical Molecular Model. *J. Am. Chem. Soc.* **1985**, *107*, 3902-3909.
- (374) Kolb, M.; Thiel, W. Beyond the MNDO Model: Methodical Considerations and Numerical Results. *J. Comp. Chem.* **1993**, *14*, 775-789.
- (375) Weber, W.; Thiel, W. Orthogonalization Corrections for Semiempirical Methods. *Theor. Chem. Acc.* **2000**, *103*, 495-506.
- (376) Hoffmann, M.; Wanko, M.; Strodel, P.; König, P. H.; Frauenheim, T.; Schulten, K.; Thiel, W.; Tajkhorshid, E.; Elstner, M. Color Tuning in Rhodopsins: The Mechanism of the Spectral Shift Between Bacteriorhodopsin and Sensory Rhodopsin II. *J. Am. Chem. Soc.* **2006**, *128*, 10808-10818.
- (377) Silva-Junior, M. R.; Thiel, W. Benchmark of Electronically Excited States for Semiempirical Methods: MNDO, AM1, PM3, OM1, OM2, OM3, INDO/S, and INDO/S2. *J. Chem. Theory. Comput.* **2010**, *6*, 1546-1564.
- (378) Löwdin, P. -O. Quantum Theory of Many-particle Systems. III. Extension of the Hartree-Fock Scheme to Include Degenerate Systems and Correlation Effects. *Phys. Rev.* **1955**, *97*, 1509.
- (379) Langhoff, S. R.; Davidson, E. R. Configuration Interaction Calculations on the Nitrogen Molecule. *Int. J. Quantum. Chem.* **1974**, *8*, 61-72.
- (380) Andersson, K.; Malmqvist, P. -Å.; Roos, B. O. Second-order Perturbation Theory with a Complete Active Space Self-consistent Field Reference Function. *J. Chem. Phys.* **1992**, *96*, 1218-1226.
- (381) Finley, J.; Malmqvist, P. -Å.; Roos, B. O.; Serrano-Andrés, L. The Multi-state CASPT2 Method. *Chem. Phys. Lett.* **1998**, *288*, 299-306.
- (382) Angeli, C.; Cimraglia, R.; Evangelisti, S.; Leininger, T.; Malrieu, J. -P. Introduction of N-electron Valence States for Multireference Perturbation Theory. *J. Chem. Phys.* **2001**, *114*, 10252-10264.
- (383) Angeli, C.; Pastore, M.; Cimraglia, R. New Perspectives in Multireference Perturbation Theory: The N-electron Valence State Approach. *Theor. Chem. Acc.* **2007**, *117*, 743-754.
- (384) Ghigo, G.; Roos, B. O.; Malmqvist, P. K. A Modified Definition of the Zeroth-order Hamiltonian in Multiconfigurational Perturbation Theory (CASPT2). *Chem. Phys. Lett.* **2004**, *396*, 142-149.
- (385) Angeli, C.; Pastore, M. The Lowest Singlet States of Octatetraene Revisited. *J. Phys. Chem.* **2011**, *134*, 184302.
- (386) Kepenekian, M.; Robert, V.; Le Guennic, B. What Zeroth-order Hamiltonian for CASPT2 Adiabatic Energetics of Fe(II)N(6) Architectures?. *J. Phys. Chem.* **2009**, *131*, 114702.
- (387) Zobel, J. P.; Nogueira, J. J.; González, L. The IPEA Dilemma in CASPT2. *J. Chem. Sci.* **2017**, *8*, 1482-1499.
- (388) Rintelman, J. M.; Adamovic, I.; Varganov, S.; Gordon, M. S. Multireference Second-order Perturbation Theory: How Size Consistent Is almost Size Consistent. *J. Chem. Phys.* **2005**, *122*, 044105.
- (389) Roca-Sanjuán, D.; Aquilante, F.; Lindh, R. Multiconfiguration Second-order Perturbation Theory Approach to Strong Electron Correlation in Chemistry and Photochemistry. *Wiley Interdiscip. Rev. Comp. Mol. Sci.* **2012**, *2*, 585-603.
- (390) Neese, F. A Spectroscopy Oriented Configuration Interaction Procedure. *J. Chem. Phys.* **2003**, *119*, 9428-9443.
- (391) Krylov, A. I. Equation-of-motion Coupled-cluster Methods for Open-shell and Electronically Excited Species: The Hitchhiker's Guide to Fock Space. *Annu. Rev. Phys. Chem.* **2008**, *59*, 433-462.
- (392) Stanton, J. F.; Bartlett, R. J. The Equation of Motion Coupled-cluster Method. A Systematic Biorthogonal Approach to Molecular Excitation Energies, Transition Probabilities, and Excited State Properties. *J. Chem. Phys.* **1993**, *98*, 7029-7039.
- (393) Stanton, J. F.; Gauss, J. Analytic Energy Derivatives for Ionized States Described by the Equation-of-motion Coupled Cluster Method. *J. Chem. Phys.* **1994**, *101*, 8938-8944.
- (394) Nooijen, M.; Bartlett, R. J. Equation of Motion Coupled Cluster Method for Electron Attachment. *J. Chem. Phys.* **1995**, *102*, 3629-3647.
- (395) Krylov, A. I. Spin-flip Equation-of-motion Coupled-cluster Electronic Structure Method for a Description of Excited States, Bond Breaking, Diradicals, and Triradicals. *Acc. Chem. Res.* **2006**, *39*, 83-91.
- (396) Krylov, A. I. Size-consistent Wave Functions for Bond-breaking: The Equation-of-motion Spin-flip Model. *Chem. Phys. Lett.* **2001**, *338*, 375-384.
- (397) Slipchenko, L. V.; Krylov, A. I. Singlet-triplet Gaps in Diradicals by the Spin-flip Approach: A Benchmark Study. *J. Chem. Phys.* **2002**, *117*, 4694-4708.
- (398) Levchenko, S. V.; Krylov, A. I. Equation-of-motion Spin-flip Coupled-cluster Model with Single and Double Substitutions: Theory and Application to Cyclobutadiene. *J. Chem. Phys.* **2004**, *120*, 175-185.
- (399) Schirmer, J. Beyond the Random-phase Approximation: A New Approximation Scheme for the Polarization Propagator. *Phys. Rev. A.* **1982**, *26*, 2395.
- (400) Trofimov, A. B.; Schirmer, J. An Efficient Polarization Propagator Approach to Valence Electron Excitation Spectra. *J. Phys. Pt. B. Atom. M. P.* **1995**, *28*, 2299.

- (401) Dreuw, A.; Wormit, M. The Algebraic Diagrammatic Construction Scheme for the Polarization Propagator for the Calculation of Excited States. *Wiley. Interdiscip. Rev. Comp. Mol. Sci.* **2015**, *5*, 82-95.
- (402) Casida, M. E. Time-dependent Density-functional Theory for Molecules and Molecular Solids. *J. Mol. Struct. THEOCHEM.* **2009**, *914*, 3-18.
- (403) Dreuw, A.; Head-Gordon, M. Failure of Time-dependent Density Functional Theory for Long-range Charge-transfer Excited States: The Zincbacteriochlorin-bacteriochlorin and Bacteriochlorophyll-spheroidene Complexes. *J. Am. Chem. Soc.* **2004**, *126*, 4007-4016.
- (404) Cordova, F.; Doriol, L. J.; Ipatov, A.; Casida, M. E.; Filippi, C.; Vela, A. Troubleshooting Time-dependent Density-functional Theory for Photochemical Applications: Oxirane. *J. Chem. Phys.* **2007**, *127*, 164111.
- (405) Laurent, A. D.; Jacquemin, D. TD-DFT Benchmarks: A Review. *Int. J. Quantum. Chem.* **2013**, *113*, 2019-2039.
- (406) Jacquemin, D.; Wathelet, V.; Perpète, E. A.; Adamo, C. Extensive TD-DFT Benchmark: Singlet-Excited States of Organic Molecules. *J. Chem. Theory. Comput.* **2009**, *5*, 2420-2435.
- (407) Gozem, S.; Melaccio, F.; Luk, H. L.; Rinaldi, S.; Olivucci, M. Learning From Photobiology How to Design Molecular Devices Using a Computer. *Chem. Soc. Rev.* **2014**, *43*, 4019-4036.
- (408) Shao, Y.; Head-Gordon, M.; Krylov, A. I. The Spin-flip Approach Within Time-dependent Density Functional Theory: Theory and Applications to Diradicals. *J. Chem. Phys.* **2003**, *118*, 4807-4818.
- (409) Marti, K. H.; Ondík, I. M.; Moritz, G.; Reiher, M. Density Matrix Renormalization Group Calculations on Relative Energies of Transition Metal Complexes and Clusters. *J. Chem. Phys.* **2008**, *128*, 014104.
- (410) Chan, G. K. -L.; Head-Gordon, M. Highly Correlated Calculations with a Polynomial Cost Algorithm: A Study of the Density Matrix Renormalization Group. *J. Chem. Phys.* **2002**, *116*, 4462-4476.
- (411) Chan, G. K. An Algorithm for Large Scale Density Matrix Renormalization Group Calculations. *J. Chem. Phys.* **2004**, *120*, 3172-3178.
- (412) Keller, S.; Dolfi, M.; Troyer, M.; Reiher, M. An Efficient Matrix Product Operator Representation of the Quantum Chemical Hamiltonian. *J. Phys. Chem.* **2015**, *143*, 244118.
- (413) Knecht, S.; Hedegård, E. D.; Keller, S.; Kovyrshin, A.; Ma, Y.; Muolo, A.; Stein, C. J.; Reiher, M. New Approaches for Ab Initio Calculations of Molecules with Strong Electron Correlation. *Chimia.* **2016**, *70*, 244-251.
- (414) Wouters, S.; Poelmans, W.; Ayers, P. W.; Van Neck, D. ChemPS2: A Free Open-source Spin-adapted Implementation of the Density Matrix Renormalization Group for Ab Initio Quantum Chemistry. *Comput. Phys. Comm.* **2014**, *185*, 1501-1514.
- (415) Guo, S.; Watson, M. A.; Hu, W.; Sun, Q.; Chan, G. K. -L. N-Electron Valence State Perturbation Theory Based on a Density Matrix Renormalization Group Reference Function, with Applications to the Chromium Dimer and a Trimer Model of Poly (p-Phenylenevinylene). *J. Chem. Theory. Comput.* **2016**, *12*, 1583-1591.
- (416) Stein, C. J.; Reiher, M. Automated Selection of Active Orbital Spaces. *J. Chem. Theory. Comput.* **2016**, *12*, 1760-1771.
- (417) Warshel, A.; Levitt, M. Theoretical Studies of Enzymic Reactions: Dielectric, Electrostatic and Steric Stabilization of the Carbonium Ion in the Reaction of Lysozyme. *J. Mol. Biol.* **1976**, *103*, 227-249.
- (418) Ferre, N.; Cembran, A.; Garavelli, M.; Olivucci, M. Complete-active-space Self-consistent-field/Amber Parameterization of the Lys296-retinal-Glu113 Rhodopsin Chromophore-counterion System. *Theor. Chem. Acc.* **2004**, *112*, 335-341.
- (419) Stoner-Ma, D.; Jaye, A. A.; Ronayne, K. L.; Nappa, J.; Meech, S. R.; Tonge, P. J. An Alternate Proton Acceptor for Excited-state Proton Transfer in Green Fluorescent Protein: Rewiring GFP. *J. Am. Chem. Soc.* **2008**, *130*, 1227-1235.
- (420) Zhang, Q.; Chen, X.; Cui, G.; Fang, W. H.; Thiel, W. Concerted Asynchronous Hula-twist Photoisomerization in the S65T/H148D Mutant of Green Fluorescent Protein. *Angew. Chem. Int. Ed. Engl.* **2014**, *53*, 8649-8653.
- (421) Valsson, O.; Campomanes, P.; Tavernelli, I.; Rothlisberger, U.; Filippi, C. Rhodopsin Absorption From First Principles: Bypassing Common Pitfalls. *J. Chem. Theory. Comput.* **2013**, *9*, 2441-2454.
- (422) Senn, H. M.; Thiel, W. QM/MM Methods for Biomolecular Systems. *Angew. Chem. Int. Ed. Engl.* **2009**, *48*, 1198-1229.
- (423) Ji, C.; Mei, Y. Some Practical Approaches to Treating Electrostatic Polarization of Proteins. *Acc. Chem. Res.* **2014**, *47*, 2795-2803.
- (424) Baker, C. M. Polarizable Force Fields for Molecular Dynamics Simulations of Biomolecules. *Wiley. Interdiscip. Rev. Comp. Mol. Sci.* **2015**, *5*, 241-254.
- (425) Daday, C.; Curutchet, C.; Sinicropi, A.; Mennucci, B.; Filippi, C. Chromophore-Protein Coupling Beyond Nonpolarizable Models: Understanding Absorption in Green Fluorescent Protein. *J. Chem. Theory. Comput.* **2015**, *11*, 4825-4839.
- (426) Jacob, C. R.; Neugebauer, J. Subsystem Density-functional Theory. *Wiley. Interdiscip. Rev. Comp. Mol. Sci.* **2014**, *4*, 325-362.
- (427) Schapiro, I.; Roca-Sanjuán, D.; Lindh, R.; Olivucci, M. A Surface Hopping Algorithm for Nonadiabatic Minimum Energy Path Calculations. *J. Comp. Chem.* **2015**, *36*, 312-320.
- (428) Singh, U. C.; Kollman, P. A. A Combined Ab Initio Quantum Mechanical and Molecular Mechanical Method for Carrying Out Simulations on Complex Molecular Systems: Applications to the CH₃Cl+ Cl- Exchange Reaction and Gas Phase Protonation of Polyethers. *J. Comp. Chem.* **1986**, *7*, 718-730.
- (429) Field, M. J.; Bash, P. A.; Karplus, M. A Combined Quantum Mechanical and Molecular Mechanical Potential for Molecular Dynamics Simulations. *J. Comp. Chem.* **1990**, *11*, 700-733.
- (430) Dapprich, S.; Komáromi, I.; Byun, K. S.; Morokuma, K.; Frisch, M. J. A New ONIOM Implementation in Gaussian98. Part I. The Calculation of Energies, Gradients, Vibrational Frequencies and Electric Field Derivatives. *J. Mol. Struct. THEOCHEM.* **1999**, *461*, 1-21.
- (431) Svensson, M.; Humbel, S.; Froese, R. D.; Matsubara, T.; Sieber, S.; Morokuma, K. ONIOM: A Multilayered Integrated MO+ MM Method for Geometry Optimizations and Single Point Energy Predictions. A Test for Diels-Alder Reactions and Pt (P (t-Bu)₃)₂+ H₂ Oxidative Addition. *J. Phys. Chem.* **1996**, *100*, 19357-19363.
- (432) Humbel, S.; Sieber, S.; Morokuma, K. The IMOMO Method: Integration of Different Levels of Molecular Orbital Approximations for Geometry Optimization of Large Systems: Test for N-butane Conformation and SN₂ Reaction: RCl+ Cl-. *J. Phys. Chem.* **1996**, *105*, 1959-1967.
- (433) Vreven, T.; Morokuma, K. On the Application of the IMOMO (integrated Molecular Orbital+ Molecular Orbital) Method. *J. Comp. Chem.* **2000**, *21*, 1419-1432.
- (434) Maseras, F.; Morokuma, K. IMOMM: A New Integrated Ab Initio+ Molecular Mechanics Geometry Optimization Scheme of Equilibrium Structures and Transition States. *J. Comp. Chem.* **1995**, *16*, 1170-1179.
- (435) Vreven, T.; Morokuma, K. in *Annual Reports in Computational Chemistry* Spellmeyer, D. C., Ed.; Elsevier: Amsterdam, 2006; *2*, pp 35-51.
- (436) Antes, I.; Thiel, W. Adjusted Connection Atoms for Combined Quantum Mechanical and Molecular Mechanical Methods. *J. Phys. Chem. A.* **1999**, *103*, 9290-9295.
- (437) Théry, V.; Rinaldi, D.; Rivail, J. -L.; Maigret, B.; Ferenczy, G. G. Quantum Mechanical Computations on Very Large Molecular Systems: The Local Self-consistent Field Method. *J. Comp. Chem.* **1994**, *15*, 269-282.
- (438) Woo, T. K.; Cavallo, L.; Ziegler, T. Implementation of the IMOMM Methodology for Performing Combined QM/MM Molecular Dynamics Simulations and Frequency Calculations. *Theor. Chem. Acc.* **1998**, *100*, 307-313.
- (439) Sherwood, P.; de Vries, A. H.; Collins, S. J.; Greatbanks, S. P.; Burton, N. A.; Vincent, M. A.; Hillier, I. H. Computer Simulation of

- Zeolite Structure and Reactivity Using Embedded Cluster Methods. *Faraday. Discuss.* **1997**, *106*, 79-92.
- (440) Berman, H. M.; Westbrook, J.; Feng, Z.; Gilliland, G.; Bhat, T. N.; Weissig, H.; Shindyalov, I. N.; Bourne, P. E. The Protein Data Bank. *Nucleic. Acids. Res.* **2000**, *28*, 235-242.
- (441) Zhou, Y.; Wang, S.; Li, Y.; Zhang, Y. Chapter Five-Born--Oppenheimer Ab Initio QM/MM Molecular Dynamics Simulations of Enzyme Reactions. *Method. Enzymol.* **2016**, *577*, 105-118.
- (442) Barbatti, M. Nonadiabatic Dynamics with Trajectory Surface Hopping Method. *Wiley. Interdiscip. Rev. Comp. Mol. Sci.* **2011**, *1*, 620-633.
- (443) Subotnik, J. E.; Jain, A.; Landry, B.; Petit, A.; Ouyang, W.; Bellonzi, N. Understanding the Surface Hopping View of Electronic Transitions and Decoherence. *Annu. Rev. Phys. Chem.* **2016**, *67*, 387-417.
- (444) Groenhof, G.; Schäfer, L. V.; Boggio-Pasqua, M.; Goette, M.; Grubmüller, H.; Robb, M. A. Ultrafast Deactivation of An Excited Cytosine-guanine Base Pair in DNA. *J. Am. Chem. Soc.* **2007**, *129*, 6812-6819.
- (445) Paterson, M. J.; Hunt, P. A.; Robb, M. A.; Takahashi, O. Non-adiabatic Direct Dynamics Study of Chromium Hexacarbonyl Photodissociation. *J. Phys. Chem. A.* **2002**, *106*, 10494-10504.
- (446) Blais, N. C.; Truhlar, D. G. Trajectory-surface-hopping Study of Na (3p 2P)+ H2 Na (3s 2S)+ H2 (v, J, Θ). *J. Chem. Phys.* **1983**, *79*, 1334-1342.
- (447) Tully, J. C. Molecular Dynamics with Electronic Transitions. *J. Phys. Chem.* **1990**, *93*, 1061-1071.
- (448) Hammes-Schiffer, S.; Tully, J. C. Proton Transfer in Solution: Molecular Dynamics with Quantum Transitions. *J. Phys. Chem.* **1994**, *101*, 4657-4667.
- (449) Granucci, G.; Persico, M. Critical Appraisal of the Fewest Switches Algorithm for Surface Hopping. *J. Phys. Chem.* **2007**, *126*, 134114.
- (450) Nakamura H. *Nonadiabatic Transitions, Concepts, Basic Theories and Applications World Scientific: Singapore, 2002.*
- (451) Zhu, C.; Kamisaka, H.; Nakamura, H. New Implementation of the Trajectory Surface Hopping Method with Use of the Zhu--Nakamura Theory. II. Application to the Charge Transfer Processes in the 3D DH2+ System. *J. Phys. Chem.* **2002**, *116*, 3234-3247.
- (452) Zhu, C.; Kamisaka, H.; Nakamura, H. Significant Improvement of the Trajectory Surface Hopping Method by the Zhu--Nakamura Theory. *J. Chem. Phys.* **2001**, *115*, 11036-11039.
- (453) Oloyede, P.; Mil'nikov, G.; Nakamura, H. Generalized Trajectory Surface Hopping Method Based on the Zhu-Nakamura Theory. *J. Chem. Phys.* **2006**, *124*, 144110.
- (454) Makri, N.; Miller, W. H. A Semiclassical Tunneling Model for Use in Classical Trajectory Simulations. *J. Chem. Phys.* **1989**, *91*, 4026-4036.
- (455) Czakó, G.; Kaledin, A. L.; Bowman, J. M. A Practical Method to Avoid Zero-point Leak in Molecular Dynamics Calculations: Application to the Water Dimer. *J. Phys. Chem.* **2010**, *132*, 164103.
- (456) Granucci, G.; Persico, M.; Zocante, A. Including Quantum Decoherence in Surface Hopping. *J. Phys. Chem.* **2010**, *133*, 134111.
- (457) Zheng, J.; Meana-Pañeda, R.; Truhlar, D. G. Including Tunneling in Non-Born-Oppenheimer Simulations. *J. Phys. Chem. Lett.* **2014**, *5*, 2039-2043.
- (458) Xu, X.; Zheng, J.; Yang, K. R.; Truhlar, D. G. Photodissociation Dynamics of Phenol: Multistate Trajectory Simulations Including Tunneling. *J. Am. Chem. Soc.* **2014**, *136*, 16378-16386.
- (459) Burghardt, I.; Meyer, H. -D.; Cederbaum, L. S. Approaches to the Approximate Treatment of Complex Molecular Systems by the Multiconfiguration Time-dependent Hartree Method. *J. Phys. Chem.* **1999**, *111*, 2927-2939.
- (460) Worth, G. A.; Burghardt, I. Full Quantum Mechanical Molecular Dynamics Using Gaussian Wavepackets. *Chem. Phys. Lett.* **2003**, *368*, 502-508.
- (461) Worth, G. A.; Robb, M. A.; Lasorne, B. Solving the Time-dependent Schrödinger Equation for Nuclear Motion in One Step: Direct Dynamics of Non-adiabatic Systems. *Mol. Phys.* **2008**, *106*, 2077-2091.
- (462) Wang, H.; Thoss, M. Multilayer Formulation of the Multiconfiguration Time-dependent Hartree Theory. *J. Chem. Phys.* **2003**, *119*, 1289-1299.
- (463) Schulze, J.; Shibl, M. F.; Al-Marri, M. J.; Kühn, O. Multi-layer Multi-configuration Time-dependent Hartree (ML-MCTDH) Approach to the Correlated Exciton-vibrational Dynamics in the FMO Complex. *J. Phys. Chem.* **2016**, *144*, 185101.
- (464) Richings, G. W.; Polyak, I.; Spinlove, K. E.; Worth, G. A.; Burghardt, I.; Lasorne, B. Quantum Dynamics Simulations Using Gaussian Wavepackets: The VMCG Method. *Int. Rev. Phys. Chem.* **2015**, *34*, 269-308.
- (465) Ben-Nun, M.; Martínez, T. J. Nonadiabatic Molecular Dynamics: Validation of the Multiple Spawning Method for a Multidimensional Problem. *J. Chem. Phys.* **1998**, *108*, 7244-7257.
- (466) Ben-Nun, M.; Quenneville, J.; Martínez, T. J. Ab Initio Multiple Spawning: Photochemistry From First Principles Quantum Molecular Dynamics. *J. Phys. Chem. A.* **2000**, *104*, 5161-5175.
- (467) Curchod, B. F.; Sisto, A.; Glowacki, D. R.; Martínez, T. J. *APS Meeting Abstracts* **2015**, *1*, 26008.
- (468) Curchod, B. F.; Rauer, C.; Marquetand, P.; González, L.; Martínez, T. J. Communication: GAIMS--Generalized Ab Initio Multiple Spawning for Both Internal Conversion and Intersystem Crossing Processes. *J. Phys. Chem.* **2016**, *144*, 101102.

