Sunlight Decatungstate Photoinduced Trifluoromethylations of (Hetero)aromatics and Electron-poor Olefins

Received 00th January 20xx,
Accepted 00th January 20xx

Sara Corsico, Maurizio Fagnoni, Davide Ravelli*

DOI: 10.1039/x0xx00000x

www.rsc.org/

Tetrabutylammonium decatungstate has been exploited to photoinduce the trifluoromethylation of (hetero)aromatics and electron-poor olefins. The process made use of the cheap Langlois' reagent ($\text{CF}_3\text{SO}_2\text{Na}$) and occurred under sunlight irradiation in an acetonitrile/water mixture. The trifluoromethylation of (hetero)aromatics required the use of potassium persulfate as an additive and the mono- or bis-trifluoromethylated adducts were obtained, depending on the actual reaction conditions and the chosen substrate.

Trifluoromethylation reactions are highly important since the incorporation of a CF_3 moiety into a pharmacophore is often known to increase lipophilicity/bioavailability, resistance to oxidation, and activity.¹ Accordingly, trifluoromethylated compounds are playing an increasingly important role in different fields, particularly in medicinal chemistry, but also in materials science. Chart 1 gathers some examples of drugs containing the trifluoromethylaryl core, including androgenic receptors (flutamide, nilutamide), antidepressants (fluoxetine) and antiarthritics (leflunomide, celcoxib). Accordingly, the introduction of a trifluoromethyl moiety in an organic compound is becoming a key issue in synthesis and different approaches have been recently developed to fulfill this goal.² Several articles have been reported in recent years on the use of photoredox catalysis for the generation of the trifluoromethyl radical and the ensuing functionalization of organic compounds.^{2,3} The strategy used is mainly that depicted in Scheme 1. The commonly used sources of $\text{F}_3\text{C}^\bullet$ are shown in Scheme 1a, including expensive Umemoto and Togni's reagents, along with cheaper CF_3I and $\text{CF}_3\text{SO}_2\text{Cl}$. The reductive cleavage of these reagents gave the $\text{F}_3\text{C}^\bullet$ radical that was used to functionalize alkenes (electron-deficient,⁴ electron-rich⁵ or styrene⁶ derivatives), alkynes⁷ and (hetero)aromatics⁸ (Scheme 1b). In the case of alkenes and alkynes, the adduct radical (**A**) was in most cases oxidized by the oxidized form of the photocatalyst (PC_{ox}) and a cationic

chemistry resulted. A radical homolytic aromatic substitution reaction (HAS) took place instead when using aromatic compounds (Scheme 1b).

Chart 1

In search for a cheaper and easier to handle trifluoromethylating agent, sodium triflinate ($\text{CF}_3\text{SO}_2\text{Na}$, the Langlois' reagent)⁹ was recently adopted.^{4b,8a,10} The triflinate reagent has the unique feature to be oxidized rather than reduced from the excited photocatalyst (Scheme 1c). This calls for photocatalysts having a reduction potential in the excited state E^*_{RED} superior to the oxidation potential of the triflinate anion ($E_{\text{OX}} = +1.05$ V vs SCE for potassium triflinate).¹¹ Albeit various photo(organo)catalysts were actually used for this purpose,^{4b,8a,10} a robust, cheap and efficient alternative is desirable. We envisaged that tetrabutylammonium decatungstate (TBADT)¹²⁻¹⁴ could be applied in this case as photoredox catalyst for the generation of the trifluoromethyl radical (Scheme 1c). In fact, TBADT was only sparsely used so far in the role of oxidant in the excited state^{13,14} despite its E^*_{RED} is quite high (a value of +2.44 V vs SCE was estimated)¹⁵ and largely sufficient to promote the reaction even under sunlight irradiation.^{12a}

^a PhotoGreen Lab, Department of Chemistry, Viale Taramelli 12, 27100 Pavia, Italy
E-mail: davide.ravelli@unipv.it

Electronic Supplementary Information (ESI) available: experimental details and copy of ^1H and ^{13}C NMR spectra of synthesized compounds. See DOI: 10.1039/x0xx00000x

COMMUNICATION

Since the temperature of the solution during irradiation was around 48 °C, we investigated the occurrence of the reaction also under thermal conditions. Thus, when the reaction was carried out on a heating plate at 50 °C, we observed a **2a** consumption of 65%, with **4** again formed in around 50% yield based on the consumption of the aromatic derivative (entry 11). We then surmised that, upon proper choice of the reaction conditions, we could drive the process towards the formation of **5**. Thus, when adopting a 3-fold amount of **1** in the presence of 2 equiv. of K₂S₂O₈, the reaction gave a mixture of **4** and **5** after irradiation for 20 h (entry 12). However, a clean reaction occurred when the excess of **1** was increased to 6 equiv., since **5** was formed in 82% GC yield (entry 13). When the reaction was repeated under the same conditions, but in the absence of TBADT, a complete consumption of **2a** was again observed, albeit a lower yield of **5** was obtained (56%, entry 14).

Thus, 1,3,5-trimethoxybenzene **2a** led to products **4** and **5** in 54 and 79% isolated yield, when subjected to the reaction conditions reported in entries 3 and 13, Table 1, respectively (Table 2). We then evaluated the scope of this trifluoromethylation procedure by subjecting different (hetero)aromatics to the optimized conditions reported in Table 1, entry 13, with the aim of maximizing the number of trifluoromethyl groups introduced in the final product.

Table 2. Trifluoromethylation of (hetero)aromatics.^a

^a Isolated yields after silica gel chromatography have been reported. ^b Reaction conditions: see entry 3, Table 1.

The trifluoromethylation of 1,4-dimethoxybenzene **2b** likewise proceeded smoothly to give product **6** in 45% yield. The presence of benzylic C-H bonds did not interfere with the desired reaction, as demonstrated by the formation of products **7** and **8** (both in 41% yield), from 3,4,5-trimethoxytoluene **2c** and mesitylene **2d**, respectively.

The reaction was next extended to heteroaromatic nuclei. The reaction of 2,6-dimethoxypyridine **2e** and 3-methylindole **2f** led to products **9** and **10** in ca. 50% isolated yield. Finally, the present protocol could be extended also to biologically relevant derivatives caffeine **2g** and theophylline **2h**, that were trifluoromethylated in the 8-position to give **11** and **12** in good yields (Table 2).

TBADT is widely used to perform radical conjugate addition reactions^{12a} and we then evaluated whether electron-poor olefins could be trifluoromethylated as well. Optimization of the reaction conditions made use of *N*-phenyl maleimide **3a** (0.1 M), that was subjected to trifluoromethylation conditions in the presence of different amounts of **1** and TBADT (4 mol%) in a MeCN:H₂O 5/1 mixture (Table 3). Thus, the performance of the process increased when passing from 1.5 to 3 equiv. of **1**, albeit a further increase (5 equiv.) did not improve the reaction yield (entries 1-3).

Table 3. Optimization of reaction conditions for the trifluoromethylation of **5a**.^a

Reaction scheme for Table 3: $\text{CF}_3\text{SO}_2^- \text{Na}^+$ (**1**) + **3a** (0.1 M) $\xrightarrow[\text{MeCN-H}_2\text{O 5/1, Additives}]{\text{TBADT (4 mol\%), hv (SolarBox)}}$ **13**

Entry	1 , M	Additives, M	Irradiation Conditions	3a Consumption, %	13 Yield, % ^b
1	0.15	-	SolarBox, 20h	60	37
2	0.3	-	SolarBox, 20h	100	42
3	0.5	-	SolarBox, 20h	100	44
4	0.3	Biphenyl, 0.1	SolarBox, 20h	80	45
5	0.3	K ₂ S ₂ O ₈ , 0.15	SolarBox, 20h	95	22
6 ^c	0.3	-	SolarBox, 20h	-	-
7 ^d	0.3	-	SolarBox, 20h	9	-
8	0.3	-	Heating, 20h (50°C)	< 5	-

^a Reaction conditions: **1** (0.15-0.5 M, 0.75-2.5 mmol), **3a** (0.5 mmol), (*n*-Bu₄N)₄[W₁₀O₃₂] (TBADT, 4·10⁻³ M, 4 mol%) in 5 mL of a MeCN-H₂O 5/1 mixture. ^b Gas Chromatography (GC) yields referred to the consumption of the limiting reagent (**3a**), using biphenyl as standard (*n*-undecane for entry 4). ^c In the absence of light, T = 35 °C. ^d No TBADT used, T = 48 °C.

Table 4. Trifluoromethylation of electron-poor olefins.^a

^a Isolated yields after silica gel chromatography have been reported. ^b Yield based on 44% consumption of **3d**.

The role of additives was next tested and neither biphenyl (1 equiv.) nor potassium persulfate (1.5 equiv.) improved the performance of the reaction (entries 4, 5). Blank experiments finally demonstrated that the present trifluoromethylation requires both light and TBADT to occur (entries 6, 7). Contrary to the trifluoromethylation of aromatics, in this case thermal conditions were not effective in promoting the formation of product **13** (entry 8).

Thus, under the optimized conditions reported in entry 2, Table 3, the reaction of **3a** allowed to isolate the expected product *N*-phenyl 2-(trifluoromethyl)succinimide **13** in 40% yield after purification with column chromatography (Table 4). The process was next extended to *N*-methyl maleimide **3b**, leading to the formation of **14** in 43% isolated yield. Finally, the reaction was tested on acyclic substrates, such as phenyl vinyl sulfone **3c** and ethyl isopropylidene cyanoacetate **3d**. The former gave 3,3,3-trifluoropropyl phenyl sulfone **15** in 43% isolated yield, while the latter allowed to prepare product **16** containing a quaternary carbon, albeit only a partial consumption of the olefin (44%) was observed. Thus, **16** was obtained in 23% overall yield (52% based on the consumption of **3d**).

As depicted in Scheme 2, the strong oxidizing properties of the excited decatungstate anion are the key to the success of this reaction. Thus, upon excitation with solar light, $[\text{W}_{10}\text{O}_{32}]^{4-*}$ is capable of oxidizing the sulfinate anion, that then loses SO_2 to liberate the trifluoromethyl radical. This is a highly reactive intermediate, prone to add to the reaction partners present in solution, either an aromatic or an electron-poor olefin. As for aromatic derivatives, radical adducts **I** are formed, while the reduced form of the photocatalyst $[\text{W}_{10}\text{O}_{32}]^{5-}$ is restored to its original form by the persulfate anion through a single electron-transfer step.¹⁶ This leads to the formation of sulfate radical $\text{SO}_4^{\cdot-}$ that is then involved in the oxidation of **I**, finally leading to the desired trifluoromethylated product. Likewise, bis-trifluoromethylated derivatives are formed in a similar way, when the original products undergo the same reaction sequence.

Scheme 2. Proposed reaction mechanism for the TBADT photoinduced trifluoromethylation of aromatics (dashed lines) and electron-poor olefins (solid lines); the steps in common to both mechanisms have been highlighted with a bold line.

Based on experimental evidences, we propose that the background reaction observed in the absence of the photocatalyst is due to thermal effects and is not related to direct absorption of light by the reagents (see Figure S1 in Electronic Supplementary Material with relevant UV-Vis spectra). This is also supported by the thermal experiments reported in Table 1 as well as by previous reports in the literature,¹⁷ in nice agreement with the temperature registered inside the SolarBox during irradiation. Accordingly, the role of the photocatalyst here is to accelerate the reaction,¹⁸ since only a partial consumption of substrate **2a** has been observed under thermal conditions (see also the kinetic profiles reported in Figure S2). At any rate, also under prolonged irradiation conditions, the presence of the photocatalyst allows to obtain a higher yield (compare the results of entries 13 and 14, Table 1). Another point is related to the formation of products either mono- (**4**, **6**, **8**, **10-12**) or bis-trifluoromethylated (**5**, **7**, **9**). In the case of **2a**, proper conditions were found to drive the process towards the clean formation of either of the trifluoromethylated derivatives. As a general trend, however, it is apparent that only markedly electron-rich nuclei (e.g. **2c** and **2e**) undergo a double trifluoromethylation process, while for the other substrates the clean introduction of a single CF_3 -group occurred.

Turning to electron-poor olefins, a different reaction course is involved, since this is a redox-neutral process and does not require the use of any external oxidant. Thus, the photogenerated $\text{F}_3\text{C}^\bullet$ intermediate undergoes addition onto electron-poor olefins **3** to give radical adducts **II**, that finally lead to the desired products **13-16** upon back-electron transfer from the reduced photocatalyst and protonation.

In conclusion, we demonstrated here a new application of TBADT as photoredox catalyst in promoting trifluoromethylation reactions of (hetero)aromatics and electron-poor olefins under sunlight irradiation. The process is

simple, versatile and occurs in a partial aqueous mixture, making use of the cheap Langlois' reagent as CF₃-source.

Conflicts of interest

There are no conflicts of interest to declare.

Acknowledgements

We thank Luca Capaldo and Edoardo Torti (University of Pavia) for preliminary experiments. D.R. thanks the MIUR for financial support (SIR Project "Organic Synthesis via Visible Light Photocatalytic Hydrogen Transfer"; Code: RBSI145Y9R).

- 1 *Fluorine in Pharmaceutical and Medicinal Chemistry. From Biophysical Aspects to Clinical Applications*; V. Gouverneur and K. Müller, Eds.; ICP: Oxford, 2012.
- 2 (a) J.-A. Ma and D. Cahard, Strategies for nucleophilic, electrophilic, and radical trifluoromethylations, *J. Fluorine Chem.*, 2007, **128**, 975–996; (b) N. Shibata, A. Matsnev and D. Cahard, Shelf-stable electrophilic trifluoromethylating reagents: A brief historical perspective, *Beilstein J. Org. Chem.*, 2010, **6**, No. 65; (c) Y. Ji, T. Brueckl, R. D. Baxter, Y. Fujiwara, I. B. Seiple, S. Su, D. G. Blackmond and P. S. Baran, Innate C-H trifluoromethylation of heterocycles, *Proc. Natl. Acad. Sci. USA*, 2011, **108**, 14411–14415; (d) A. Studer, A "Renaissance" in Radical Trifluoromethylation, *Angew. Chem. Int. Ed.*, 2012, **51**, 8950–8958; (e) S. Barata-Vallejo, S. M. Bonesi and A. Postigo, Photocatalytic fluoroalkylation reactions of organic compounds, *Org. Biomol. Chem.*, 2015, **13**, 11153–11183; (f) S. Barata-Vallejo, S. M. Bonesi and A. Postigo, Perfluoroalkylation reactions of (hetero)arenes, *RSC Adv.*, 2015, **5**, 62498–62518; (g) C. Alonso, E. Martínez de Marigorta, G. Rubiales and F. Palacios, Carbon Trifluoromethylation Reactions of Hydrocarbon Derivatives and Heteroarenes, *Chem. Rev.*, 2015, **115**, 1847–1935; (h) T. Chatterjee, N. Iqbal, Y. You and E. J. Cho, Controlled Fluoroalkylation Reactions by Visible-Light Photoredox Catalysis, *Acc. Chem. Res.*, 2016, **49**, 2284–2294; (i) T. Koike and M. Akita, Fine Design of Photoredox Systems for Catalytic Fluoromethylation of Carbon-Carbon Multiple Bonds, *Acc. Chem. Res.*, 2016, **49**, 1937–1945; (j) X. Pan, H. Xia and J. Wu, Recent advances in photoinduced trifluoromethylation and difluoroalkylation, *Org. Chem. Front.*, 2016, **3**, 1163–1185.
- 3 D. Ravelli, S. Protti and M. Fagnoni, Carbon-Carbon Bond Forming Reactions via Photogenerated Intermediates, *Chem. Rev.*, 2016, **116**, 9850–9913.
- 4 (a) X.-J. Tang and W. R. Dolbier, Jr., Efficient Cu-catalyzed Atom Transfer Radical Addition Reactions of Fluoroalkylsulfonyl Chlorides with Electron-deficient Alkenes Induced by Visible Light, *Angew. Chem. Int. Ed.*, 2015, **54**, 4246–4249; (b) Q. Lefebvre, N. Hoffmann and M. Rueping, Photoorganocatalysed and visible light photoredox catalysed trifluoromethylation of olefins and (hetero)aromatics in batch and continuous flow, *Chem. Commun.*, 2016, **52**, 2493–2496.
- 5 (a) W. J. Choi, S. Choi, K. Ohkubo, S. Fukuzumi, E. J. Cho and Y. You, Mechanisms and applications of cyclometalated Pt(II) complexes in photoredox catalytic trifluoromethylation, *Chem. Sci.*, 2015, **6**, 1454–1464; (b) D. Cantillo, O. de Frutos, J. A. Rincón, C. Mateos and C. O. Kappe, Continuous Flow α -Trifluoromethylation of Ketones by Metal-Free Visible Light Photoredox Catalysis, *Org. Lett.*, 2014, **16**, 896–899.
- 6 (a) Y. Yasu, T. Koike, and M. Akita, Three-component Oxytrifluoromethylation of Alkenes: Highly Efficient and Regioselective Difunctionalization of C=C Bonds Mediated by Photoredox Catalysts, *Angew. Chem. Int. Ed.*, 2012, **51**, 9567–9571; (b) G. Dagousset, A. Carboni, E. Magnier and G. Masson, Photoredox-Induced Three-Component Azido- and Aminotrifluoromethylation of Alkenes, *Org. Lett.*, 2014, **16**, 4340–4343; (c) Q.-Y. Lin, X.-H. Xu and F.-L. Qing, Chemo-, Regio-, and Stereoselective Trifluoromethylation of Styrenes via Visible Light-Driven Single-Electron Transfer (SET) and Triplet-Triplet Energy Transfer (TTET) Processes, *J. Org. Chem.*, 2014, **79**, 10434–10446.
- 7 (a) N. Iqbal, J. Jung, S. Park and E. Jin Cho, Controlled Trifluoromethylation Reactions of Alkynes through Visible-Light Photoredox Catalysis, *Angew. Chem. Int. Ed.*, 2014, **53**, 539–542; (b) R. Tomita, T. Koike and M. Akita, Photoredox-Catalyzed Stereoselective Conversion of Alkynes into Tetrasubstituted Trifluoromethylated Alkenes, *Angew. Chem. Int. Ed.*, 2015, **54**, 12923–12927.
- 8 (a) L. Li, X. Mu, W. Liu, Y. Wang, Z. Mi, and C.-J. Li, Simple and Clean Photoinduced Aromatic Trifluoromethylation Reaction, *J. Am. Chem. Soc.*, 2016, **138**, 5809–5812; (b) D. A. Nagib, D. W. C. MacMillan, Trifluoromethylation of arenes and heteroarenes by means of photoredox catalysis, *Nature*, 2011, **480**, 224–228; (c) N. J. W. Straathof, H. P. L. Gemoets, X. Wang, J. C. Schouten, V. Hessel and T. Noël, *ChemSusChem*, Rapid Trifluoromethylation and Perfluoroalkylation of Five-Membered Heterocycles by Photoredox Catalysis in Continuous Flow, 2014, **7**, 1612–1617; (d) J. Xie, X. Yuan, A. Abdukader, C. Zhu and J. Ma, Visible-Light-Promoted Radical C–H Trifluoromethylation of Free Anilines, *Org. Lett.*, 2014, **16**, 1768–1771.
- 9 (a) B. R. Langlois, E. Laurent and N. Roidot, "Pseudo-cationic" trifluoromethylation of enol esters with sodium trifluoromethanesulfinate, *Tetrahedron Lett.*, 1992, **33**, 1291–1294; (b) B. R. Langlois, E. Laurent and N. Roidot, Trifluoromethylation of aromatic compounds with sodium trifluoromethanesulfinate under oxidative conditions, *Tetrahedron Lett.*, 1991, **32**, 7525–7528.
- 10 (a) D. J. Wilger, N. J. Gesmundo and D. A. Nicewicz, Catalytic hydrotrifluoromethylation of styrenes and unactivated aliphatic alkenes via an organic photoredox system, *Chem. Sci.*, 2013, **4**, 3160–3165; (b) L. Cui, Y. Matusaki, N. Tada, T. Miura, B. Uno and A. Itoh, Metal-Free Direct C–H Perfluoroalkylation of Arenes and Heteroarenes Using a Photoredox Organocatalyst, *Adv. Synth. Catal.*, 2013, **355**, 2203–2207; (c) B. Li, D. Fan, C. Yang and W. Xia, UV light-mediated difunctionalization of alkenes with CF₃SO₂Na: synthesis of trifluoromethyl phenanthrene and anthrone derivatives, *Org. Biomol. Chem.*, 2016, **14**, 5293–5297; (d) L. Zhu, L.-S. Wang, B. Li, B. Fu, C.-P. Zhang and W. Li, Operationally simple hydrotrifluoromethylation of alkenes with sodium trifluoromethanesulfinate enabled by Ir photoredox catalysis, *Chem. Commun.* 2016, **52**, 6371–6374; (e) X.-L. Yu, J.-R. Chen, D.-Z. Chen and W.-J. Xiao, Visible-light-induced photocatalytic azotrifluoromethylation of alkenes with aryldiazonium salts and sodium trifluoromethanesulfinate, *Chem. Commun.*, 2016, **52**, 8275–8278; (f) H.-T. Qin, S.-W. Wu, J.-L. Liu and F. Liu, Photoredox-catalysed redox-neutral trifluoromethylation of vinyl azides for the synthesis of α -trifluoromethylated ketones, *Chem. Commun.*, 2017, **53**, 1696–1699; (g) Q. Lefebvre, Toward Sustainable Trifluoromethylation Reactions: Sodium Trifluoromethanesulfinate under the Spotlight, *Synlett*, 2017, **28**, 19–23.
- 11 J.-B. Tommasino, A. Brondex, M. Médebielle, M. Thomalla, B. R. Langlois and T. Billard, Trifluoromethylation Reactions with Potassium Trifluoromethanesulfinate under Electrochemical Oxidation, *Synlett*, 2002, 1697–1699.
- 12 (a) D. Ravelli, S. Protti and M. Fagnoni, Decatungstate Anion for Photocatalyzed "Window Ledge" Reactions, *Acc. Chem.*

- Res., 2016, **49**, 2232–2242 and references therein; (b) I. N. Lykakis, E. Evgenidou and M. Orfanopoulos, Photo-catalysis and Polyoxo-anion Decatungstate in Organic Chemistry: A Manifold Concept for Green Chemistry, *Curr. Org. Chem.*, 2012, **16**, 2400–2414; (c) M. D. Tzirakis, I. N. Lykakis and M. Orfanopoulos, Decatungstate as an efficient photocatalyst in organic chemistry, *Chem. Soc. Rev.*, 2009, **38**, 2609–2621; (d) C. L. Hill, Introduction of Functionality into Unactivated Carbon-Hydrogen Bonds. Catalytic Generation and Nonconventional Utilization of Organic Radicals, *Synlett*, 1995, 127–132.
- 13 (a) T. S. Symeonidis, I. Tamiolakis, G. S. Armatas and I. N. Lykakis, Green photocatalytic organic transformations by polyoxometalates vs. mesoporous TiO₂ nanoparticles: selective aerobic oxidation of alcohols, *Photochem. Photobiol. Sci.*, 2015, **14**, 563–568; (b) I. Texier, J. A. Delaire and C. Giannotti, Reactivity of the charge transfer excited state of sodium decatungstate at the nanosecond time scale, *Phys. Chem. Chem. Phys.*, 2000, **2**, 1205–1212.
- 14 (a) L. Capaldo, L. Buzzetti, D. Merli, M. Fagnoni and D. Ravelli, Smooth Photocatalyzed Benzylolation of Electrophilic Olefins via Decarboxylation of Arylacetic Acids, *J. Org. Chem.*, 2016, **81**, 7102–7109; (b) S. Montanaro, D. Ravelli, D. Merli, M. Fagnoni and A. Albini, Decatungstate As Photoredox Catalyst: Benzylolation of Electron-Poor Olefins, *Org. Lett.*, 2012, **14**, 4218–4221.
- 15 V. De Waele, O. Poizat, M. Fagnoni, A. Bagno and D. Ravelli, Unraveling the Key Features of the Reactive State of Decatungstate Anion in Hydrogen Atom Transfer (HAT) Photocatalysis, *ACS Catal.*, 2016, **6**, 7174–7182.
- 16 M. C. Quattrini, S. Fujii, K. Yamada, T. Fukuyama, D. Ravelli, M. Fagnoni and I. Ryu, Versatile cross-dehydrogenative coupling of heteroaromatics and hydrogen donors via decatungstate photocatalysis, *Chem. Commun.*, 2017, **53**, 2335–2338.
- 17 (a) W. Wei, J. Wen, D. Yang, X. Liu, M. Guo, R. Dong and H. Wang, Metal-Free Direct Trifluoromethylation of Activated Alkenes with Langlois' Reagent Leading to CF₃-Containing Oxindoles, *J. Org. Chem.*, 2014, **79**, 4225–4230; (b) D. Wang, G.-J. Deng, S. Chen and H. Gong, Catalyst-free direct C–H trifluoromethylation of arenes in water–acetonitrile, *Green Chem.*, 2016, **18**, 5967–5970.
- 18 I. Ryu, A. Tani, T. Fukuyama, D. Ravelli, S. Montanaro and M. Fagnoni, Efficient C–H/C–N and C–H/C–CO–N Conversion via Decatungstate-Photoinduced Alkylation of Diisopropyl Azodicarboxylate, *Org. Lett.*, 2013, **15**, 2554–2557.