Revisiting *Ursus etruscus* (Carnivora, Mammalia) from the Early Pleistocene of Greece
with description of new material

George D. Koufos ^{a,*}, George E. Konidaris ^b, Katerina Harvati ^b

^a Aristotle University of Thessaloniki, Department of Geology, Laboratory of Geology and Palaeontology, 54124 Thessaloniki, Greece.

^b Eberhard-Karls University of Tübingen, Palaeoanthropology, Senckenberg Centre for Human Evolution and Palaeoenvironment, Rümelinstr. 23, 72070 Tübingen, Germany.

* Corresponding author. E-mail address: koufos@geo.auth.gr (G.D. Koufos).

Abstract

The Villafranchian ursids of Greece are scanty, represented by a few isolated teeth and bones. During our last field campaigns in the Mygdonia Basin (Macedonia, Greece) we discovered an almost complete cranium, as well as some cranial, dental and postcranial remains, which are presented in this article. The new material originates from the locality Tsiotra Vryssi (TSR), dated to the late Villafranchian (1.8–1.2 Ma). The specimens are described and compared with ursids from various European localities; their morphological characters and dimensions place it with *Ursus etruscus* Cuvier, 1823, while the dental features are most similar to the specimens from Pietrafitta (Italy). The TSR cranium was CT-scanned in order to create a 3D virtual model and study its internal anatomy. Its observed endocranial traits exhibit primitive conditions, confirming the basal position of *U. etruscus* among the Pleistocene European ursids. Most of these features are shared with *U. arctos* and clearly differ from *U. deningeri* and *U. spelaeus*. Finally, we provide a revision of the known Greek material of *U. etruscus* and the biostratigraphic distribution of the Miocene–Pleistocene ursids of Greece. The earliest secure appearance of *Ursus etruscus* in Greece is found in the locality Dafnero (Western Macedonia), dated to the end of the middle Villafranchian at ~2.0 Ma, whereas its probable last occurrence is known from the locality Apollonia 1 of the Mygdonia Basin, dated to the latest Villafranchian at ~1.2 Ma.

Keywords: Greece, Villafranchian, *Ursus*, systematic, endocranium, biostratigraphy

1. Introduction

Although the Neogene/Quaternary fossil mammal archive of Greece is quite rich, the known ursids are relatively scarce. The oldest ursid remains from this country have been recovered from the late Miocene localities Pikermi, Perivolaki, Samos and Halmyropotamos, and are attributed to the genera *Ursavus* and *Indarctos* (Koufos, 2013). After a long temporal interval, ursids re-occurred in the late Pliocene of Greece. During the Early Pleistocene bears were represented by *Ursus etruscus*, while during the Middle and Late Pleistocene by the cave bears, continuing their presence until today with the brown bear *U. arctos* (Kostopoulos and Vassileiadou, 2006; Koufos and Kostopoulos, 2016 and cited references).

Ursus etruscus is a common Villafranchian ursid known from several Greek localities, but in most cases it is represented by fragmentary material, sometimes leading to insecure determinations. The new material, described in this article, is to date the richest known Greek sample. It comprises both cranial and postcranial remains, thus allowing confident attribution and confirmation of the presence of the species in the Early Pleistocene mammal record of Greece. It was recovered during excavation by a joint team from the Aristoteleion University of Thessaloniki and the University of Tübingen (field seasons 2014-2017) from the locality Tsiotra Vryssi (TSR) of Mygdonia Basin (Macedonia, Greece), about 90 km southeast of Thessaloniki (Fig. 1a; Konidaris et al., 2015). Stratigraphically, this fossiliferous site is located in the upper part of the Gerakarou Formation, and consists mainly of sands, gravels, sandy-clays and red clays (Koufos et al., 1995; Konidaris et al., 2015). The TSR fauna determined so far comprises several mammalian taxa (including the carnivores *Pachycrocuta brevirostris* and *Canis etruscus*), as well as some reptiles (Chelonii) and birds. The preliminary biochronological correlation of the mammal fauna suggests a late Villafranchian age, between 1.8 and 1.2 Ma. Further details about the stratigraphy, locality and fauna are reported in Konidaris et al. (2015, 2016).

2. Materials and methods

The ursid material from Tsiotra Vryssi studied in this article is stored at the Laboratory of Geology and Palaeontology, Aristotle University of Thessaloniki (LGPOT). Other bear specimens from Greek localities included here originate from Dafnero 1, Vassiloudi, Apollonia 1, Libakos (all at LGPUT) and Sesklon (Athens Museum of Palaeontology and Geology, AMPG). The uppercase letters C, P and M are used to indicate the upper canine, premolars and molars, and the lowercase letters c, p and m the corresponding lower teeth. The dental nomenclature used follows that of Rabeder (1999). The cranial, dental and postcranial measurements are taken according to Mazza and Rustioni (1992); the endocast and frontal sinuses measurements are taken according to García et al. (2007). The software PAST 3.1 (Hammer et al., 2001) has been used for the discriminant analysis in Fig. 4.

The cranium TSR-E21-50 was scanned using a v|tomex|l GE CT scanner at the “Forschungsinstitut Edelmetalle + Metallchemie” (Schwäbisch Gmünd, Germany). The cranium was subjected to high resolution CT scanning with the following parameters: scanner energy 260 Kv, 111 mA, slice thickness 0.15924875 mm, inter-slice spacing 0.15924875 mm, 2024 × 2024 pixel Detector and 2013 × 850 × 772 pixels matrix size. 772 slices were obtained, in .vol format with an image resolution of 0.15924875 mm. Digital cleaning was conducted to distinguish the bones and teeth from the matrix in each individual slice (segmentation), subsequently creating the cranium’s three dimensional computer model. The virtual endocast of the cranium, as well as the linear and volumetric measurements, were obtained using the Avizo Lite 9.0.1 software.

3. Systematic palaeontology

Order Carnivora Bowdich, 1821

Family Ursidae Fischer de Waldheim, 1817

Genus *Ursus* Linnaeus, 1758

Ursus etruscus Cuvier, 1823

Locality. Tsiotra Vryssi, Mygdonia Basin, Macedonia, Greece (Fig. 1a).

Age. Late Villafranchian; Early Pleistocene; 1.8–1.2 Ma (Konidaris et al., 2015).

Material. Cranium, TSR-E21-50; right maxillary fragment with P4–M2, TSR-E16-18; maxillary fragment with both I1 and the left canine, TSR-E21-63; left P3, TSR-E18-12; right m1, TSR-E19-34; atlas + axis, TSR-E21-52a, b; proximal part of right ulna, TSR-C16-26; calcaneum + MtI + MtII + MtIII + MtIV + MtV + (PhI + PhII first digit) + (PhI + PhII + PhIII second digit) + (PhI + PhII third digit) + PhI fourth digit + (PhI + PhII + PhIII fifth digit), TSR-160.

Description

Cranium TSR-E21-50 (Fig. 2a–e; virtual 3D model shown in Fig. 8a–c). It is relatively well preserved, lacking only the premaxillae with the incisors and the left canine, the left part of the nasal cavity and the distal half of the right zygomatic arch. It is dorso-laterally compressed and the right side of the face is slightly deformed.

The face and the neurocranium are elongated. Although prosthion is not preserved, the estimated length frontal midpoint–prosthion is slightly smaller than the neurocranial length (Table 1, meas. 5 and 7). The right nasal bone is partially preserved and appears elongated and narrow. The maxilla is flattened in front of the orbit and bears a slight bulging at the area of the canine's root. The infraorbital foramen is large and situated in front of the ventral border of the orbit with its distal margin above the metacone of the M1. The right

orbit is well-preserved and rounded with its mesial margin above the mesial border of the M2. The left zygomatic arch, although compressed and deformed, is complete, robust and slightly laterally projecting. Its root is placed above the M2. The forehead is broad and flat with a slight concavity in the middle. The temporal lines are weak and visible only in the distal part of the neurocranium, converging far back on the distal end of the parietals. The sagittal crest is very short, restricted to the area of the external occipital protuberance (~30 mm from its end) and is only slightly projecting. The occiput is well-preserved, high, and fairly inclined backwards. The nuchal crest is prominent and continues ventrally and then mesially to connect with the zygomatic arch. The occipital condyles are small and the foramen magnum large; the dimensions of the latter are 31.5×22.0 mm. The paraoccipital processes are short and relatively wide; their height slightly exceeds that of the occipital condyles. The auditory bullae are small, fairly concave with a large external acoustic meatus. The palate is elongated and relatively narrow; this narrowness is partially due to the lateral compression of the frontal part of the skull. The choanae are elongated, wide, elliptical and deep, with their mesial border far behind the distal border of the M2; the distance from the mesial border of choanae to the middle of the line connecting the distal end of the M2s is 20.0 mm.

The cranium preserves the right tooth row C–M2 which, however, lacks the P1 (its alveolus is well distinguished), and the left tooth row P3–M2 with the alveoli of the P1 and P2 (Fig. 2d–e; virtually reconstructed in Fig. 8h, i). All teeth are well-preserved and moderately worn. The canine is slender, sharp, flattened bucco-lingually and slightly curved backwards; it bears a disto-buccal and a mesio-lingual crest, both running from the base to the apex. The exposed part of the root indicates that it is large and elongated. With the exception of the P4, the premolars are small and bear a diastema between them, as well as between the P1 and the canine; the longest diastema is between the P2 and P3 and measures

5.72 mm. The P2 is monocuspid and single-rooted with elliptical occlusal outline and weak lingual cingulum. The morphologically similar but larger P3 is situated laterally to the P4 (its mesio-distal axis forms an angle of $\sim 40^\circ$ with that of the P4). The P4 shows a triangular occlusal outline with a large and high paracone, a metacone that is smaller and lower than the paracone and a small protocone, which is situated more distally. A weak vertically oriented metastyle is present on the distal end of the metacone. It has pronounced lingual and distal cingula. The M1 is rectangular in occlusal aspect with pronounced and equally-sized buccal cusps (paracone and metacone). The lingual cusps (protocone and hypocone) are also equal-sized but lower than the buccal ones. The M1 bears a small parastyle and metastyle, as well as a large and clear metaconule between the protocone and metacone. There is a weak lingual cingulum and a stronger buccal one. The talon is short and inclines abruptly backwards. In occlusal aspect, the M2 is elongated with a narrow talon; the talon is detached from the trigonid by an abrupt step and twists lingually in buccal aspect. The buccal cusps are lower than those of the M1, and both bear a distal crest-like projection; the distal projection of the metacone bears a cuspule. The lingual cusps are large and low; the hypocone extends distally and bears a second cuspule. The margin of the talon is elevated and separates a flat occlusal surface. There is a small metaconule, which is worn and connected to the protocone. The tooth has strong lingual cingulum in its mesial half but lacks buccal cingulum.

Maxilla TSR-E16-18 (Fig. 2h-i). This specimen is a right maxillary fragment, preserving part of the orbit, the tooth row P4–M2 and the P1–P3 alveoli. The maxillary bone is well-preserved with a strong bulging at the canine's root. The mesial margin of the orbit is situated above the mesial border of the M2. The large infraorbital foramen is situated in front of the ventral margin of the orbit and its distal margin lies above the metacone of the M1. The teeth are less worn than those of TSR-E21-50 but they have similar morphology. The less worn teeth allow the better distinction of some dental cusps, like the metaconule of the M1 and the

talon morphology of the M2. The latter is unworn and bears several ridges in its occlusal surface.

Maxillary fragment, TSR-E21-63 (Fig. 2f-g). This specimen preserves the most frontal part of the cranium with both first incisors and the left canine in place. It was found very close to the TSR-E21-50 cranium (Fig. 1b) and likely belongs to the same individual. The partially preserved nasal cavity is wide and abruptly inclined. The I1 is well preserved and moderately worn; in occlusal view the lingual and labial borders are convex and the mesial and distal ones almost straight; it bears a pronounced labial cusp and a series of cusplets across the lingual margin; a fissure separates them from the labial cusp. The canine has the same morphology and size to that to that of the cranium TSR-E21-50.

Left P3 TSR-E18-12 (Fig. 2j-l). The morphology closely resembles that of the P3 of the cranium TSR-E21-50. The preserved root is single and strong with rounded transverse section; its length is 13.5 mm.

Left m1 TSR-E19-34 (Fig. 2m). This isolated m1 preserves a little-worn crown with all its cuspids. It is elongated and slender with wider talonid than trigonid. The paraconid is large and well separated from the rest of the trigonid by a deep, wide and lingually open valley. The protoconid is the highest cuspid; the metaconid is small and low, situated slightly behind the protoconid; the metastylid is a small cuspid situated in front of the metaconid. The talonid is well-separated from the trigonid by a wide and low basin; the hypoconid and entoconid are of equal height but the former is larger; the entoconid is single with a weak notch in its mesial border, indicating the origin of a secondary one; the hypoconulid is small and forms the distal margin of the talonid; in the valley between the protoconid and hypoconid there are two small cusplets.

Postcranial elements. The atlas and axis, TSR-E21-52a, b (Fig. 3a-d) were found articulated with each other and close to the TSR-E21-50 cranium (Fig. 1b). They likely belong to the

same individual. The atlas bears two large and concave articular facets for the cranium and two other, flat facets, for the axis. The axis, which lacks its distal part, has a high and long spinal process and its odontoid process is cylindrical and large. The ulna TSR-C16-26 (Fig. 3e–f) preserves its proximal half, which lacks the lateral surface of the olecranon; the latter is relatively short; the beak is quadrangular. The calcaneus + metatarsus + phalanges, TSR-160 (Fig. 3g–j) were found articulated (Fig. 3i). The calcaneus is well-preserved having the general ursid morphology; it is short with large sustentaculum tali; the articular facet for the cuboid is perpendicular to the longitudinal axis of the calcaneus and its surface is concave; the calcaneus tuber is moderately wide and its proximal surface is flattened with a slight depression. In dorsal view the metatarsals are straight; in lateral view they are slightly convex; the condyle of the distal epiphysis is robust and bears a strong keel ventrally. The first phalanges are straight and flattened dorso-ventrally; the proximal articular surface is wide and strongly concave; the dorsal one has two condyles separated by a wide groove; in ventral view they bear two strong projections just above the distal epiphysis at the lateral borders of the diaphysis.

4. Results of the comparative analyses

The history of the species *Ursus etruscus* is long and rather complicated (see Mazza and Rustioni, 1992; Baryshnikov, 2007). Mazza and Rustioni (1992) based their amended diagnosis on the male cranium VA 1119 (Naturhistorisches Museum Basel, Switzerland) and the female cranium IGF 906 (Museum of Geology and Palaeontology of Florence, Italy) from Upper Valdarno (Italy). The convex dorsal profile of the cranium in the parietals, the elongated neurocranium, the small occipital condyles, the broad, flat and slightly convex forehead, the elongated face, the very small upper premolars separated by large diastemas, the rectangular M1 with weak parastyle and metastyle, and the elongated M2 are the main diagnostic features of *U. etruscus* (Mazza and Rustioni, 1992). All these characters are

present in the studied cranium TSR-E21-50. Additionally, the morphology of the TSR upper teeth and m1, as well as the morphology and size of the studied postcranial elements coincide with that of *U. etruscus* as described by Mazza and Rustioni (1992) and Petrucci et al. (2013), (Table 3).

The metric comparison of TSR-E21-50 with *U. etruscus* from different European localities is mainly based on the P4–M2 dimensions, as the available cranial dimensions are limited due to its deformation and the absence of some crucial parts (e.g., prosthion). However, the cranial measurements that can be securely assessed fall within the range of variability of *U. etruscus* from Upper Valdarno and Olivola (Italy) and are slightly smaller than those of the sample from Saint-Vallier (France) (Table 1). We compared the dental dimensions of the TSR sample with those of *U. etruscus* from different European localities by discriminant analysis and classification using the software PAST 3.10 (Hammer et al., 2001). The P4, M1 and M2 length and maximal breadth of only complete P4–M2 rows were included as variables in the analysis. *Ursus arctos*, *U. deningeri*, *U. spelaeus* and *U. rodei* were used as comparative material. Axis 1 clearly separates *U. etruscus* from all the other species. The TSR sample is very close to the convex hull of *U. etruscus* from Upper Valdarno / Olivola (group a) and to the Pirro Nord (Italy) sample (group d) (Fig. 4); the TSR ursid is classified as *U. etruscus* from Upper Valdarno / Olivola (overall classification success of 84.62%; see table of Fig. 4). Like the TSR sample, the Pietrafitta (Italy) specimen (group b) is also outside but close to the convex hull of *U. etruscus* from Upper Valdarno / Olivola, indicating metrical similarities between these samples. Further, the Saint-Vallier and Ceyssaguet (France) ursids (group c and e) differ slightly from the typical *U. etruscus* from Italy (Fig. 4). In order to check possible differences between each tooth, we used Simpson's logarithmic diagrams comparing the mean upper dental dimensions (Fig. 5). Compared to the Upper Valdarno sample, the TSR P3 is slightly longer, and the M1 and M2 are more

compressed laterally, like those of Olivola and Pietrafitta. The isolated m1 TSR-E19-34 falls within the range of size variation of *U. etruscus* (Fig. 6d).

Viret (1954) described a rich sample of *U. etruscus* from the locality Saint-Vallier, dated to the middle Villafranchian, at ~2.0 Ma (Guérin et al., 2004). Later, Mazza and Rustioni (1992) considered the Saint-Vallier ursid different from the typical *U. etruscus* as it possesses some distinct cranial and dental characters (e.g., more convex dorsal profile of the cranium, short muzzle, larger premolars with short diastemas between them, squarish (shorter mesiodistally) occlusal outline on M1 and short M2 (Fig. 5a). The same opinion was expressed by Baryshnikov (1992), who later described the Saint-Vallier ursid as a separate subspecies (Baryshnikov, 2007). Meanwhile, Argant (2004) considered that the small differences observed in the Saint-Vallier sample can be attributed to individual or geographic variability of *U. etruscus*. The TSR m1 is similar in size to the Upper Valdarno / Olivola *U. etruscus*, but it is longer than the Saint-Vallier specimen (Fig. 6d); this size difference is also noted between Saint-Vallier and the Upper Valdarno / Olivola sample (Mazza and Rustioni, 1992).

Ursus etruscus is also known from Pirro Nord (Italy). The Pirro Nord fauna is correlated to the late Villafranchian and its age is estimated from 1.5 to 1.2 Ma on the basis of the presence of the vole *Allophaiomys ruffoi* (Arzarello et al., 2016). The morphology of the TSR upper teeth fits well with that of the Pirro Nord Etruscan bear as it is described by Petrucci et al. (2013); the dental dimensions of these two samples are also similar (Fig. 5a). However, despite their similarities, the TSR sample differs from that of Pirro Nord by displaying a narrower M1 and M2, as well as a longer and narrower m1 (Figs 5a, 6d). The locality Pietrafitta (Italy) has also yielded ursid remains, which have been described as *U. etruscus* by Rustioni and Mazza (1993). They found that some upper carnassials (except specimen 7), the M2, 3, the p4 and m2, 3 are more laterally compressed, bear higher and

sharper cusps, and show more extensive development of crests and ridges than the teeth from Upper Valdarno. These authors concluded that the Pietrafitta ursid appears to be an advanced form of *U. etruscus*, and the last ascertained representative of this taxon in Europe (Rustioni and Mazza, 1993; Mazza and Rustioni, 1994). The TSR upper carnassial, the M1, M2 and m1 are narrower than those of Upper Valdarno and Pirro Nord (Figs 5a, 6) and thus close to the more advanced form of *U. etruscus* from Pietrafitta. *Ursus etruscus* is also known from the Iberian localities Venta Micena [ca. 1.6–1.5 Ma (Martínez-Navarro, 1991; Oms et al., 2000)], Barranco León [1.43 ± 0.38 Ma (Toro-Moyano et al., 2013)] and Fuente Nueva-3 [1.19 ± 0.21 Ma (Duval et al., 2012)]. The morphology and size of the upper dentition of *U. etruscus* from these localities (Medin et al., 2017) is similar to that of TSR.

The new species *U. dolinensis* has been described from the TD4 level of Trincheria Dolina, Atapuerca (Spain), originally dated between 0.9 and 0.78 Ma (García and Arsuaga, 2001). Recent palaeomagnetic studies suggest an older age for this level, at ~1.0 Ma (Parés et al., 2013). In addition to its larger dental size than TSR and other samples of *U. etruscus* (Fig. 5b), *U. dolinensis* is distinguished by the trapezoidal occlusal outline of the M1 with visible but not large styles oriented vertically, and by the larger M2 with weak internal structure (García and Arsuaga, 2001). The P4 of *U. dolinensis* is metrically within the range of variation of *U. etruscus*, but the great overlap with *U. deningeri* (Fig. 6b) prevents a clear distinction. Moreover, the fourth (length: 92.0 mm) and fifth (length: 89.0–98.5 mm) metatarsal from Atapuerca are significantly longer than those from TSR (Table 3). Another new species, named *U. rodei*, was described from the Epivillafranchian locality Untermassfeld (Germany) (Musil, 2001). This species shares several cranio-dental features with *U. dolinensis* and thus they are considered synonyms (García, 2004; Kahlke, 2006). Compared to the TSR ursid it has quite larger dental dimensions, being closer to *U. deningeri* in size (Fig. 5b). Rabeder (2010) assigned the ursids from Deutsch-Altenburg (Austria) to the

arctoid lineage, noting their larger size, wider P4 and longer metatarsals compared to *U. etruscus*. All these metric differences from the Deutsch-Altenburg ursid are also present in the TSR material, separating it from this taxon.

5. Sex, age and body mass determination of the TSR specimens

The available material of the TSR *ursus etruscus* suggests the presence of at least two individuals (MNI= 2).

Sex. Following the observations and criteria of Mazza and Rustioni (1992), the small overall cranial size, the widely spaced and weakly marked temporal lines, the short and slightly developed sagittal crest, and the rather weak zygomatic arch of the cranium TSR-E21-50 point to a female individual (Fig. 8a–c). Moreover, the dimensions of the upper canine, which are used to separate sexes in cave bears (e.g., Koby, 1949; Kurtén, 1955), plot clearly in the lower limit of the *U. etruscus* sample (Fig. 6a), also supporting a female designation.

Age. Applying the age-scoring technique of Stiner (1998) based on the wear stages of P4–M2, the slightly worn teeth of TSR-E21-50, with exposed dentine on the lingual cusps of the M1 and M2, as well as the incompletely fused cranial sutures, indicate that the animal was at the transition to adulthood (stage IV) and thus can be regarded as prime adult. The maxilla TSR-E16-18 belongs to a younger individual (stage III), as evidenced by the only slightly worn occlusal surface without dentine exposure, and is considered a juvenile. The slightly worn m1 TSR-E19-34 without exposed dentine (stage III) also belongs to a juvenile.

Body mass. In order to estimate body mass (BM), we applied the regression equations proposed by Van Valkenburgh (1990) for ursids. For the cranium TSR-E21-50 we used the occiput-to-orbit length (OOL = 186 mm) [$\log BM = 1.98 \log(OOL) - 2.38$] and we obtained a body mass of 129.9 kg. This value falls within the body mass variability observed in females of the European *U. arctos* (Swenson et al., 2007, tab. 4). For the m1 TSR-E19-34 we used its

total length (M1L = 25.6 mm) [$\log BM = 0.49 \log(M1L) + 1.26$] and we obtained a body mass of 89.1 kg. However, M1L is considered as less reliable body weight predictor compared to OOL (Van Valkenburgh, 1990).

6. Virtual model of the TSR-E21-50 *U. etruscus* cranium

In order to complete the exocranial study of the *U. etruscus* cranium TSR-E21-50 with its endocranial morphology, we performed a high resolution CT scan and created its 3D virtual model. The internal cranial anatomy of *U. etruscus* has to date not been described and therefore this study aims to add new information on this species and provide a more detailed comparison of *U. etruscus* with *U. dolinensis* (Trinchera Dolina; late Early Pleistocene), *Ursus* sp. (Deutsch-Altenburg 4; late Early Pleistocene), *U. deningeri* (Sima de los Huesos, Petralona; Middle Pleistocene), *U. spelaeus* (Spain; Late Pleistocene), *U. ingressus* (Drachenhöhle near Mixnitz, Austria; Late Pleistocene) and *U. arctos* (Holocene), whose endocranial morphology (either through CT scans or natural sections/casts of brains and frontal sinuses) has already been described (Drexler, 1931; Groiss, 1994; García et al., 2006, 2007; Rabeder et al., 2010; Santos et al., 2014). Although the TSR cranium was totally filled internally with sediment, it preserves most of its endocranial structures (Fig. 7). These are described and compared below. Measurements of the endocast and the frontal sinuses are reported in Table 4.

Forehead. In lateral view, the forehead of the TSR-E21-50 *U. etruscus* cranium is relatively straight (Fig. 8a). This is clearly different from the step-profile of *U. deningeri* and the even more stepped and vaulted forehead of *U. spelaeus* and *U. ingressus*, which show a deep glabella. The straight profile of the TSR specimen is possibly related also to its female sex, as male *U. etruscus* seem to have a more stepped forehead (e.g., Mazza and Rustioni, 1994: pl. 4, fig. 3a), although not as pronounced as that shown by cave bears. The forehead profile of

TSR-E21-50 is similar to that of *U. arctos* females (e.g., Mazza and Rustioni, 1994: pl. 6, fig. 3a)

Frontal sinuses. The preserved morphology of the TSR-E21-50 frontal sinuses shows that they are caudo-rostrally elongated, extending rostrally until the level of the distal half of the m1, while caudally they cover approximately the 1/3 to half of the brain's dorsal surface (Fig. 8e, f). The rostral frontal sinus is rather long and flat, while the caudal one is relatively thin and weak (Fig. 8j). This morphology is similar to that of the brown bear *U. arctos*. It differs from the condition present in the cave bears *U. deningeri*, *U. spelaeus* and *U. ingressus*, which show dorso-ventrally enlarged frontal sinuses with a rounded and robust rostral part, reaching the level of the m2, and an higher and stronger caudal part. Furthermore, in *U. spelaeus* and *U. ingressus* the sinuses are caudally extended, covering most of the dorsal brain's surface. The caudal part of the frontal sinuses of the Deutsch-Altenburg 4 ursid is thin and weak, thus resembling the morphology of *U. etruscus* and *U. arctos*.

Palate. In sagittal view, the palate of the TSR-E21-50 *U. etruscus* is thin and rather straight, as it is in *U. arctos* (Figs 7a, 8g, 9a). In *U. deningeri* the palate is thicker and dorsoventrally concave, while in *U. spelaeus* it is even thicker, but less concave. This feature differentiates also the TSR *U. etruscus* from *U. dolinensis*, which presents a thicker palate (Fig. 9a).

Basioccipital. Like *U. arctos* the TSR-E21-50 *U. etruscus* shows a thin and non-pneumatized basioccipital, in contrast to *U. spelaeus*, where this bone is thick and pneumatized (Figs 7d, 8d, 9b). *U. deningeri* also shows a rather thick, but non-pneumatized, basioccipital.

Antero-dorsal shape of the brain cavity. The antero-dorsal profile of the TSR-E21-50 brain cavity displays a convex shape (Figs 7a, b, 8g). This is most similar to the curved profile of *U. arctos*, whereas the profile of *U. deningeri*, *U. spelaeus* and *U. ingressus* is straight or only slightly curved.

Cribriform plate. The TSR-E21-50 cranium shows a large and hemispheric cribriform plate with a slight constriction separating it from the brain cavity (Fig. 7b). It thus most closely resembles the morphology of *U. arctos*. The cribriform plate is smaller and ellipsoid in *U. deningeri* and *U. spelaeus*, and a marked constriction is present, clearly dividing the anterior cranial fossa from the rest of the brain cavity.

Virtual endocast. In lateral view (Fig. 8k), the shape of the TSR-E21-50 endocast is more rounded compared to the caudo-rostrally elongated endocast of *U. deningeri* and *U. spelaeus*, and thus closer to the shape of *U. arctos* (García et al., 2007: fig. 8). The antero-dorsal profile of the TSR-E21-50 endocast shows only a slight curvature, similar to *Ursus* sp. from Deutsch-Altenburg 4, whereas in *U. arctos* this profile is more convex; in *U. deningeri* and *U. spelaeus* the profile is straight (García et al., 2007; Rabeder et al., 2010). The most rostral part of the brain cavity is not well preserved and therefore the olfactory bulbs could not be reconstructed.

García et al. (2007) summarized the primitive and derived endocranial traits of the Middle and Late Pleistocene European ursids. Our study of the Early Pleistocene *Ursus etruscus* TSR specimen found that all the features observed show the primitive state (straight forehead, caudo-rostrally elongated and forward placed frontal sinuses, narrow rostral frontal sinus, thin and straight palatine, thin and non-pneumatized basioccipital, hemispheric cribriform plate with slight constriction, arched anterodorsal profile of the brain cavity), confirming the basal position of *U. etruscus* among the Pleistocene European ursids. No spelaeoid (derived) characters were recognized, indicating that the endocranial morphology of *U. etruscus* is different from that of the cave bears. On the other hand, *Ursus etruscus* is more similar with the brown bear *U. arctos*, a species retaining the primitive morphology. The relatively thick palate of *U. dolinensis* from Gran Dolina differentiates *U. etruscus* from the former species; however, other endocranial features could be compared. The morphology

of the caudal part of the frontal sinuses and the antero-dorsal profile of the endocast show similarities with *Ursus* sp. from Deutsch-Altenburg 4.

7. The Greek fossil record of *U. etruscus*

Ursus etruscus, accompanied by *Agriotherium* sp., is reported from the early Villafranchian locality Milia (Grevena Basin, Western Macedonia) by an isolated lower canine and some postcranial bones (Tsoukala et al., 2014). This could represent the first appearance of *U. etruscus* in Greece, but additional diagnostic specimens are required to confirm this observation. Regardless of the specific attribution, Milia signals the earliest occurrence of the genus *Ursus* in Greece. The earliest secure occurrence of *U. etruscus* in Greece is in the localities Dafnero 1 (Western Macedonia) and possibly in Sesklon (Thessaly), both dated to the middle Villafranchian, at ~2.0 Ma (Koufos and Kostopoulos, 2016) (Fig. 10). Only a single P4 (DFN-193) is known from Dafnero 1 (Koufos and Kostopoulos, 1997). Its morphology fits well with that of the TSR sample and its size is within the range of variation of *U. etruscus* from the various European sites (Fig. 6b). In comparison with the P4 TSR-E21-50, its protocone is smaller and situated more mesially. Some ursid remains from Sesklon have been described as *U. cf. etruscus*; that material includes two mandibular fragments (No 84Σ) and another mandibular fragment with m1–m2 sin, No 88Σ (Symeonidis, 1992). One of these specimens, probably No 88Σ, was illustrated by Symeonidis (1992: pl. 9, figs 5, 5a, 5b), but the teeth are partially broken and worn and therefore do not provide information on their morphology. The author did not give a description and the provided measurements do not fit with those of the lower dentition of *U. etruscus*; e.g., the given maximal breadth is 25.0 and 24.0 mm for the m1 and m2 respectively; this is probably a typographical error confounding length with breadth. New measurements of the teeth, kindly provided to us by Dr. A. Athanassiou, suggest that both m1

and m2 fall within the *U. etruscus* ranges from Europe (Fig. 6d, e). As the occlusal surface of the teeth is not preserved their morphology cannot be compared with that of other samples. Therefore we believe that the Sesklon ursid must be referred to with its original determination as *U. cf. etruscus*. In the original description of the carnivores from the locality Volax (Drama Basin, Eastern Macedonia), dated to the middle Villafranchian, at ~2.0 Ma (Koufos and Kostopoulos, 2016), a new ursid genus and species *Bosdagius felinus* was described (Sickenberg, 1968: pl. 1; figs 1a–c). This new taxon is based on a single DP3, whose morphology indicates clearly that it is a deciduous carnassial of a hyaenid (Spassov, 2003). We fully agree with this designation and do not further consider this specimen as representing an ursid.

Ursus etruscus has also been reported from Vassiloudi (Mygdonia Basin), correlated to the beginning of the late Villafranchian (~1.8 Ma) (Fig. 10), (Koufos and Kostopoulos, 1997, 2016). This material comprises only a single upper canine (VSL-20), whose morphology and size (Fig. 6a) fits with those of *U. etruscus* from TSR, Upper Valdarno and Olivola. One upper (APL-549) and one lower (APL-550) canine, recovered from the latest Villafranchian (~1.2 Ma) locality Apollonia 1 (Mygdonia Basin) have similar morphology and size (Fig. 6a, c) to those of *U. etruscus* (Koufos and Kostopoulos, 1997, 2016). However, as the material from these localities is scanty it is better to refer them as *U. cf. etruscus* until further specimens confirm the presence of the taxon. One calcaneus (LIB-179) and one third phalanx (LIB-255) are known from the late Villafranchian locality Libakos (Western Macedonia), which were described as *Ursus* sp. aff. *arctos/deningeri* by Steensma (1988). The Libakos calcaneus is larger than that of TSR and the *U. etruscus* specimens from Pirro Nord and Ceyssaguet (France), and it is closer to *U. deningeri* (Fig. 6f). The dimensions of the third phalanx ($DAP_{prox.} = 24.5+$, $DT_{prox.} = 15.1$ mm) are slightly larger than those ($DAP_{prox.} = 24.4–25.8$, $DT_{prox.} = 12–12.8$ mm; Mazza and Rustioni, 1992) of the specimen

IGF-915 of *U. etruscus* from Upper Valdarno. Due to the limited material and the large size of the Libakos calcaneus and third phalanx, we propose that they are best referred to as *Ursus* sp. The species *U. etruscus* is possibly also present in two other Early Pleistocene localities of Southern Greece, Kastritsi near Patras (Achaia, Peloponnese) and Makinia (Etoloakarnania, Central Greece). The material described as *U. cf. etruscus* there includes an upper canine and a fragment of a humerus from the former locality, and an upper third incisor from the latter one (Symeonidis et al., 1985/86).

The biostratigraphic distribution of the Greek ursids is shown in Fig. 10. The earliest ursid evidence is a mandibular fragment preserving the m1–m2 from the middle Turolian (late Miocene) locality Pikermi, described as *Hyaenarctos atticus* and later transferred to *Indarctos* (see Weithofer, 1888; Roussiakis, 2001). The ursids *Ursavus cf. depereti* and *Indarctos atticus* were recognized in the late Miocene faunas of Samos Q1 (Helbing, 1932; Bernor et al., 1996), while later the species *Ursavus ehrenbergi* and *Ursavus depereti* were discovered from the localities of Halmyropotamos and Perivolaki, respectively (Brunner, 1941; Thenius, 1947; Koufos, 2006). All the above localities are dated between ~7.3 and 7.1 Ma (Fig. 10). Ursids re-appeared in the late Pliocene (early Villafranchian) with the genus *Ursus*, while *U. etruscus* was well known during the middle and late Villafranchian, and probably present until the end of the Villafranchian. During the Middle Pleistocene, *U. deningeri* was represented in the archaic fauna of the Petralona cave (Tsoukala, 1989, 1991; Baryshnikov and Tsoukala, 2010). In the Late Pleistocene, bears of the spelaeoid group are known from various Greek caves (e.g., Mitzopoulos, 1960; Symeonidis and Theodorou, 1979; Koufos, 1981; Tsoukala et al., 2006; Baryshnikov and Tsoukala, 2010). The brown bear *U. arctos* is recognized during the Late Pleistocene in the Vraona and Petralona caves (Symeonidis et al., 1980; Baryshnikov and Tsoukala, 2010), and possibly in the Cave Z of

Almopia, Macedonia, Greece (Tsoukala et al., 2006). *Ursus arctos* is still present in the Greek mammal fauna, living in some mountainous areas of Northern Greece.

8. Conclusions

The study of the ursid material from the locality TSR provides new information for *U. etruscus*. Although the species has previously been recognized in Greece, its presence was questionable, as the known material was limited and fragmentary. The sample from TSR is to date the richest and most complete found in Greece and confirms the presence of *U. etruscus* in the Villafranchian of Greece and in Southeastern Europe. The morphological characters of the TSR *U. etruscus* suggest similarities with the youngest forms from Italy and specifically with the late Villafranchian locality Pietrafitta. The endocranial morphology of *U. etruscus* was described here in detail for first time, indicating primitive features, which clearly differ from the more derived *U. deningeri* and *U. spelaeus*, and are more similar to those observed in *U. arctos*. Finally, the species *U. etruscus* first occurred in Greece in the middle Villafranchian, at ~2.0 Ma, in the localities of Dafnero 1 and possibly Sesklon, and survived probably until the latest Villafranchian, making its last appearance in the locality Apollonia 1, biochronologically dated to the latest Villafranchian, at ~1.2 Ma.

Acknowledgements

The excavations in the locality Tsiotra Vryssi were supported from 2014 to 2016 by the ERC STG no. 283503 PaGE ('Paleoanthropology at the Gates of Europe: Human evolution in the southern Balkans') and in 2017 by the ERC COG no. 724703 CROSSROADS ('Human Evolution at the Crossroads'). We wish to thank all the field team members for their contribution during the fieldwork. Many thanks to Dr. H. Scherf (University of Tübingen) for the CT scanning process, Dr. A. Athanassiou (Greek Ministry of Culture) for the photos and

measurements of the Sesklon ursids, D. Giusti (University of Tübingen) for the excavation map of TSR and to M. Ioannidou (University of Tübingen) for help with the Avizo software. We are grateful to the two reviewers P. Mazza and J. Wagner for their constructive comments and suggestions that improved significantly the manuscript.

References

- Argant, A., 2004. Les Carnivores du gisement Pliocène final de Saint-Vallier (Drôme, France). *Geobios* 37, S133–S182.
- Arzarello, M., De Weyer, L., Peretto, C., 2016. The first European peopling and the Italian case: Peculiarities and “opportunism”. *Quaternary International* 393, 41–50.
- Baryshnikov, G.F., 1992. *Ursus mediterraneus* in the Pleistocene of the Caucasus and comments on the history of small bears of Eurasia. *Proceeding of the Zoological institute USSR AS*, 238, 3–60 (in Russian with English summary).
- Baryshnikov, G.F., 2007. Ursidae. In: *Fauna of Russia and neighboring countries, Mammals*. Nauka, Saint Petersburg, pp. 1–554. (in Russian).
- Baryshnikov, G.F., Tsoukala, E., 2010. New analysis of the Pleistocene carnivores from Petralona cave (Macedonia, Greece) based on the collection of the Thessaloniki Aristotle University. *Geobios* 43, 389–402.
- Bernor, R.L., Solounias, N., Swisher III, C.C., van Couvering, J.A., 1996. The correlation of the three classical “Pikermian” mammal faunas - Maragheh, Samos and Pikermi - with the European MN unit system. In: Bernor, R.L., Fahlbusch, V., Mittman, H.W. (Eds.), *The Evolution of Western Eurasian Neogene Mammal Faunas*. Columbia University Press, New York, pp. 137–154.

- Brunner, J., 1941. Ein Bärenrest aus dem Unterpliozän von Euboea. Anzeiger der Akademie der Wissenschaften in Wien, Mathematisch-naturwissenschaftliche Klasse 3, 24–27.
- Drexler, H., 1931. Über Hirnschädelausgüsse von *Ursus spelaeus*. In: Abel, O., Kyrle, G. (Eds.), Die Drachenhöhle von Mixnitz. Speläologische Monographien, pp. 498–536.
- Duval, M., Falguères, C., Bahain, J.J., Grün, R., Shao, Q., Aubert, M., Dolo, J.M., Agustí, J., Martínez-Navarro, B., Palmqvist, P., Toro-Moyano, I., 2012. On the limits of using combined U-series/ESR method to date fossil teeth from two Early Pleistocene archaeological sites of the Orce area (Guadix-Baza basin, Spain). Quaternary Research 77, 482–491.
- García, N., 2004. New results on the remains of Ursidae from Untermaßfeld: comparisons with *Ursus dolinensis* from Atapuerca and other Early and Middle Pleistocene sites, 18th International Senckenberg Conference. Terra Nostra - Schriften der Alfred-Wegener-Stiftung, Weimar, pp. 112–113.
- García, N., Arsuaga, J.L., 2001. *Ursus dolinensis*: a new species of Early Pleistocene ursid from Trinchera Dolina, Atapuerca (Spain). Comptes Rendus Academie Sciences Paris, Sciences de la Terre et des planètes 332, 717–725.
- García, N., Santos, E., Arsuaga, J.L., Carretero, J.M., 2006. High-resolution x-ray computed tomography applied to the study of some endocranial traits in cave and brown bears. Scientific Annals, School of Geology, Aristotle University of Thessaloniki 98, 141–146.
- García, N., Santos, E., Arsuaga, J.L., Carretero, J.M., 2007. Endocranial morphology of the *Ursus deningeri* von Reichenau 1904 from the Sima de Los Huesos (Sierra de Atapuerca) Middle Pleistocene Site. Journal of Vertebrate Paleontology 27, 1007–1017.
- Groiss, J.T., 1994. Untersuchungen der Gehirnmorphologie von *Ursus deningeri* v. Reichenau und von *Ursus spelaeus* Rosenmüller (Mammalia, Ursidae) an

- Schädelausgüssen quartärer Funde aus österreichischen Höhlen. Abhandlungen der Geologischen Bundesanstalt 50, 115–123.
- Guérin, C., Faure, M., Argant, A., Argant, J., Grégut-Bonnoure, É. et al., 2004. Le gisement pliocène supérieur de Saint-Vallier (Drôme, France): synthèse biostratigraphique et paléoécologique. *Geobios* 37, S349–S360.
- Hammer, Ø., Harper, D.A.T., Ryan, P.D., 2001. PAST: paleontological statistics software package for education and data analysis. *Palaeontologia Electronica* 4, 1–9.
- Helbing, H., 1932. Über einen *Indarctos*-Schädel aus dem Pontien der Insel Samos. Nebst einem Anhang: *Hyaenarctos* spec. aus dem Pliocaen von Viallette (Haute-Loire). *Abhandlungen der Schweizerischen Palaeontologischen Gesellschaft* 52, 1–18.
- Kahlke, R.D., 2006. Untermassfeld. A late Early Pleistocene (Epivillafranchian) fossil site near Meiningen (Thuringia, Germany) and its position in the development of the European mammal fauna. *Bar International series* 1578, 1–141.
- Koby, F.E., 1949. Le dimorphisme sexuel des canines d'*Ursus arctos* et d'*U. spelaeus*. *Revue Suisse de Zoologie* 56, 675–687.
- Konidaris, G.E., Tzouroukakis, V., Kostopoulos, D.S., Thompson, N., Giusti, D., Michailidis, D., Koufos, G.D., Harvati, K., 2015. Two new vertebrate localities from the Early Pleistocene of Mygdonia Basin (Macedonia, Greece): Preliminary results. *Comptes Rendus Palevol* 14, 353–362.
- Konidaris, G.E., Kostopoulos, D.S., Koufos, G.D., Tzouroukakis, V., Harvati, K., 2016. Tsiotra Vryssi: A new vertebrate locality from the Early Pleistocene of Mygdonia Basin (Macedonia, Greece). XIV Annual Meeting of the European Association of Vertebrate Palaeontologists. Koninklijke Nederlandse Akademie van Wetenschappen, Haarlem, The Netherlands, p. 37.

- Kostopoulos, D.S., Vasileiadou, K., 2006. The Greek late Neogene-Quaternary ursids in relation to palaeogeography and palaeoenvironment. *Scientific Annals, School of Geology, Aristotle University of Thessaloniki* 98, 285–292.
- Koufos, G.D., 1981. A new Late Pleistocene (Wurmian mammal locality from the basin of Drama (Northern Greece). *Scientific Annals, Faculty of Sciences, Aristotle University of Thessaloniki*, 21, 129–148.
- Koufos, G.D., 2006. Carnivora. In: Koufos, G.D. (Ed.), *The late Miocene vertebrate locality of Perivolaki, Thessaly, Greece. Palaeontographica Abt. A*, Stuttgart, pp. 39–74.
- Koufos, G.D., 2013. Neogene mammal biostratigraphy and chronology of Greece. In: Wang, X., Flynn, L.J., Fortelius, M. (Eds.), *Fossil mammals of Asia. Neogene biostratigraphy and chronology*. Columbia University Press, New York, pp. 595–621.
- Koufos, G.D., Kostopoulos, D.S., 1997. New carnivore material from the Plio-Pleistocene of Macedonia (Greece) with the description of a new canid. *Münchner Geowissenschaftliche Abhandlungen* 34, 33–63.
- Koufos, G.D., Kostopoulos, D.S., 2016. The Plio-Pleistocene large mammal record of Greece: Implications for early human dispersals into European. In: Harvati, K., Roksandic, M. (Eds.), *Human evolution in the southern Balkans*. Springer, Dordrecht, pp. 269–280.
- Koufos, G.D., Syrides, G.E., Kostopoulos, D.S., Koliadimou, K.K., 1995. Preliminary results about the stratigraphy and the palaeoenvironment of Mygdonia Basin, Macedonia, Greece. *Geobios* 18, 243–249.
- Kurtén, B., 1955. Sex dimorphism and size trends in the cave bear, *Ursus spelaeus* Rosenmüller and Heinroth. *Acta Zoologica Fennica* 90, 1–48.

- Magnin, R., 1959. Révision de la faune de vertébrés pliocènes de Perpignan: Le Pliocène en France et en Italie. Diplôme d'études supérieures de Géologie, Faculte des Sciences, Université Lyon, pp. 1–164.
- Martínez Navarro, B., 1991. Revisión Sistemática y estudio cuantitativo de la fauna de macromamíferos del yacimiento de Venta Micena (Orce, Granada). Ph.D. Thesis, Universitat Autònoma de Barcelona, Bellaterra.
- Mazza, P., Rustioni, M., 1992. Morphometric revision of the Eurasian species *Ursus etruscus* Cuvier. *Palaeontographia Italica* 79, 101–146.
- Mazza, P., Rustioni, M., 1994. On the phylogeny of Eurasian bears. *Palaeontographica Abt. A* 230, 1-38.
- Medin, T., Martínez-Navarro, B., Rivals, F., Madurell-Malapeira, J., Ros-Montoya, S., Espigares, M.P., Figueirido, B., Rook, L., Palmqvist, P., 2017. Late Villafranchian *Ursus etruscus* and other large carnivorans from the Orce sites (Guadix-Baza basin, Andalusia, southern Spain): Taxonomy, biochronology, paleobiology, and ecogeographical context. *Quaternary International* 431, 20–41.
- Mitzopoulos, M.K., 1960. Über das Vorkommen von *Ursus spelaeus* in Agrapha Gebirge (Griechenland). *Annales Géologiques des Pays Helléniques* 11, 150–153.
- Moullé, P.E., 1992. Les grands mammifères du Pléistocène inférieur de la grotte du Vallonnet (Roquebrune-Cap-Martin, Alpes-Maritimes). Étude paléontologique des Carnivores, Equidé, Suidé et Bovidé. Ph.D. Thesis, Muséum National d'Histoire Naturelle, Paris.
- Musil, R., 2001. Die Ursiden-Reste aus dem Unterpleistozän von Untermassfeld. In: Kahlke, R.D. (Ed.), *Das Pleistozän von Untermassfeld bei Meiningen (Thüringen), Teil 2. Monographien des Römisch-Germanischen Zentralmuseums Mainz*, pp. 633–658.

- Parés, J.M., Arnold, L., Duval, M., Demuro M., Pérez-González, A., Bermúdez de Castro, J.M., Carbonell, E., Arsuaga, J.L., 2013. Reassessing the age of Atapuerca-TD6 (Spain): new paleomagnetic results. *Journal of Archaeological Science* 40, 4586–4595.
- Oms, O., Parés, J.M., Martínez-Navarro, B., Agustí, J., Toro, I., Martínez-Fernández, G., Turq, A., 2000. Early human occupation of Western Europe: Paleomagnetic dates for two paleolithic sites in Spain. *Proceedings of the National Academy of Sciences of the United States of America* 97, 10666–10670.
- Petrucchi, M., Cipullo, A., Martínez-Navarro, B., Rook, L., Sardella, R., 2013. The Late Villafranchian (Early Pleistocene) carnivores (Carnivora, Mammalia) from Pirro Nord (Italy). *Palaeontographica A* 298, 113–145.
- Quiles, J., 2003. Les Ursidae du Pléistocène moyen et supérieur en Midi méditerranéen: Apports paléontologiques, biochronologiques et archéozoologiques. Ph.D. Thesis, Museum National d'Histoire Naturelle, Paris.
- Rabeder, G., 1999. Die Evolution des Höhlenbärengebisses. *Mitteilungen der Kommission für Quartärforschung der österreichischen Akademie der Wissenschaften* 11, 1–102.
- Rabeder, G., Pacher, M., Withalm, G., 2010. Early Pleistocene bear remains from Deutsch-Altenburg (Lower Austria). *Mitteilungen der Kommission für Quartärforschung der Österreichischen Akademie der Wissenschaften* 17, 1–135.
- Roussiakis, S.J. 2001. Postcranial remains of *Indarctos atticus* (Ursidae, Mammalia) from the classical locality of Pikermi (Attica, Greece), with a description of the front limb. *Senckenbergiana lethaea* 81, 347–358.
- Rustioni, M., Mazza, P., 1993. The Late Villafranchian bear from Pietrafitta (Perugia, Central Italy). *Palaeontographia Italica* 80, 51–62.
- Santos, E., García, N., Carretero, J.M., Arsuaga, J.L., Tsoukala, E., 2014. Endocranial traits of the Sima de los Huesos (Atapuerca, Spain) and Petralona (Chalkidiki, Greece)

- Middle Pleistocene ursids. Phylogenetic and biochronological implications. *Annales de Paléontologie* 100, 297–309.
- Sickenberg, O., 1968. Die unterpleistozäne Fauna von Wolaks (Griech.-Mazedonien). II. Die Carnivoren. *Annales Géologiques des Pays Helleniques* 19, 621–646.
- Spassov, N., 2003. The Plio-Pleistocene vertebrate fauna in South-Eastern Europe and the megafaunal migratory waves from the east to Europe. *Revue Paléobiologique Genève*, 22, 197–229.
- Steensma, K.J., 1988. Plio-/Pleistozäne Großäugetiere (Mammalia) aus dem Becken von Kastoria/Grevena, südlich von Neapolis-NW Griechenland. Ph.D. Thesis, Technische Universität Clausthal, Clausthal.
- Stiner, M.C., 1998. Mortality analysis of Pleistocene bears and its paleoanthropological relevance. *Journal of Human Evolution* 34, 303–326.
- Swenson, J.E., Adamič, M., Huber, D., Stokke, S., 2007. Brown bear body mass and growth in northern and southern Europe. *Oecologia* 153, 37–47.
- Symeonidis, N., 1992. Fossil mammals of Early Pleistocene (Villafranchian) age from Sesklon Basin (Volos). *Annales Géologiques des Pays Helleniques* 35, 1–21 (in Greek with English summary).
- Symeonidis, N., Theodorou, G.E., 1979. Ein neuer Nachweis des Höhlenbären in Nord-Griechenland. *Annales Géologiques des Pays Helleniques* 29, 704–715.
- Symeonidis, N., Bachmayer, F., Zapfe, H., 1980. Ergebnisse weiterer Grabungen in der Höhle von Vraona (Attica, Griechenland). *Annales Géologiques des Pays Helleniques* 30, 291–299.
- Symeonidis, N., Theodorou, G.E., Schütt, H., Velitzelos, E., 1985/86. Palaeontological and stratigraphical observations in the area Achaia and Etoloakarnania (W. Greece). *Annales Géologiques des Pays Helleniques* 33, 329–365 (in Greek with English summary).

- Thenius, E., 1947. *Ursavus ehrenbergi* aus dem Pont von Euböa (Griechenland). Sitzungsberichte der Österreichischen Akademie der Wissenschaften 156, 225–249.
- Torres, T., 1988. Osos (Mammalia, Carnivora, Ursidae) del Pleistoceno de la Península Ibérica. Boletín Geológico y Minero 99, 1–316.
- Toro-Moyano, I., Martínez-Navarro, B., Agustí, J., Souday, C., Bermúdez de Castro, J.M., Martín-Torres, M., Fajardo, B., Duval, M., Falguères, C., Oms, O., Parés, J.M., Anadón, P., Julià, R., García-Aguilar, J.M., Moigne, A.M., Espigares, M.P., Ros-Montoya, S., Palmqvist, P., 2013. The oldest human fossil in Europe, from Orce (Spain). Journal of Human Evolution 65, 1–9.
- Tsoukala, E., 1989. Contribution to the study of the Pleistocene fauna of large mammals (Carnivora, Perissodactyla, Artiodactyla) from Petralona cave (Chalkidiki, N. Greece). Ph.D. Thesis, Scientific Annals School of Geology, Aristotle University of Thessaloniki, Thessaloniki (in Greek with English summary).
- Tsoukala, E., 1991. Contribution to the study of the Pleistocene fauna of large mammals (Carnivora, Perissodactyla, Artiodactyla) from Petralona Cave (Chalkidiki, N. Greece). Preliminary report. Comptes Rendus de l'Académie des Sciences Paris 312, 331–336.
- Tsoukala, E., Bonifay, M.F., 2004. The Early Pleistocene carnivores (Mammalia) from Ceyssaguet (Haute-Loire). Paléo 16, 193–242.
- Tsoukala, E., Chatzopoulou, K., Rabeder, G., Pappa, S., Nagel, D., Withalm, G., 2006. Paleontological and stratigraphical research in Loutra Arideas bear cave (Almopia speleopark, (Pella, Macedonia, Greece). Scientific Annals of the School of Geology, Aristotle University of Thessaloniki 98, 41–67.
- Tsoukala, E., van Loghem, W., Lazaridis, G., Mol, D., 2014. Carnivores of the early Villafranchian site of Milia, (Grevena, Macedonia, Greece). Abstracts of the 6th

- International conference on mammoths and their relatives. Scientific Annals of the School of Geology, Aristotle University of Thessaloniki 102, 6–7.
- Van Valkenburgh, B., 1990. Skeletal and dental predictors of body mass in carnivores. In: Damuth, J., MacFadden, B.J. (Eds.), Body size in mammalian paleobiology. Estimation and biological implications. Cambridge University Press, Cambridge, pp. 181–205.
- Viret, J., 1954. Le loess à bancs durcis de Saint-Vallier (Drôme) et sa faune de mammifères villafranchiens. Nouvelles Archives Museum Histoire naturelle Lyon 4, 1–200.
- Weithofer, A. 1888. Beiträge zur Kenntniss der Fauna von Pikermi bei Athen. Beiträge zur Paläontologie Österreich-Ungarns 6, 225–292.

FIGURES CAPTIONS

Figure 1. **a**, Sketch map indicating the position of the locality Tsiotra Vryssi in the Mygdonia Basin; the map was taken from www.shaded-relief.com; **b**, excavation map of Tsiotra Vryssi indicating the points where the studied ursid remains were found. The pedal remains are not shown, as they were recovered during the test excavation of the site.

Figure 2. *Ursus etruscus*, Tsiotra Vryssi, Mygdonia Basin, Macedonia, Greece, late-Villafranchian. **a–e**, cranium, TSR-E21-50; **a**, right lateral; **b**, left lateral; **c**, dorsal; **d**, ventral; **e**, tooththrow's occlusal view; **f–g**, maxillary fragment with both I1 and left canine, TSR-E21-63; **f**, dorsal, and **g**, ventral view; **h–i**, right maxillary fragment with P4–M1, TSR-E16-18; **h**, lateral; **i**, occlusal view; **j–l**, right isolated P3, TSR-E18-19; **j**, buccal, **k**, lingual and **l**, occlusal view; **m**, left isolated m1, TSR-E19-34; occlusal view.

Figure 3. *Ursus etruscus*, Tsiotra Vryssi, Mygdonia Basin, Macedonia, Greece, late Villafranchian. **a–b**, atlas, TSR-E21-52a; **a**, ventral; **b**, dorsal view; **c–d**, axis, TSR-E21-52b; **c**, ventral; **d**, dorsal view; **e–f**, proximal part of right ulna, TSR-C16-26; **e**, ventral; **f**, dorsal

view; **g–h**, left calcaneus, TSR-160; **g**, medial; **h**, ventral view; **i**, left posterior foot, TSR-160, ventral view *in situ*; **j**, left metatarsals with the preserved phalanges, TSR-160; dorsal view.

Figure 4. Discriminant analysis of the dental dimensions of *U. etruscus*, *U. rodei*, *U. deningeri*, *U. spelaeus* and *U. arctos* from various European localities. The used specimens are complete P4–M2 tooth rows. The table gives the classification of the TSR specimens in relation to the samples used. Data sources: [Mazza and Rustioni \(1992\)](#) for Upper Valdarno, Olivola, Pietrafitta and Saint-Vallier *U. etruscus*; [Petrucci et al. \(2013\)](#) for Pirro Nord *U. etruscus*; [Tsoukala and Bonifay \(2004\)](#) for Ceyssaguet *U. etruscus*; [Musil \(2001\)](#) for *U. rodei*; [Viret \(1954\)](#) for *U. arctos*; [Quiles \(2003\)](#) and [Santos et al. \(2014\)](#) for *U. deningeri*; [Quiles \(2003\)](#) for *U. spelaeus*.

Figure 5. **a**, Simpson’s log-ratio diagram comparing the dimensions of the TSR upper teeth with *U. etruscus* from various European localities. Reference sample: *U. arctos*, Spain, n = 18–52 ([Torres, 1988](#)); **b**, Simpson’s log-ratio diagram comparing the dimensions of the TSR upper teeth with various European ursid taxa. Reference sample: *U. arctos*, Spain ([Torres, 1988](#)). Data sources: [Mazza and Rustioni \(1992\)](#) for Upper Valdarno, Olivola, Pirro Nord, Pietrafitta, Senèze, Saint-Vallier and Perrier *U. etruscus*; [Petrucci et al. \(2013\)](#) for Pirro Nord *U. etruscus*; [Magnin \(1959\)](#) for *U. ruscinensis*; [Tsoukala \(1989\)](#) for Petralona *U. deningeri*; [Musil \(2001\)](#) for Untermassfeld *U. rodei*; [Garcia and Arsuaga \(2001\)](#) for *U. dolinensis*; [Quilles \(2003\)](#) for Vallonnet, Mosbach and Spanish *U. deningeri*; [Tsoukala and Bonifay \(2004\)](#) for Ceyssaguet *U. etruscus*.

Figure 6. Bivariate plot (length/breadth) comparing the C, P4, c, m1, m2 and calcaneus dimensions of the Greek ursids with samples from various European localities. Data sources as in [Fig. 4](#) plus [Rabeder et al. \(2010\)](#) for Deutsch-Altenburg ursid and [Moullé \(1992\)](#) for Vallonnet *U. deningeri*.

Figure 7. CT-scan slices of the *Ursus etruscus* cranium from Tsiotra Vryssi (TSR-E21-50). **a–b**, parasagittal slices; **c**, horizontal slice, showing the frontal sinuses; **d**, coronal slice; not to scale.

Figure 8. a–c, Virtual reconstruction of the *Ursus etruscus* cranium from Tsiotra Vryssi (TSR-E21-50); **a**, lateral view with mirror-imaged the zygomatic arch of the opposite side; **b**, dorsal view; **c**, nuchal view; **d**, coronal section at the level of the tympanic bulla; **e–f**, virtual casts of the brain and the frontal sinuses in anatomical position; **e**, dorsal view; **f**, lateral view (transparent); **g**, sagittal section; **h–i**, close view of the virtually reconstructed right tooth row; **h**, occlusal view; **i**, lateral view; **j**, virtual reconstruction of the frontal sinuses (reversed), lateral view; **k**, virtual reconstruction of the brain, lateral view.

Figure 9. Box plot diagrams comparing (mean and standard deviation) the palate (a) and the basioccipital thickness (b) of *U. etruscus* (Tsiotra Vryssi), *U. arctos* (Iberian localities), *U. dolinensis* (Trinchera Dolina), *U. deningeri* (Sima de los Huesos and Petralona) and *U. spelaeus* (Iberian and Greek localities). Data from Santos et al. (2014).

Figure 10. Biostratigraphic table with the chronological distribution of ursid taxa in the Neogene/Quaternary of Greece.

LEGENDS OF TABLES

Table 1. *Ursus etruscus*, Tsiotra Vryssi (TSR), cranial measurements; according to Mazza and Rustioni (1992).

1. length prosthion–akrokranium; 2. length prosthion–occipital condyles; 3. length prosthion–basion; 4. length basion–nasion; 5. length frontal midpoint–akrokranium; 6. length prosthion–

nasion; **7.** length prosthion–frontal midpoint; **8.** length nasion–rhinion; **9.** length prosthion–orbital cavity; **10.** length prosthion–staphylion; **11.** length staphylion–palatino-orale; **12.** breadth otion–otion; **13.** least breadth between temporal crests; **14.** greatest breadth of the occipital condyles; **15.** greatest breadth of the paroccipital processes; **16.** greatest breadth of the supraoccipital crest; **17.** greatest breadth of the foramen magnum; **18.** greatest breadth between the euryons; **19.** greatest zygomatic breadth; **20.** least postorbital breadth; **21.** greatest frontal breadth; **22.** least breadth between the orbits; **23.** greatest palatal breadth; **24.** least breadth behind the canine alveoli; **25.** greatest breadth of the canine alveoli; **26.** greatest height of the orbital cavities.

Table 2. *Ursus etruscus*, Tsiotra Vryssi (TSR), dental measurements.

Table 3. *Ursus etruscus*, Tsiotra Vryssi (TSR), postcranial measurements according to **Mazza and Rustioni (1992)**.

Table 4. *Ursus etruscus*, Tsiotra Vryssi (TSR), measurements of the endocast and the frontal sinuses; according to **García et al. (2007)**.

a

b

e

f

g

5cm

h

j

k

l

m

1cm

i

a

b

e

5cm

f

c

d

i

g

h

2cm

j

	Given group	Classification	Jackknifed		Given group	Classification	Jackknifed
TSR -E21-50 dex	?	a		Ceyssagueut	e	e	e
TSR -E21-50 sin	?	a		Ceyssagueut	e	e	e
TSR -E16-18	?	a		Ceyssagueut	e	e	e
Upper Valdarno	a	d	d	Ceyssagueut	e	e	e
Upper Valdarno	a	a	a	Ceyssagueut	e	e	e
Upper Valdarno	a	a	a	<i>U. rodei</i>	f	f	f
Upper Valdarno	a	a	c	<i>U. rodei</i>	f	f	f
Upper Valdarno	a	a	b	<i>U. deningeri</i>	g	g	h
Upper Valdarno	a	a	a	<i>U. deningeri</i>	g	h	h
Olivola	a	a	e	<i>U. deningeri</i>	g	g	g
Olivola	a	e	e	<i>U. deningeri</i>	g	g	h
Pietrafitta	b	b	a	<i>U. spelaeus</i>	h	h	h
Saint-Vallier	c	a	a	<i>U. spelaeus</i>	h	h	g
Saint-Vallier	c	c	c	<i>U. spelaeus</i>	h	h	h
Saint-Vallier	c	c	c	<i>U. spelaeus</i>	h	h	g
Saint-Vallier	c	c	i	<i>U. spelaeus</i>	h	h	g
Saint-Vallier	c	c	c	<i>U. spelaeus</i>	h	g	g
Saint-Vallier	c	c	a	<i>U. arctos</i>	i	i	h
Pirro Nord	d	d	a	<i>U. arctos</i>	i	i	i
Ceyssagueut	e	e	e	<i>U. arctos</i>	i	i	i
Ceyssagueut	e	e	e	<i>U. arctos</i>	i	c	c
				% correctly classified		84.62	

a

b

a **b** **c** **d** **e** **f**

a

b

c

d

e

f

10 cm
a-g

g

h

i

j

k

5 cm
h-k

a

b

EPOCH		AGE (Ma)	CHRONS	LOCALITIES	TAXA	
HOLOCENE						
PLEISTOCENE	LATE	0.0		Vraona, Attica Cave Agrapha Cave Perama, Ioannina Cave A (Aridea) Cave Maara, Drama Petalona Cave (young fauna)	<i>Ursus arctos</i> ' <i>Ursus spelaeus</i> group' ' <i>Ursus spelaeus</i> group' <i>Ursus ingressus</i> <i>Ursus spelaeus</i> (out of the cave) <i>Ursus spelaeus</i>	
	MIDDLE	0.5	C1n	Petalona Cave (archaic fauna)	<i>Ursus deningeri</i>	
	EARLY	1.0	C1r	Apollonia 1	<i>Ursus cf. etruscus</i>	
		1.5		TSIOTRA VRYSSI (TSR) Libakos	<i>URSUS ETRUSCUS</i> <i>Ursus sp.</i>	
		2.0		Vassiloudi Dafnero 1, Sesklon	<i>Ursus cf. etruscus</i> <i>Ursus etruscus</i>	
		2.5				
	PLIOCENE	VILLAFRANCHIAN	3.0	C2An	Milia	<i>Ursus sp., Agriotherium</i>
			3.5			
		RUSCINIAN	4.0	C2Ar		
			4.5			
TUROLIAN		5.0	C3n			
		5.5				
		6.0				
		6.5				
MIOCENE		TUROLIAN	7.0	C3r		
			7.5	C3An		
	8.0		C3Ar			
	8.5		C3Bn			
	9.0		C3Br			
		9.5	C4n	Samos Q1 (~7.1Ma) → Perivolaki (7.3-7.1 Ma) → Pikermi (7.37-7.11 Ma)	<i>Ursavus cf. depereti</i> , <i>Indarctos atticus</i> <i>Ursavus depereti</i> <i>Ursavus sp., Indarctos atticus</i>	

?

Halmypopotamos

?

Halmypopotamos