

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**REVISION: JNS-D-18-00762R2**

**Journal of Neurological Sciences - Letter to the Editor**

**Epileptic chorea: another window into neural networks?**

Fernando Tenório Gameleira, MD;<sup>1</sup> Karina Torres Alúcio;<sup>1</sup> Maria Luísa Maia Nobre de Paiva;<sup>1</sup> Kathleen Caroline de Lima Carlos;<sup>1</sup> Kedma Mayara de Lima;<sup>1</sup> Mariana Holanda Gameleira;<sup>2</sup> Marcelo A Kauffman, MD, PhD;<sup>3</sup> Alberto J. Espay, MD, MSc.<sup>4</sup>

1. Programme of Epilepsy Surgery of the University Hospital of the Federal University of Alagoas, Brazil.
2. Tiradentes University of Alagoas, Brazil.
3. Consultorio y Laboratorio de Neurogenética, Centro Universitario de Neurología José María Ramos Mejía. Buenos Aires, Argentina
4. James J. and Joan A. Gardner Center for Parkinson disease and Movement Disorders, Department of Neurology and Rehabilitation Medicine, University of Cincinnati, Cincinnati, Ohio, USA.

**Word count:** 888; Title character count (with spaces): 54; Figures: 1; References: 10;

Supplemental material: 4 (Video: 3; Additional EEG data: 1)

**Keywords:** chorea, epilepsy, paroxysmal nonkinesigenic dyskinesia

**All correspondence to:**

Dr. Alberto J Espay, Gardner Center for Parkinson's disease and Movement Disorders

University of Cincinnati Academic Health Center, 260 Stetson St., Suite 2300

Cincinnati, OH 45267-0525 - E-mail: [alberto.espay@uc.edu](mailto:alberto.espay@uc.edu)

**Funding source:** None

1 **Financial disclosures related to research covered in this article:** The authors report  
2 no conflict of interest as related to research covered in this article.  
3  
4  
5  
6  
7

8  
9 **Complete Disclosures**

10  
11  
12 **Drs. Tenório Gameleira, Torres Alúcio, Maia Nobre de Paiva; de Lima Carlos;**  
13 **de Lima; Gameleira** have nothing to disclose.  
14  
15  
16

17  
18 **Dr. Kauffman** is an employee of the CONICET. He has received grant support from  
19  
20 Ministry of Science and Technology of Argentina and Ministry of Health of Buenos  
21  
22 Aires.  
23  
24  
25

26 **Dr. Espay** has received grant support from the NIH, Great Lakes Neurotechnologies  
27  
28 and the Michael J Fox Foundation; personal compensation as a consultant/scientific  
29  
30 advisory board member for Abbvie, Neuroderm, TEVA, Impax, Acadia, Acorda,  
31  
32 Cynapsus/Sunovion, Lundbeck, Osmotica Pharmaceutical, and USWorldMeds;  
33  
34 publishing royalties from Lippincott Williams & Wilkins, Cambridge University  
35  
36 Press, and Springer; and honoraria from Abbvie, UCB, USWorldMeds, Lundbeck,  
37  
38 Acadia, the American Academy of Neurology, and the Movement Disorders Society.  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Dear Editor,

Paroxysmal abnormalities of motor or non-motor function are defined as epileptic when they co-occur with specific patterns of organized electrical activity as measured by scalp electroencephalography (EEG). Certain forms of abdominal pain and migraine may be therefore epileptic.<sup>1</sup> Stereotypical movements at night can be the result of mesial frontal epilepsy, despite a negative EEG and *prolonged* focal paroxysmal dystonia can result from unilateral putaminal hemorrhage.<sup>2</sup> Short-lasting motor activity categorized clinically as dystonia or athetosis, meeting criteria for paroxysmal non-kinesigenic dyskinesia (PNKD), may result from caudate nucleus rather than cortical discharges.<sup>3,4</sup> These examples suggest that epileptic activity affecting different neural networks may manifest unique motor and non-motor clinical phenotypes, beyond the classic myoclonic or tonic-clonic motor phenomena attributed to cortical epilepsy. To this end, using video-EEG, we documented ictal discharges in the left temporal lobe of a patient with episodic chorea, initially suspected to be within the PNKD spectrum, suggesting that paroxysmal chorea can also be of epileptic origin in a subtype of patients.

This 40-year-old woman had brief 90-second episodes of generalized choreoathetosis since the age of 6 months (Video 1). These episodes (10-15/day) included a brief paresthetic aura, were more prominent upon awakening, and alternated with minor spells (3-4/day) that included chorea in the perioral and pharyngeal-laryngeal musculature, sometimes including the upper limbs, associated with dysarthria (Video 2). Consciousness was invariably preserved. She consistently leaned to her right in the immediate post-ictal period. She felt exhausted but coherent in the post-ictal period. Episodes did not occur during each of her three pregnancies but returned within days

1 from delivery. There was no effect by caffeinated or alcoholic beverages but physical  
2 or emotional stress increased their frequency. Prior treatment with antidepressants,  
3  
4  
5 anxiolytics and neuroleptics had been of no benefit. The episodes became more  
6  
7 frequent after the age of 20 years.  
8  
9

10  
11 Despite subjective global weakness, the interictal neurological exam was normal. On  
12  
13 neuropsychological evaluation, she exhibited frontal dysfunction, with most severe  
14  
15 impairment in tasks measuring inhibitory control (Stroop Test II and III), alternating  
16  
17 attention (20<sup>th</sup> percentile); long-term visual memory (Wechsler Memory Scale-III  
18  
19 delayed visual memory; <1 percentile); and mental flexibility (Trail Making Test B,  
20  
21 without errors; <1 percentile). Video-EEG showed initial epileptiform activity in the  
22  
23 left temporal region, 7 seconds after clinical onset (Figure, A). The interictal EEG  
24  
25 showed wicket spikes-type pattern on the left frontotemporal region (Figure, B),  
26  
27 generally considered a benign variant. Full ictal and interictal EEG is available online  
28  
29 (supplementary material). Brain MRI was normal. Whole Exome Sequencing was  
30  
31 negative for any genetic findings of etiologic relevance. Phenytoin 300 mg/day and  
32  
33 clonazepam 2 mg/day reduced the frequency and severity of episodes by about 80%.  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43

44 Preserved consciousness and normal interictal exam had suggested a functional  
45  
46 etiology. However, onset since early childhood and a high frequency of episodes  
47  
48 permitted video-EEG assessments to disclose its epileptic nature (Video 3). Unlike  
49  
50 classic PNKD, whereby dystonic or athetotic movements are predominantly  
51  
52 appendicular, unilateral or asymmetric, and often triggered by alcohol or caffeinated  
53  
54 beverages, the epileptic events documented in this patient were generalized, the  
55  
56 phenomenology was chorea rather than dystonia, and there were no identifiable  
57  
58  
59  
60  
61  
62  
63  
64  
65

triggers.

1  
2  
3  
4  
5 Whereas an epileptic mechanism has been demonstrated in some types of paroxysmal  
6  
7 kinesigenic dyskinesia (PKD),<sup>3,5</sup> to our knowledge this is the first case of a PNKD-  
8  
9 like disorder with an abnormal electrographic correlate. This observation is consistent  
10  
11 with the growing recognition that some genetic disorders can manifest both a  
12  
13 movement and an epileptic phenotype. *PRRT2* mutations have been recognized as the  
14  
15 etiology of benign familial infantile epilepsy (BFIE), infantile convulsions with  
16  
17 choreoathetosis (ICCA) syndrome, and PKD.<sup>6</sup> Similarly, *SCL2A1* gene mutations  
18  
19 (GLUT1 deficiency syndrome) can lead to manifestations ranging from early-onset  
20  
21 epilepsy with myoclonic seizures to paroxysmal exercise-induced dystonia,<sup>7</sup> and  
22  
23 *KCNMA1* gene mutations allows the coexistence of epilepsy and PNKD in the same  
24  
25 individual or family.<sup>8</sup> Furthermore, classic epileptic phenotypes and movement  
26  
27 disorders can be ascertained by the same mutations in *ATPIA3*<sup>9</sup> and *CACNA1A*.<sup>10</sup>  
28  
29  
30  
31  
32  
33  
34 Despite the lack of a defined genetic etiology, which likely reflects our incomplete  
35  
36 understanding of the spectrum of genetic disorders, our patient highlights the complex  
37  
38 relationship between cortical and basal ganglia circuits and raises the possibility that  
39  
40 neural networks may have nodes (ictal zones) of differential susceptibility at various  
41  
42 cortical and subcortical levels, capable of yielding relative non-classical seizure  
43  
44 phenotypes.  
45  
46  
47  
48  
49  
50

51 Our case also demonstrated cognitive abnormalities in the frontal lobe, suggesting  
52  
53 chronic dysfunction beyond ictal chorea and arguing for an ictal onset in the frontal  
54  
55 mesial regions, beyond the reach of the scalp EEG (explaining the EEG “silence” for  
56  
57 a few seconds after the clinical onset). The paroxysmal and stereotypical nature of the  
58  
59  
60  
61  
62  
63  
64  
65

1 short episodes, presence of aura, and response to antiepileptic drugs supported their  
2 epileptic nature. Negative video-EEG evaluations in most patients with paroxysmal  
3 chorea or other PNKD-like disorders, with abnormal frontal mesial-caudate neural  
4 network, may be explained by the relative predominance of electrical abnormalities in  
5 the basal ganglia over the cortex, beyond the resolution of scalp electrodes.<sup>4</sup>  
6  
7  
8  
9  
10

11  
12  
13 While we do not have connectivity data to prove that this manifestation reflects a  
14 window into the frontal-basal ganglia network, paroxysmal chorea has not before  
15 been shown to arise from “pure” cortical lesions but rather from basal ganglia  
16 abnormalities. Given the close relationship between the brain cortex and the basal  
17 ganglia, it remains a mystery why chorea and other hyperkinetic movement disorders  
18 are not more commonly documented as ictal (network-based) epileptic manifestation.  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## Authors' Contributions

A. Drafting of the case report; B. acquisition of data; C. analysis and interpretation;  
D. critical revision of the manuscript for important intellectual content.

Fernando Tenório Gameleira: A, B, C, D

Karina Torres Alúcio: B, C, D

Maria Luísa Maia Nobre de Paiva: B, D

Kathleen Caroline de Lima Carlos: B, D

Kedma Mayara de Lima: B, D

Mariana Holanda Gameleira: B, C, D

Marcelo Kauffman: B, C

Alberto J. Espay: A, C, D

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
**Figure legend. A.** Ictal EEG, with abnormal epileptiform activity arising on the left temporal region, approximately 7 seconds after the onset of chorea. Given delay in EEG onset in relationship to clinical onset, the ictal onset may have been in a separate region, probably fronto-mesial. A background rhythmic theta becomes prominent toward the end of the clinical episode (see also Supplementary material). **B.** Interictal EEG showing wicket spikes arising in the left frontotemporal region.

17  
18  
**Video legends.**

19  
20  
**Video 1.** Major choreoathetotic event demonstrated during clinic visit.

21  
22  
23  
24  
**Video 2.** Minor choreoathetotic episode with perioral and peri-pharyngeal involvement.

25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
**Video 3.** Video EEG-documented episode. The initial leg movements occurred a few seconds before the EEG onset (theta-delta activity) on the left temporal region. The onset of chorea co-occurred with facial expression changes. The yawn was an epileptic automatism. She retained consciousness throughout the event.

37  
38  
39  
40  
**Supplementary material.**

41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
**EEG of the preictal, ictal, and immediate postictal period.** The first EEG alteration was a rhythmic theta activity on the left temporal area followed by alpha rhythm for a few seconds before evolving back into theta rhythm (mixed with artifacts) prior to the end of the seizure, 70 seconds after onset, shown as global flattening.

51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
**Ethical Compliance Statement:** We confirm that we have read the Journal's position on issues involved in ethical publication and affirm that this work is consistent with



those guidelines. The authors confirm that patient consent was obtained for this work  
but no IRB approval was necessary.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## References

1. Gameleira FT, Ataide L, Jr., Raposo MC. Relations between epileptic seizures and headaches. *Seizure* 2013;22:622-626.
2. Lee DH, Kim DW, Kim HJ. Movement-induced focal tonic-clonic seizure-like movements after ipsilateral basal ganglia hemorrhage. *J Clin Neurosci* 2011;18:285-286.
3. Akiyama T, Ohtsuka Y, Kobayashi K, Oka E. Kinesigenic attacks with ictal electroencephalographic abnormalities. *Pediatr Neurol* 2004;31:357-359.
4. Lombroso CT. Paroxysmal choreoathetosis: an epileptic or non-epileptic disorder? *Ital J Neurol Sci* 1995;16:271-277.
5. Madhusudanan M, Bhatia KP, Balint B. Paroxysmal craniocervical dyskinesia as manifestation of frontal lobe epilepsy. *Movement disorders : official journal of the Movement Disorder Society* 2011;26:2580-2582.
6. Heron SE, Dibbens LM. Role of PRRT2 in common paroxysmal neurological disorders: a gene with remarkable pleiotropy. *J Med Genet* 2013;50:133-139.
7. Urbizu A, Cuenca-Leon E, Raspall-Chaure M, et al. Paroxysmal exercise-induced dyskinesia, writer's cramp, migraine with aura and absence epilepsy in twin brothers with a novel SLC2A1 missense mutation. *J Neurol Sci* 2010;295:110-113.
8. Du W, Bautista JF, Yang H, et al. Calcium-sensitive potassium channelopathy in human epilepsy and paroxysmal movement disorder. *Nat Genet* 2005;37:733-738.
9. Heinzen EL, Arzimanoglou A, Brashear A, et al. Distinct neurological disorders with ATP1A3 mutations. *The Lancet Neurology* 2014;13:503-514.
10. Damaj L, Lupien-Meilleur A, Lortie A, et al. CACNA1A haploinsufficiency causes cognitive impairment, autism and epileptic encephalopathy with mild cerebellar symptoms. *Eur J Hum Genet* 2015;23:1505-1512.

Figure



**e-component**

[Click here to download e-component: Supplementary material\\_Preictal and ictal EEG\\_REV.pptx](#)

**e-component**

[Click here to download e-component: Patients consent.pdf](#)

## Video

[Click here to download Video: Video 1.mov](#)

## Video

[Click here to download Video: Video 2.mov](#)

## Video

[Click here to download Video: Video 3.wmv](#)