Alleviating tension in Λ CDM and the local distance ladder from first principles with no free parameters

Maurice H.P.M. van Putten

Department of Physics and Astronomy
Sejong University, Seoul

MNRAS 491 L6 (2020)

zenodo.org/record/2597378 (March 18 2020)

AAS234 Press Conf.: Cold Quasars & **Hot Cosmology** (June 13 2019), <https://www.youtube.com/watch?v=bNdscedDCnQ>

CQUEST, 서남대학교
August 21 2020

Cosmological Expansion 1.0

Distant nebulae recede in Doppler shift

$$z = \frac{v}{c} = 0.004$$

$$z = 0.05$$

$$z = 0.0733$$

$$z = 0.13$$

$$z = 0.2033$$

Vesto M. Slipher
1875-1969

Slipher, V., 1917, Amer. Phil. Soc., 56, 403

Lemaitre

Conformal expansion by scale factor $a(t)$

$$D = D_0 \frac{a}{a_0}$$

Lemaitre, G., 1927, "Un Univers homogene de masse constante et de rayon croissant, rendant compte de la vitesse radiale des nebuleuses extra-galactiques," Ann. Soc. Sc. Bruxelles, A47, 49-59; 1931, "A homogeneous universe of constant mass and increasing radius accounting for the radial velocity of extra-galactic nebulae," MNRAS, 91, 443

G.H.E.J. Lemaitre (1894-1966)

$$V = D_0 H$$

Expansion parameter:

$$H = \frac{\dot{a}}{a}$$

Lemaitre's Big Bang

E.g. “coasting”: linear expansion in time

$$\frac{a}{a_0} = t : H = \frac{\dot{a}}{a} = \frac{1}{t}$$

Age of the Universe:

$$\Delta t = \frac{1}{H_0}$$

where $H_0 = H(0)$.

Cepheid Variables

- Fairly luminous ($-7 < M < -4$) high-mass stars on the instability strip in the HR-diagram: observable out to few tens of Mpc (notably NGC 4603)

Leavitt, H.S, Pickering, E.C., 1912, "Periods of 25 Variable Stars in the Small Magellanic Cloud". Harvard College Observatory Circular. 173, 1

https://www.atnf.csiro.au/outreach/education/senior/astrophysics/variable_cepheids.html

M31

Observation of the very faint - extragalactic - cepheid variable

Photometry on M31 (1924)

One month periodicity of Hubble's Cepheid V1

M31: $D = 2.3 \text{ Mly}$

“Here is the [Hubble's] letter that destroyed my universe [Milky Way].” Harlow Shaply (1885-1972)

Photometry and spectroscopy on 22 extragalactic nebulae

Edwin Hubble

Spectroscopic data ->

Hubble, E.P.,
1929, PNAS
15, 168

Photometric data ->

Approximately linear relation of receding velocity with distance

Hubble-Lemaitre law

$$V = H_0 D : D = z R_H \quad (R_H = c/H_0)$$

Hubble constant: new parameter of observational cosmology

$$[H_0] = \mathbf{s^{-1}}$$

$$[H_0] = \mathbf{km \ s^{-1} \ Mpc^{-1}}$$

Velocity

Distance scale
between galaxies

Type Ia supernovae

Absolute magnitude $M_B = -19.76 \pm 0.13$ SN1937C (Type Ia): SNe seen much further out than Cepheids ($M_B \gtrsim -5$)

Light curves of supernovae of Type Ia extend over a few weeks during which time they are powered largely by decay of ^{56}Ni and ^{56}Co .

Type Ia: spectra without H, He,
with SiII absorption lines

Mean rise time 17.5d
(over 13-23d in B-band)

Philips normalization

To leading order, Ia light curves effectively show 1 DOF, apparent in a correlation of peak magnitude and decay (defined over 15d)

Philips, M.M., 1993, ApJ, 413, L105

$$\Delta m_{15} = m_{15d} - m_{max}$$

SN Ia: bright standard candles (residual $\lesssim 7\%$)

CE 3.0: Local distance ladder

Distance measurement by bootstrapping:

Riess, A.G., et al., 2016, ApJ, 826, 56

Observational results

CE 1.0 Galaxies (not standard candles) [Slipher 1917]

CE 2.0 $H_0 \simeq 501 \text{ km s}^{-1} \text{Mpc}^{-1}$ [Hubble 1929]

CE 3.0 $H_0 \simeq 74.03 \pm 1.46 \text{ km s}^{-1} \text{Mpc}^{-1}$ [Riess et al. 2019]

Age of the Universe

Quality of measurements evolves with improved understanding of statistical and systematic errors

e.g. Cepheids in Virgo: $H_0 = 72 \pm 2 \pm 7 \text{ km s}^{-1} \text{ Mpc}^{-1}$

Garnavich, P. M., et al., 1998, ApJ, 509, 74

Sandage parameters

THE ASTROPHYSICAL JOURNAL

AN INTERNATIONAL REVIEW OF SPECTROSCOPY AND
ASTRONOMICAL PHYSICS

VOLUME 133

MARCH 1961

NUMBER 2

THE ABILITY OF THE 200-INCH TELESCOPE TO DISCRIMINATE
BETWEEN SELECTED WORLD MODELS

ALLAN SANDAGE

Mount Wilson and Palomar Observatories
Carnegie Institution of Washington, California Institute of Technology
Received October 14, 1960; revised November 5, 1960

Allen Sandage (1926-2010)

*“... the future of observational cosmology would be the search for two parameters: the **Hubble constant** H_0 and the **deceleration parameter** q_0 .”*

https://en.wikipedia.org/wiki/Allan_Sandage

- Cosmological scale factor $a(t)$ be governed by Friedmann equations
- Evolve $H = \frac{\dot{a}}{a}$ and $\frac{d}{dt}H(t)$
- Express results in $H(z)$ and deceleration parameter $q(z)$:

$$q = -\frac{\ddot{a}a^2}{\dot{a}^2} = -1 + (1+z)H^{-1}(z)H'(z), \quad a = \frac{a_0}{1+z}$$

q captures curvature in the graph of $H(z)$ versus z

Context

A three-flat four-covariant spacetime satisfies the Hamiltonian energy constraint of GR (first Friedmann equation)

$$\Omega_M + \Omega_\Lambda = 1 \quad \text{Then} \quad q(z) = \frac{1}{2}\Omega_M - \Omega_\Lambda$$

Expected: $q = \frac{1}{2}$

Matter-dominated era

Observed: $q(z) < 0$

Λ -dominated era

The Nobel Prize in Physics 2011

© The Nobel Foundation. Photo: U. Montan

Saul Perlmutter

Prize share: 1/2

© The Nobel Foundation. Photo: U. Montan

Brian P. Schmidt

Prize share: 1/4

© The Nobel Foundation. Photo: U. Montan

Adam G. Riess

Prize share: 1/4

"Dat doet de Lambda"

de Sitter, AH 1930

Cosmological spacetime

Expanding spacetime is unique to cosmology

Beyond Poincare-invariant Minkowski spacetime with Lorentz invariant light cones:

Minkowski spacetime

Symmetry breaking

Breaking translation-invariance

Minkowski spacetime

$$T = 0$$

Rindler spacetime

($h = \partial D^-$ at distance ξ)

$$k_B T \sim \frac{c}{\xi} \hbar$$

Cosmological spacetime

“A day without yesterday”

$$k_B T \sim H \hbar$$

"Observers were not present at the Big-Bang instant" (Crease & Mann 1996)

Equivalent temperatures by breaking translation invariance in space- or time

de Sitter temperature

de Sitter spacetime
(eternal)

$$k_B T \sim H_0 \hbar = \frac{c}{R_H} \hbar$$

FLRW-cosmologies

$$k_B T \sim H \hbar = \frac{c}{R_H} \hbar$$

Evolving de Sitter temperature in FLRW-cosmologies

Heat comes with mass-energy...

Spacetime thermodynamics

Spacetime thermodynamics

J. Bekenstein
(1947-2015)

Recall that
mass is
energy !

Mass-energy in heat

J. Bekenstein
(1947-2015)

Unit of energy flows “in/out” by diffusion: dM “in/out” of a BH (based on propagators)

$$dM = T_H dS, \quad S = \frac{1}{4} A_H : \quad dM = T_H (8\pi M dM) \rightarrow T_H = \frac{1}{8\pi M}$$

$Q = Mc^2$ is total heat content of a Schwarzschild BH spacetime

Spacetime Q

- Black hole spacetimes: $Q = Mc^2$

By the Bekenstein-Hawking entropy and Hawking temperature of the event horizon.

- Rindler spacetimes: $Q = mc^2$

$$Q = \int_0^\xi T_U dS = U_h = \int_0^\xi F ds = mc^2$$

$$F = ma, \xi = c^2/a$$

$$S = 2\pi\Delta\varphi_C, \Delta\varphi_C = k_C s$$

$$T_U = a\hbar/2\pi c$$

(van Putten, 2017, ApJ, 848, 28; 2018, MNRAS, 281, L26)
(Diffusion constant $D = \hbar c^2/2\pi k_B T$ (Son & Starinets, 2007, Ann Rev. Nucl. Part. Sci., 57, 95): $D = \xi c \rightarrow T = T_U$)

- Cosmological spacetime?

Q at de Sitter temperature of the Hubble horizon is on par with closure density $\rho_c c^2$.

Include Q of cosmological spacetime in Friedmann equation

Reference phase

Boundary term from cosmological horizon:

$$S_0[\mathcal{N}] = 0 \text{ (or some } f(t) \text{) in Minkowski}$$

$S_0[\mathcal{H}] \neq 0$ in FLRW cosmologies

Proper total phase *difference* in path integral formulation:

$$e^{i(S-S_0)/\hbar}$$

Van Putten, 2019, MNRAS, 491, L6

Equivalent energy density

$$S_0[\mathcal{H}] = \int 2\Lambda_{FLRW} = \lambda \int R_{FLRW}$$

$$R_Q = R(a, \dot{a}) = 6 (1 - q) H^2$$

$\lambda = \frac{1}{3}$ is uniquely fixed by horizon temperature $T_H = \frac{1}{2}(1 - q)T_{dS}$:

$$\Lambda_Q = (1 - q) H^2$$

van Putten, 2019, MNRAS, 491, L6;
2017, ApJ, 848, 28; 2015, MNRAS, 450, L48

Analytic solution

Friedman equations of three-flat FLRW with $\Lambda_Q = (1 - q) H^2$

$$H(z) = H_0 \frac{\sqrt{1 + 6\Omega_{M,0}(z + 2z^2 + 2z^3 + z^4 + \frac{1}{5}z^5)}}{1 + z}$$

$$w = \frac{2q - 1}{1 - q}, \quad q = 3\Omega_p$$

Van Putten, 2019, MNRAS, 491, L6;
2017, ApJ, 848, 28; 2015, MNRAS, 450, L48

Same parameters as late-time Λ CDM

$$H(z) = H_0 \sqrt{1 - \Omega_{M,0} + \Omega_{M,0}(1 + z)^3}$$

For the first time, can now compare apples with apples...

Cosmological evolution

Λ_Q -driven evolution is very similar to Λ CDM from RD to MD $\left(\frac{1}{2} < q < 1\right)$

Distinct evolution at late times, since dS is now unstable

N.B. Turning point in $H(z)$:

$$z_* = \left(\frac{5 - 6\Omega_{M,0}}{9\Omega_{M,0}} \right)^{1/5} - 1$$

O Colgain, van Putten,
Yavartanoo, 2019, PLB, 793 126

General scaling

$$\alpha = \frac{3 - 2\lambda}{\lambda} :$$

$$\frac{H(z)}{H_0} = h(\alpha z, \Omega_{M,0}/\alpha)$$

Results on H_0

- Fit Λ_Q to $H(z)$ compiled by Farooq et al. (2017) vs Λ CDM/CMB (Planck):

$$\frac{\Delta H_0}{H_0} = 10.6 \pm 3.6 \%$$

- Fit Λ_Q and Λ CDM to $H(z)$ compiled by Farooq et al., 2017) vs

$$\frac{\Delta H_0}{H_0} = 11.4 \pm 4.6 \%$$

based on

$$\Lambda_Q : H_0 = 74.9 \pm 2.6 \text{ km s}^{-1} \text{Mpc}^{-1}$$

$$\Lambda\text{CDM} : H_0 = 66.8 \pm 1.9 \text{ km s}^{-1} \text{Mpc}^{-1}$$

- H_0 from local distance ladder vs Λ CDM/CMB (Planck):

$$\frac{\Delta H_0}{H_0} = 9.38 \pm 2.2 \%$$

Riess et al., 2019, ApJ, 876, 85

<https://www.youtube.com/watch?v=bNdscedDCnQ>

van Putten, 2019, MNRAS, 491, L6
2017, ApJ, 848, 28

High-resolution low z

van Putten, 2020, MNRAS, 491, L6

$$\mu = 5 \log_{10} D_L + C_0 - 5 \log_{10} H_0, \quad D_L(z, \Omega_{M,0}) = (1+z) \int_0^z \frac{dz}{h(z, \Omega_{M,0})}$$

Results on $\Omega_{M,0}$

Fits to the Pantheon data,
1048 SNe by $[0, z_{max}]$ -
analysis

van Putten, 2019,
10.5281/zenodo.2597378
(March 18 2019)

Van Putten, 2020, MNRAS, 491, L6

Matter density more consistent ("flat") with Λ_Q than Λ CDM by a factor > 2

Low-high z approach?

- Λ CDM versus Λ_Q have distinct high- z asymptotic behavior:

$$H(z) \sim H_0 \sqrt{\Omega_{M,0}} (1+z)^{\frac{3}{2}}$$

$$H(z) \sim H_0 \sqrt{\frac{6}{5} \Omega_{M,0}} (1+z)^{\frac{3}{2}} \simeq 1.0954 \times H_0 \sqrt{\Omega_{M,0}} (1+z)^{\frac{3}{2}}$$

- Given present H_0 -tension of $\sim 10\%$, this tension is evolving according to

$$\frac{H(z)}{H(z)|_{\Lambda\mathbf{CDM}}} \simeq 1.2 \left(\frac{\Omega_{M,0}}{\Omega_{M,0,\Lambda\mathbf{CDM}}} \right)^{\frac{1}{2}} \simeq 1.34$$

May be observed to very high z , e.g., by GRBs with the planned ESA mission THESEUS.

- Expect a relatively high matter density:

$$\Omega_{M,0}[\Lambda_Q] \simeq 0.3707 \pm 0.015$$

van Putten, 2020, MNRAS, 491, L6

$$\Omega_{M,0}[\Lambda\mathbf{CDM}] = 0.2979 \pm 0.022$$

Planck Λ CDM/CMB

Cosmology by GRBs

FIG. 10.— Ω_M for a flat universe with $w = -1$. The best value and its asymmetric 1σ uncertainty are $0.39^{+0.12}_{-0.08}$.

Schaefer, B.E., 2007, ApJ, 660, 16

Conclusions

“It’s all about heat”

- We identify heat Q in cosmological spacetime as a relic of the Big Bang (breaking time-translation symmetry over a Hubble time at the onset to inflation), analogous to Q in black hole and Rindler spacetimes.
- Q is parameterized by the cosmological horizon with an equivalent energy density $\Lambda_Q = (1 - q)H^2$ of zero wave-length modes in the Lagrangian.
- A new analytic solution in (H_0, q_0) or, equivalently, $(H_0, \Omega_{M,0})$, recovers H_0 of the Local Distance Ladder from the $H(z)$ data of Farooq et al. (2017), reproducing the $\sim 10\%$ tension with Λ CDM.

H_0 -tension identifies Λ_Q : a cosmic thermal background of spacetime as a relic of the onset to inflation

EXTRA SLIDES

$j(z)$ -parameter

$$Q(z) = \frac{dq(z)}{dz}, \quad j(t) = \frac{\ddot{a}a^2}{\dot{a}^3}$$

Harrison, T., 1976, Nat., 260, 591

Sahni, V., Saini, T.D., Starobinski, A.A., Alam, U., 2003, JETP Lett., 77, 201;

Blandford, 2004, in Carnegie Obs. Astroph. Ser., 2 (Cambridge Univ. Press); 2005, in Observing Dark Energy, eds. S.C.Soff and T.R. Lauer, ASP Conf. Ser. 339

Shiba, T., Nakamura, T., 1998, PTP, 100, 1077

Mukherjee, A., & Banerjee, N., 2017, Class. Quant. Grav., 34 035016

Mamon, A.A., Bamba, K., 2018, EPJC, 78, 862

$$j(z) = q(z)(2q(z) + 1) + (1 + z) Q(z)$$

$$j(z) = 1 - (1 + z) h^{-2}(z) [h^2(z)]' + \frac{1}{2} (1 + z) [h^2(z)]''$$

(q_0, j_0) -parameters

Consider Λ CDM versus finite temperature Λ

$$\Lambda\mathbf{CDM} : h'(0) = \frac{3}{2}\Omega_{M,0}$$

$$\Lambda_Q : h'(0) = 3\Omega_{M,0} - 1$$

Example:

$$q_0 = -1 + h'(0) = \frac{3}{2}\Omega_{M,0} - 1 : \Omega_{M,0} = 0.3 \rightarrow q_0 = -0.55, \quad j_0 = 1$$

$$q_0 = -1 + h'(0) = 3\Omega_{M,0} - 2 : \Omega_{M,0} = 0.37 \rightarrow q_0 \rightarrow -0.89, \quad j_0 = 3.78$$