Journal Pre-proof

Initial quartz OSL and dust mass accumulation rate investigation of the Kisiljevo loess sequence in north-eastern Serbia

Zoran M. Perić, Slobodan B. Marković, Anca Avram, Alida Timar-Gabor, Christian Zeeden, Janina J. Nett, Peter Fischer, Kathryn E. Fitzsimmons, Milivoj B. Gavrilov

PII: S1040-6182(20)30665-0

DOI: <https://doi.org/10.1016/j.quaint.2020.10.040>

Reference: JQI 8585

To appear in: *Quaternary International*

Received Date: 26 July 2020

Revised Date: 3 October 2020

Accepted Date: 13 October 2020

Please cite this article as: Perić, Z.M., Marković, S.B., Avram, A., Timar-Gabor, A., Zeeden, C., Nett, J.J., Fischer, P., Fitzsimmons, K.E., Gavrilov, M.B., Initial quartz OSL and dust mass accumulation rate investigation of the Kisiljevo loess sequence in north-eastern Serbia, *Quaternary International* (2020), doi: <https://doi.org/10.1016/j.quaint.2020.10.040>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier Ltd.

Initial quartz OSL and dust mass accumulation rate investigation of the Kisiljevo loess sequence in north-eastern Serbia

Zoran M. Perić^{a, b*}, Slobodan B. Marković^b, Anca Avram^{c, d}, Alida Timar-Gabor^{c, d},
Christian Zeeden^e, Janina J. Nett^f, Peter Fischer^g, Kathryn E. Fitzsimmons^a, Milivoj B.
Gavrilov^b

^a Research Group for Terrestrial Paleoclimates, Max Planck Institute for Chemistry, Hahn-Meitner Weg 1, 55128 Mainz, Germany

^b Chair of Physical Geography, Faculty of Science and Mathematics, University of Novi Sad, Trg Dositeja Obradovića 3, 21000 Novi Sad, Serbia

^c Interdisciplinary Research Institute on Bio-Nano-Sciences, Babeş-Bolyai University, Treboniu Laurian 42, 400271 Cluj-Napoca, Romania

^d Faculty of Environmental Science and Engineering, Babeş-Bolyai University, Fantanele 30, 400327 Cluj-Napoca, Romania

^e Leibniz Institute for Applied Geophysics, Stilleweg 2, Hannover 30655, Germany

^f Department of Geography, Chair of Physical Geography and Geo-Ecology, RWTH Aachen University, Willnerstr. 5b, 52062 Aachen, Germany

^g Institute of Geography, Johannes Gutenberg University Mainz, Johann-Joachim-Becher-Weg 21, 55099 Mainz, Germany

*Corresponding author: Research Group for Terrestrial Paleoclimates, Max Planck Institute for Chemistry, Hahn-Meitner Weg 1, 55128 Mainz, Germany.

E-mail address: zoranperic.geo@gmail.com (Z.M. Perić)

Abstract

The thick and apparently continuous loess-palaeosol sequences in the Vojvodina region of northern Serbia are recognized and well understood as some of the oldest and most complete terrestrial European palaeoclimatic archives. By contrast, there are few published records for loess profiles from other regions in Serbia. Here we address this knowledge gap by investigating an 8 m thick loess sequence exposed near the village of Kisiljevo in north-eastern Serbia, describing the pedostratigraphy and environmental magnetic signatures in detail and placing these within a chronologic framework using quartz optically stimulated luminescence (OSL) on the 4-11 and 63-90 µm size fractions. Our results show dust accumulation over the last c. 32 ka, with substantial primary loess accumulation during the Early Holocene prior to the formation of the modern soil. We applied two age-depth

modelling approaches to estimate dust mass accumulation rates: the Bacon.r software and ADmin model. Both yield high accumulation rates, especially during MIS 2, averaging 550-600 g m⁻² a⁻¹ which exceed estimates for other investigated loess sequences in the region.

Key words: loess, dust mass accumulation rates, OSL dating, environmental magnetism, eastern Middle Danube Basin

1. Introduction

Loess-palaeosol sequences (LPS) are considered as some of the most important terrestrial archives of climatic change and past atmospheric dust activity. These sequences represent some of the oldest and most complete archives of environmental change on the European continent (Marković et al., 2012). The physical and geochemical characteristics of loess deposits yield valuable data about past environmental conditions and atmospheric circulation. The last decade has overseen the development of increasingly sophisticated and quantitative approaches for such reconstructions (e.g. Prud'homme et al., 2015; Obreht et al., 2019; Újvári et al., 2019), as well as improvements in geochronological control and age models providing the framework for correlations with global change (e.g. Zeeden et al., 2018; Timar-Gabor et al., 2015, 2017; Moine et al., 2017; Perić et al., 2019). However, significant geographical gaps in our understanding of terrestrial palaeoenvironmental change remain, in particular in regions of transition between dominant climatic regimes. Loess deposits often drape the land surface of such regions and provide enormous potential for insight into past climate dynamics.

The Middle Danube Basin, spanning Hungary, Croatia, Serbia and western Bulgaria and Romania, is one region that has overseen changes in the influence of different climatic systems (Obreht et al., 2016). In Serbia, loess sequences are thickest in the Vojvodina region (northern Serbia), where aeolian deposits blanket 95% of the land surface; deposits in the Vojvodina are the most intensively studied in the Middle Danube Basin (Marković et al., 2005, 2006, 2007, 2008, 2009, 2011). Loess in other parts of Serbia are localised and

discontinuous, overlying older slope deposits and river terraces, most commonly in the vicinity of the Danube river and its tributaries (Marković et al., 2014a; Perić et al., 2020). Aeolian deposits in north-eastern part of the country drape the westward slopes of the Carpathian-Balkan mountain chain and provide a useful insight into climate dynamics, but are the subject of only a few studies (e.g. Jovanović et al., 2006; Marković et al., 2014a; Perić et al., 2020). The loess in north-eastern Serbia is typical aeolian steppe sediment, containing rhizomorphs and pedogenic carbonates, and preserving crotonids, terrestrial gastropod and steppe rodent remains.

Here we investigate the Kisiljevo LPS, a typical loess deposit from north-eastern Serbia and the easternmost section so far discovered in the eastern Carpathian Basin. Its location on the western flanks of the Carpathian-Balkan mountains, ~90 km east of the Vojvodinan loess, presents a rare opportunity to understand regional climate evolution in a poorly investigated part of the Middle Danube Basin. We present a combination of pedostratigraphy and environmental magnetic analyses within a quartz optically stimulated luminescence (OSL) dating framework and age model for the exposed 8 m at Kisiljevo.

2. Regional setting

The Kisiljevo loess sequence is situated on the eastern edge of the village of the same name (44°44'0'' N and 21°25'0'' E), on the southern banks of the Silver Lake (Srebrno Jezero) and 29 km north-east of the city of Požarevac. The site lies at an altitude of 69 m above mean sea level (Fig. 1). The Silver Lake is an oxbow lake (former branch of the Danube) along its right riverbank and was anthropogenically disconnected from the Danube in 1971 by the construction of two dams (the Iron Gate) (Bertić et al., 2007). Previous fieldwork at the nearby village of Topolovnik (3 km south-east of Kisiljevo) showed that Quaternary sediments are widespread and overlie Pliocene deposits (Mitrović and Jovanović, 2000). Nevertheless, the loess of this area is not as thick as in the Vojvodina. Research conducted at

LPS at the nearby Nosak Drmno/Kostolac open-cut mine (Marković et al., 2014a; Perić et al., 2020) and Požarevac brickyard (Jovanović et al., 2006) suggest local predominance of grassland ecosystems during loess deposition over the Late Pleniglacial period.

Kisiljevo lies within hilly lowlands, bounded by low mountain ranges to the south. The profile extends east-west and is intersected by a road connecting the village Kisiljevo with the nearby settlements of Ram and Biskuplje, which divides the loess exposure in two parts. The thicker, northern profile is the subject of this study. It can be assumed that the Kisiljevo profile was more extensive in the past, but has been reduced to its present dimensions by fluvial erosion.

3. Methods

3.1 Sampling

Kisiljevo LPS was discovered during reconnaissance in May 2015. During June 2015, the upper 8 m profile was cleaned and sampled for environmental magnetism, colour description and luminescence dating. Samples for rock magnetic properties and colour determination were collected at 5 cm resolution. Four samples for luminescence dating – at depths of 390, 500, 700 and 790 cm - were obtained by hammering black PVC tubes (20 cm length, 5 cm diameter) into the freshly cleaned profile. The outer 3 cm of the material that had likely been exposed to sunlight was subsampled for analyses of radionuclide activity and water content. The inner, non-exposed sediment was used for OSL measurements.

3.2. Litho- and pedostratigraphic description

The profile was cleaned and described in detail (in situ observations) on a 8 m high vertical loess cliff (Fig. 2). Most previous stratigraphic studies of loess sections at various exposures in northern Serbia have used lithological, χ and pedogenic interpretations, along with variations in amino acid racemization geochronology, and magnetostratigraphy, usually combined with luminescence dating, as the primary basis for comparison (Marković et al.,

2005, 2006, 2007, 2008, 2009, 2011). The final result has been the establishment of a chronostratigraphy for the various loess-palaeosol units in Vojvodina. The nomenclature for this chronostratigraphy follows the Chinese loess stratigraphic system, with one exception - we used the prefix “V-” to refer to the standard Pleistocene loess-palaeosol stratigraphy in Vojvodina (Marković et al., 2008). Recently, a Danubian loess stratigraphic model (Marković et al., 2015) has been developed to correlate the Danubian loess-palaeosol units with the Chinese stratotype sections of Louchuan and Xiefeng. Nonetheless, in our paper the use of the Chinese “L” and “S” labels are applied similarly (e.g. Kukla, 1987; Kukla and An, 1989) without using any regional prefix. The profile description here follows this Danubian loess stratigraphy model by Marković et al. (2015).

3.3. Environmental magnetic properties and colour description

Samples for initial low-field magnetic susceptibility were ground in an agate mortar, compressed into plastic boxes and fixed with cotton wool in order to prevent movement of the sediment during the measurement. Individual sample density was used as a normalisation factor. Initial low-field magnetic susceptibility (χ , in $10^{-8} \text{m}^{-3} \text{kg}^{-1}$) was measured using a portable Bartington MS2 susceptibility meter at the Department of Geography, University of Novi Sad. Sediment colour was determined on uniformly dried samples in the laboratory using the Munsell Soil Colour Chart (1994).

3.4. Luminescence dating

3.4.1. Sample preparation

OSL dating sample preparation was carried out under subdued red light conditions in the luminescence laboratory in Cluj-Napoca, Romania. The ends of the sample tubes were subsampled and the wet and dry masses weighed for water content determination. The material from the inner part was used for both fine (4-11 μm) and coarse (63-90 μm) quartz extraction.

All samples were treated with hydrochloric acid (10% concentration) and hydrogen peroxide (30% concentration) in order to remove calcium carbonate and organic matter, respectively. The finer ($<63\ \mu\text{m}$) and coarser ($>63\ \mu\text{m}$) material was obtained through wet sieving. The fine (4-11 μm) quartz fractions were obtained by settling using Stoke's law, followed by etching in hydrofluorosilicic acid (35%) for 10 days centrifugation with distilled water. The 63-90 μm fraction was obtained by dry sieving. The 63-90 μm coarse quartz was separated from the polymineral mixture using an LST heavy liquid solution (density 2.62 and 2.75 g cm^{-3}). A treatment with 49% hydrofluoric acid digestion for 40 minutes was performed in order to isolate the quartz extracts from the plagioclase feldspars followed by a rinse with hydrochloric acid (10%) for 60 minutes in order to remove fluoride precipitates. The fine quartz grains were settled on aluminium disks with acetone ($\sim 2\ \text{mg ml}^{-1}$) using the pipette method while the coarse grained aliquots were mounted onto stainless steel disks using silicon spray.

3.4.2. Equivalent dose determination

Luminescence measurements were performed on an Risø TL/OSL DA-20 reader equipped with EMI 9235QA photomultiplier detection and blue-green LED stimulation head (Lapp et al., 2015). Quartz signals were detected using a 7.5 mm thick Hoya U-340 UV filter. A calibrated ^{90}Sr - ^{90}Y beta source was used for laboratory irradiation, calibrated for both fine and coarse grain fractions using gamma-irradiated calibration quartz (Hansen et al., 2015). The dose rate delivered for fine grains was 0.1161 Gy s and 0.1395 Gy s for coarse grains.

Equivalent doses were measured using the Single Aliquot Regenerative Dose (SAR) protocol (Wintle and Murray, 2000, 2003) on both size fractions using preheat and cutheat temperatures of 220°C and 180°C respectively. Optical stimulation was carried out for 40 s at 125 °C using blue LEDs. The signal of interest used for luminescence analysis was integrated over the first 0.308 s of the decay curve, and an early background (1.69 - 2.30 s interval) was subtracted from the net signal. The correction for sensitivity changes was made by using the

OSL signal to a test dose. For all measurements a test dose of 17 Gy was used. The temperature employed for the preheat was 220°C for 10 s. A cut-heat (ramp heating) to 180°C was employed before the measurement of the signal induced by the test dose. At the end of each SAR cycle, a high temperature bleach of 280°C was performed for 40 s (Murray and Wintle, 2003). The intrinsic tests (recycling and recuperation) of the SAR protocol (Murray and Wintle, 2003) were included in each measurement. The purity of the luminescence signal yielded by the quartz grains was investigated through the IR depletion test (Duller, 2003). A 10% deviation from unity was accepted for the recycling and IR depletion tests. For the recuperation signal, a value of 2% of the natural signal was considered acceptable. The aliquots used for the equivalent dose determination passed all the intrinsic tests. The results of the tests are presented in Table 1.

3.4.3. Dosimetry

The beta and gamma components of the dose rates were calculated from the activities of radionuclides K, Th and U, measured using high-resolution gamma spectrometry and converted to annual dose rates according to grain size using published conversion factors (Guérin et al. (2011)). A water content of 15% with a relative error of 25% was used. An α efficiency (α -value) of 0.04 ± 0.02 was considered for the fine (4-11 μm) quartz fraction (Rees-Jones, 1995). The beta attenuation and etching factor for the 63-90 μm quartz fraction was assumed to be 0.94 ± 0.05 (Mejdahl, 1979). The contribution of cosmic radiation was taken into account and calculated according to Prescott and Hutton (1994).

3.5. Dust flux estimation

The small number of OSL ages does not allow the construction of a sufficiently precise age-depth model for millennial-scale changes in the sedimentation rate (SR). However, the dataset is adequate for constraining the age and general trend of dust accumulation during MIS 2, for comparison with other loess sites in the region.

There are a number methods for developing continuous age-depth models from luminescence chronologies (e.g. Kang et al., 2015; Stevens et al., 2016; Újvári et al., 2014; Zeeden et al., 2018) and most of them use different approaches. In order to calculate reliable sedimentation rates and assess possible inconsistencies between different approaches, we generated two age-depth models, each based on contrasting assumptions: 1) the Bacon age-depth modelling software version 2.4.2 (Blaauw and Christen 2011) and 2) the ADmin model (Zeeden et al., 2018).

For the reconstruction of accumulation rates for deposits, Bacon uses Bayesian statistics, by combining numerical age dates with prior information such as the accumulation rate, its variability (and its memory change with depth). Accumulation rates are controlled using a gamma autoregressive semiparametric model with any defined number of subdivisions along the sediment profile (Blaauw and Christen, 2011). Prior knowledge about accumulation rates can lead to reduced uncertainty and more realistic modelling (Blaauw and Christen, 2011). The priors for accumulation rate and memory (autocorrelation) are modelled by gamma and beta distributions, and the latter defines how much the accumulation rate at a given depth depends on the depth above it (memory effect). The section thickness dictates the flexibility of the age-depth model to some degree, which is obtained by Markov Chain Monte Carlo (MCMC) iterations (Blaauw and Christen, 2011). Here, the resolution was set to 5 cm and the thickness to 40 cm. Because there was an observable discordance in the profile stratigraphy (where the tephra layer is assumed) a boundary was set at 700 cm depth.

Contrary to the Bacon code, the ADmin model by Zeeden et al. (2018) was designed specifically for the application on luminescence ages and does not make any assumptions on the sedimentation rates. It assumes that the real ages are stratigraphically consistent, and that seeming age inversions are the result of random uncertainty of the ages and associated uncertainty. For the application of the model on the Kisiljevo section, a density function for

random and systematic uncertainty from the data set was created with the assumption that both ages and uncertainties are correct.

For an estimation of the past atmospheric dust flux as well as for comparisons of dust archives between different regions, it is necessary to calculate the Mass Accumulation Rate (MAR; (Kohfeld and Harrison, 2003)). The MARs were calculated using the following equation:

$$\text{MAR} = \text{SR} \times \text{BD} \times f_{\text{eol}} \quad (1)$$

where SR is the sedimentation rate (m a^{-1}), BD is the dry bulk density (g cm^{-3}) and f_{eol} is the fraction of the sediment that is of aeolian origin (Kohfeld and Harrison, 2001). Here we assume that the origin of loess is entirely aeolian.

The BD values for loess in Europe vary in literature ranging from 1.65 g cm^{-3} (Frechen et al., 2003) to $1.497 \pm 0.079 \text{ g cm}^{-3}$ (Papp, 2009). Recently, Perić et al. (2020) measured the bulk density of the nearby Nosak loess-palaeosol sequence using the volumetric cylinder method and arrived at a mean value of $1.49 \pm 0.01 \text{ g cm}^{-3}$. We therefore used this BD value for our calculations.

4. Results

4.1. Litho- and pedo-stratigraphy

In the area surrounding Kisiljevo, the Holocene soil cover is a typical chernozem. This recent soil (S0) is 80 cm thick at our site, comprising a Ck horizon with plentiful CaCO_3 nodules (1-5 cm diameter), numerous crotovinas and root channels with humic material. The transitional AC horizon (10YR 5/1-3/3) is 20 cm thick, porous, with a silty loam texture and granular structure. The silty loam Ah horizon (10YR 6/3-4/4) is 40 to 60 cm thick and has a granular structure, along with some carbonate pseudomycelia. The underlying loess unit (L1) is 715 cm thick, light yellow to grey (5Y 7/3 5/3), coarse, porous and massive, without any visual

evidence of soil formation. Many spherical, soft carbonate nodules and humic infiltrations in old root channels are observed at the contact with the Holocene soil (S0). At a depth of ~7 m, a distinctively different layer, possibly a tephra, of approximately 50 cm thickness is interbedded with the loess. This layer is densely intersected with insect channels and fresh, recent cavities. The colour information is presented in Table S1.

4.2. Environmental magnetic record

Absolute χ values range between 18 and 42 x 10⁻⁸ m⁻³ kg⁻¹ (Fig. 3; Table S1). The S0 soil complex yields higher χ values than the L1 loess, and displays a low amplitude of variation ranging from 36 to 42 x 10⁻⁸ m⁻³ kg⁻¹. The L1 loess unit generally is more variable and ranges from 18 to 37 x 10⁻⁸ m⁻³ kg⁻¹ ($\bar{\chi}$ = 15 x 10⁻⁸ m⁻³ kg⁻¹). Two magnetic susceptibility peaks are observed at of ~5.5 and 7 m, with values increasing from 20 to 34 x 10⁻⁸ m⁻³ kg⁻¹ and from 24 to 37 x 10⁻⁸ m⁻³ kg⁻¹ respectively. This χ lf pattern usually suggests magnetic enhancements caused by pedogenesis (e.g. Maher and Thompson, 1992), although we observed no obvious soil development at these depths.

4.3. Luminescence properties

The quartz OSL signal is dominated by the fast component, as illustrated for both size fractions of sample ZAT 1/0.47 in Fig. 4 (inset). The rapid signal decay observed for both natural and regenerative signals from the Kisiljevo samples is similar to that of the calibration quartz, which is known to be dominated by the fast component (Hansen et al., 2015). The laboratory growth curves for sample ZAT 1/0.47 for the 4-11 and 63-90 μ m quartz fractions are presented in Fig. 4. The dose response curve was best fitted by a sum of two exponential functions in the form:

$$I(D) = I_0 + A_1 * (1 - \exp(-D/D_{01})) + A_2 * (1 - \exp(-D/D_{02})) \quad (2)$$

where I is the intensity of the signal for a given dose D ; I_0 is intercept; A_1 and A_2 represent the saturation amplitudes of the two exponential components; D_{01} and D_{02} are doses representing the onset of saturation of each exponential function.

The response to a zero dose (recuperation) is less than 0.3% from the corrected natural signal. The correction for sensitivity changes occurred throughout SAR measurement cycles are accurately performed (recycling ratio is close to unity).

4.3.1. Preheat plateau

In order to investigate the dependency of the equivalent doses on the preheat temperature, five aliquot sets from sample ZAT1/0.47 for both fine and coarse quartz were measured. For each set of aliquots, a preheat treatment ranging from 200 to 280°C in combination with a cut-heat of 180°C were employed in order to determine the equivalent doses. Fig. 5 shows that the equivalent doses measured using these intervals of preheat temperatures do not display a systematic variation. For equivalent dose calculation, only the aliquots for which recycling and recuperation ratios were within 10% deviation from unity were taken into account.

4.3.2. Dose recovery test

Further analysis focused on testing the applicability of the SAR protocol on fine grain (4-11 μm) and coarse grain (63-90 μm) quartz by applying dose recovery test. Five fresh aliquots of fine quartz from samples ZAT 2/0.395 and ZAT 1/0.47 and five aliquots of coarse (63-90 μm) quartz from sample ZAT 1/0.47 were bleached twice using the blue LEDs for 250 s at room temperature. Between the two bleaching treatments, a pause of 10 ks was used. The aliquots were then irradiated with a known dose chosen to be approximately equal with the estimated equivalent dose and then measured using the SAR protocol. Fig. 6 shows the given dose plotted against the recovered dose. In the case of fine quartz the ratios between the measured and the given dose for sample ZAT 2/0.395 and ZAT 1/0.47 are 1.00 ± 0.02 and 1.01 ± 0.03 , respectively. For coarse quartz from sample ZAT 1/0.47, a ratio of 1.02 ± 0.03 was obtained.

The results from the dose recovery test indicate that the SAR protocol is suitable for measuring laboratory doses given prior to any heat treatment for both fine and coarse grain quartz fractions.

4.3.3. Equivalent doses

Equivalent doses obtained by applying SAR protocol on fine and coarse quartz fractions are shown in Table 2. The equivalent doses for the fine quartz are ranging between 22 ± 3 (sample ZAT 2/0.395) and 97 ± 3 Gy (sample ZAT 1/0.41). In the case of coarse quartz, for the 63-90 μm fraction the measured equivalent doses range from 33 ± 1 (sample ZAT 2/0.395) to 90 ± 7 Gy (sample ZAT 1/0.41).

4.3.4. Dosimetry

The observed radionuclide concentrations as well as the annual doses for both fine and coarse quartz fractions are presented in Table 2. There is very little variation in the concentration of ^{232}Th , ^{226}Ra and ^{40}K with depth. For the 4-11 μm fine grain fraction the total dose rates range from 3.2 ± 0.15 (sample ZAT 2/0.395) to 3.6 ± 0.17 Gy ka^{-1} (sample ZAT 1/0.23); 2.7 ± 0.14 (sample ZAT 2/0.395) to 3.0 ± 0.16 Gy ka^{-1} (sample ZAT 1/0.23) for the 63-90 μm fraction. The calculated total doses rates show no observable trend with depth.

4.3.5. OSL ages

The calculated ages are presented in Table 2 and Fig. 7 as a function of depth. The quartz ages range from 27 ± 3 to 7 ± 1 ka for 4-11 μm fraction and 32 ± 3 to 12 ± 1 ka for the 63-90 μm coarse grains. The 63-90 μm ages display a steady increase with depth with no jumps or inversions. However, the 4-11 μm quartz shows an inversion at 760 cm depth of 1 ka although stratigraphically consistent within uncertainty. From Table 2 it is also noticeable that the 4-11 μm fraction displays slightly lower ages when compared to the 63-90 μm quartz. The observed age underestimation varies from 13% for sample ZAT 1/0.41 to 42% for the uppermost sample (ZAT 2/0.395). While the 13% underestimation at about 30 ka is to be

expected based on our previous observations on other sites (Constantin et al., 2015; Timar-Gabor et al., 2017) we currently have no explanation for the significant underestimation in the case of the age obtained using fine grains for sample ZAT 2/0.395, as usually a good agreement is reported between fine and coarse ages for that particular age range (Constantin et al., 2019; Tecsá et al., 2020). In the study of Timar-Gabor et al. (2015), similar age underestimation of the 4-11 μm quartz was observed for the Orlovat LPS in northern Serbia for ages older than ~ 30 ka. The discrepancy was later investigated for sites around the world and ascribed to a dose dependent phenomena, given that generally for samples with equivalent doses lower than about 100 Gy coarse and fine grain ages agree (see Timar-Gabor et al. 2017). At Kisiljevo the dependency is less pronounced, probably because the samples investigated are not old enough, with equivalent doses lower than 100 Gy. Consequently, until further measurements are conducted, we do not reject the 4-11 μm chronology for the Kisiljevo site. However, based on the overall better accuracy of the 63-90 μm quartz ages (Timar-Gabor et al., 2017), the age depth modelling was carried out using the chronology obtained on this fraction.

4.4. Mass accumulation rates

The modelling results of the Bacon and ADmin age-depth modelling methods are presented in Fig.8A and Fig. 8B respectively. It can be seen that both dataset display no inversions of the mean ages which is why the age-depth models were generated with only few resampling attempts and yielded stratigraphically consistent data.

The Bacon age-depth model is slightly less sensitive to changes in luminescence age with depth than the ADmin method and, to some degree, resulted in a more linear age-depth function. However both models indicate variable sedimentation rates (SRs), at least within the uncertainty limits of the techniques.

The calculated SRs and MARs based on the OSL ages are shown in Table S1. The MAR values in this study are presented without uncertainty estimates, mainly because there is no agreement on how to calculate accurate estimates of dust MAR uncertainty from loess deposits. Apart from potential uncertainties in bulk density estimates and the impact of post-depositional modification, simple Gaussian error propagation from luminescence ages might not be valid as uncertainty may not be strictly Gaussian, or fully random. However, it is assumed that the error on MAR estimates is higher than the luminescence age uncertainty in percentage terms (Perić et al., 2019).

The SR values calculated according to the Bacon age-depth model results vary from 275 to 962 mm ka⁻¹ with a median ($\tilde{x}$) and mean ($\bar{x}$) of 368 and 403 mm ka⁻¹, respectively. The MARs range from 409 to 1433 g m⁻² a⁻¹ peaking around 25 ka ($\tilde{x} = 548$ g m⁻² a⁻¹; $\bar{x} = 600$ g m⁻² a⁻¹). A MAR decrease was observed around ~20 ka where values of 438 g m⁻² a⁻¹ are recorded (Fig 9B). However, the decline is not as rapid as indicated by the ADmin model (Fig. 9A) and displayed a more gradual decrease which is most probably the result of the applied bacon.r code which assumes linear sedimentation over time.

According to the ADmin modelling results, the SRs at the Kisiljevo site range from 74 to 651 mm ka⁻¹ ($\tilde{x} = 385$ mm ka⁻¹; $\bar{x} = 373$ mm ka⁻¹). The resulting MARs (Fig. 9A) display highly variable values during the investigated period ranging from 110 to 970 g m⁻² a⁻¹ ($\tilde{x} = 573$ g m⁻² a⁻¹; $\bar{x} = 557$ g m⁻² a⁻¹). The highest MARs were observed at the onset of MIS 2 (~31 – 27 ka) with an average value of ~875 g m⁻² a⁻¹. A rapid decrease is observed at the transition between MIS 2 and MIS 1 (~12 ka) where the MAR values drop to 110 g m⁻² a⁻¹. The weighed mean for the Last Glacial period (~12 – 27 ka) is ~620 g m⁻² a⁻¹. Extremely high MARs were also recorded near the end of MIS 3 (31 - 32 ka) averaging at ~668 g m⁻² a⁻¹.

5. Discussion

5.1. Mass accumulation rates in loess deposits on the western Carpathian-Balkan mountain piedmonts

Magnetic susceptibility and final OSL ages are shown in Fig. 3; all calculated ages are presented in in Fig. 7 as a function of depth. The OSL ages show a steady increase with depth within uncertainty limits. The chronology suggests accumulation of the entire loess sequence from late MIS 3 to the present.

MARs from both age model outputs are generally very similar, with average values of c. 600 and 557 g m⁻² a⁻¹ for Bacon and ADmin respectively. Sedimentation varies between the models in the lower part of the section; the Bacon model produced estimates of 548 (25 ka) - 1406 g m⁻² a⁻¹ (26 ka), whereas ADmin yielded 366 (~27 ka) - 970 g m⁻² a⁻¹ (~31 ka). Peak MARs occur earlier (2-5 ka) in the ADmin model (Fig 9B). Whilst this may be the result of the model approach itself, which relies solely on luminescence ages without taking into account surrounding age depth information, given that the uncertainties on MAR estimates usually exceed luminescence age errors in percentage terms, such short-term discrepancies should be negligible.

Both models indicate both high and variable MARs during the Last Glacial, which contrasts with LPS in the Vojvodina, where linear sedimentation is assumed (e.g. Stevens et al. 2011; Perić et al. 2019). This variability at Kisiljevo may reflect changes in sediment production at the source, wind direction, deactivation or activation of an additional sources (Fenn et al. 2020), and/or local variability in trapping and preservation conditions (palaeotopography; Sprafke et al 2018; Fenn et al 2020). It is also possible that the linear sedimentation of previous studies reflects the different methodologies for calculating accumulation rates and/or the number of samples.

Our youngest age at 3.9 m depth (12 ± 1 ka) suggests extremely high rates of dust accumulation subsequent to that time (during the Holocene), which is not observed elsewhere in Serbia with the exception of Orlovat (Marković et al., 2014b, Timar-Gabor et al., 2015) and

Surduk (Hatte et al., 2013). The Kisiljevo chronology also suggests high dust accumulation rates around ~25-31 ka, as shown by MAR values of $>745 \text{ g m}^{-2} \text{ a}^{-1}$.

Generally, the most important palaeoenvironmental characteristic of the investigated area is significant dust input during the MIS 2 stage and the Holocene. Comparable accumulation rates in the Middle Danube basin have been observed only at Surduk and Titel in Serbia (Újvári et al. 2010), which yielded mean MAR values for MIS 2 (12 – 28 ka) of 434 and $510 \text{ g m}^{-2} \text{ a}^{-1}$, respectively, and in Hungary at Süttő - 584; Mende - 761; Tápiósüly - 504; Dunaszekcső - 707 and Dunaújváros ($1238 \text{ g m}^{-2} \text{ a}^{-1}$) (Újvári et al., 2017). We propose that high dust deposition rates are primarily controlled by a site's proximity to dust source, combined with prevailing wind circulations and palaeotopography (e.g. Marković et al., 2018).

5.2. Enviromagnetic record

The two peaks in the MS record observed at the depths of ~5.5 m and ~7 m are consistent with the transition of MIS 3/MIS 2 and the final stage of MIS 3, respectively. MIS 3 in this region is usually associated with considerable soil formation in loess deposits (e.g. Marković et al., 2008). Therefore it is somewhat unexpected that we observed no visible pedogenesis at Kisiljevo for this time period. The absence of MIS3 pedogenic horizons may be explained by locally continuous dust deposition preventing significant alteration, although the MS peak at ~5.5 m may indicate weakly developed palaeosols.

The peak at ~7 m coincides with the depth where a tephra layer is assumed (Fig. 2; Fig. 3) and may explain the increase in MS. However, our OSL ages do not support this hypothesis, since there is no known tephra typical in this region of this age (e.g. Campanian Ignimbrite tephra dated 39-40 ka (De Vivo et al., 2001; Pyle et al., 2006; Veres et al., 2013)). It is possible that the Kisiljevo LPS preserves a tephra that has not yet been reported.

However, until further studies are conducted (e.g. geochemical analysis), we can neither confirm nor deny the existence of tephra at this site.

5.3. Dust Source

Most likely the majority of silty material is provided by the deposition onto the Danube alluvial plain prior its passage through the Iron Gate gorge (e.g. Buggle et al., 2008; Smalley et al., 2009; Marković et al., 2015). These fine alluvial silts were most probably transported by the frequent and strong south-easterly Košava wind (Fig. 1) (Gavrilov et al. (2018). Two winds are posited to have dominated in this region over the last glacial period: persistent south-easterly winds in the wider area of the Banatska Peščara, resulting in the massive accumulation of aeolian sand ~16 km north-west of Kisiljevo (Fig. 1), and north-westerly winds over Northern Serbia during the recent and last glacial period (Marković et al. 2008). Gavrilov et al. (2018) concluded that the Košava wind is responsible for the formation of sand dunes in Banatska Peščara and perhaps for much of the loess across northern Serbia, and strongly support our interpretation about the nature of loess deposition at Kisiljevo.

6. Conclusions

In this study, we investigate the Kisiljevo loess section at the eastern edge of the Middle Danube Basin. Our OSL measurements on both quartz size fractions yield ages within error, although the fine-grained quartz slightly underestimates the coarser ages.

The calculated ages suggest continuous loess accumulation since late MIS 3, and continued high dust input into the early Holocene. Consistently high accumulation rates are likely the result of the interplay between proximity to the Danube alluvial plain as a silt source and the site's exposure to the Košava wind, which is responsible for most of the loess deposition in north-eastern Serbia. Further investigations, such as higher resolution luminescence dating and geochemical analyses, are required to gain more detailed insight into

past environmental conditions in this transitional zone between the Carpathian-Balkan mountains and the Vojvodina region.

Acknowledgements: The authors thank Marko Perić and Miloš Ranković for their help during field work and sampling. We are also grateful to naturalearthdata.com for the use of the river distribution in Fig.1 and Bruno Boemke from RWTH Aachen University for the construction of the map. SBM thanks the Serbian Academy of Sciences and Arts for financial support (grant # F178). AA and ATG acknowledge funding from the European Research Council Starting Grant 678106 (INTERTRAP, Horizon 2020). JN acknowledges funding from the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) - Project Number 57444011 - SFB 806.

Author contributions: Sample collection: ZMP, SBM and MBG; gamma spectrometry measurements: AA and ATG; sample preparation: AA and ZMP; luminescence measurements: AA and ATG; age-depth modelling: ZMP and CZ; data analysis: ZMP, SBM, AA and ATG; data presentation: ZMP and SBM; original draft preparation: ZMP, SBM and MBG; review and editing: all authors.

References

- Aitken, J.M. and Alldred, C.J., 1972. The assessment of error limits in thermoluminescent dating. *Archaeometry* 14, 257-267.
- Bertić, I., Šehić, D., Šehić, D., 2007. Atlas Srbije [The Atlas of Serbia]. Monde Neuf, Belgrade, 31-34.
- Blaauw, M., Christen, J.A., 2011. Flexible palaeoclimate age-depth models using an autoregressive gamma process. *Bayesian Analysis* 6 (3), 457-474.
- Buggle, B., Glaser, B., Hambach, U., Marković, S.B., Gerasimenko, N., Glaser, I., Zöller, L. 2008. Geochemical characterization and provenance of South-East and East European loesses (Serbia, Romania, Ukraine). *Quaternary Science Reviews* 27, 1058-1075.
- Constantin D., Camenita A., Panaiotu C., Necula C., Codrea V., Timar-Gabor A., 2015. Fine and coarse-quartz SAR-OSL dating of Last Glacial loess in Southern Romania. *Quaternary International*, 357, 33-43.
- Constantin, D., Veres, D., Anechitei-Deacu, V., Groza, S.M., Begy, R., Kelemen, S., Buylaert, J.-P., Panaiotu, C., Hambach, U., Marković, S.B., Gerasimenko, N., Timar-Gabor, A., 2019. Luminescence age constraints on the Pleistocene-Holocene transition recorded in loess sequences across SE Europe—*Quaternary Geochronology*, 49, 71-77.

- 476 De Vivo, B., Rolandi, G., Gans, P.B., Calvert, A., Bohrsen, W.A., Spera, F.J., Belkin, H.E.,
477 2001. New constraints on the pyroclastic eruptive history of the Campanian volcanic
478 Plain (Italy). *Mineralogy and Petrology* 73, 47–65.
- 479 Duller, G.A.T., 2003. Distinguishing quartz and feldspars in single grain luminescence
480 measurements. *Radiation Measurements* 37, 161-165.
- 481 Ehlers, J., Gibbard, P.L., Hughes, P.D. (Eds) 2011: *Quaternary Glaciations - Extent and*
482 *Chronology*. Elsevier, Amsterdam.
- 483 Fenn, K., Durcan, J.A., Thomas, D.S.G., Banak, A., 2020. A 180 ka record of environmental
484 change at Erdut (Croatia): a new chronology for the loess–palaeosol sequence and its
485 implications for environmental interpretation. *Journal of quaternary science*, 1-12.
- 486 Frechen, M., Oches, E.A., Kohfeld, K.E., 2003: Loess in Europe - mass accumulation rates
487 during the Last Glacial Period. *Quaternary Science Reviews* 22, 1835-1857.
- 488 Gavrilov, M. B., Marković, S. B., Schaetzl, R. J., Tošić, I., Zeeden, C., Obreht, I., Sipos, G.,
489 Ruman, A., Putniković, S., Emunds, K., Perić, Z., Hambach, U., Lehmkuhl, F., 2018.
490 Prevailing surface winds in Northern Serbia in the recent and past time periods;
491 modern- and past dust deposition. *Aeolian Research* 31, 117–129.
- 492 Guérin, G., Mercier, N., Adamiec, G., 2011. Dose-rate conversion factors: update. *Ancient*
493 *TL* 29, 5-8.
- 494 Hansen, V., Murray, A., Buylaert, J-P., Yeo, E-Y., Thomsen, C., 2015. A new irradiated
495 quartz for beta source calibration. *Radiation Measurements* 81, 123-127.
- 496 Hatté, C., Gauthier, C., Rousseau, D.D., Antoine, P., Fuchs, M., Lacroix, F., Marković, S.B.,
497 Moine, O., Sima, A., 2013. Excursions to C4 vegetation recorded in the Upper
498 Pleistocene loess of Surduk (Northern Serbia): an organic isotope geochemistry study.
499 *Climate of the Past* 9, 1001–1014.
- 500 Kohfeld, K.E. and Harrison, S.P., 2001. DIRTMAP: the geological record of dust. *Earth-*
501 *Science Reviews* 54, 81-114.

- 502 Kohfeld, K.E. and Harrison, S.P., 2003. Glacial-interglacial changes in dust deposition on the
503 Chinese Loess Plateau. *Quaternary Science Reviews* 22, 1859-1878.
- 504 JAXA EORC, 2016. ALOS Global Digital Surface Model “ALOS World 3D - 30m”.
- 505 Jovanović, M., Marković, S. B., Gaudenyi, T., Oches, E. A., Hambach, U., Zöller, L.,
506 Machalett, B., 2006. “Warm” glacial climate during loess deposition recorded at
507 exposures of the Požarevac brickyard, NE Serbia. *Geophysical Research Abstracts* 8,
508 10599.
- 509 Kukla, G., 1987. Loess stratigraphy in Central China. *Quaternary Science Reviews* 6, 191-
510 219.
- 511 Kukla, G. and An, Z., 1989. Loess stratigraphy in Central China. *Palaeogeography,*
512 *Palaeoclimatology, Palaeoecology* 72, 203-225.
- 513 Lapp, T., Kook, M., Murray, A.S., Thomsen, K., Buylaert, J.P., Jain, M., 2015. A new
514 Luminescence Detection and Stimulation Head for the Risø TL/OSL reader. *Radiation*
515 *Measurements* 81, 178-184.
- 516 Lehmkuhl, F., Böskén, J., Hošek, J., Sprafke, T., Marković, S.B., Obrecht, I., Hambach, U.,
517 Sümegi, P., Thiemann, A., Steffens, S., Lindner, H., Veres, D., Zeeden, C., 2018.
518 Loess distribution and related Quaternary sediments in the Carpathian Basin. *Journal*
519 *of Maps* 14, 661–670.
- 520 Maher, B.A. and Thompson, R., 1992. Paleoclimatic significance of the mineral magnetic
521 record of the Chinese loess and palaeosols. *Quaternary Research* 37, 155-170.
- 522 Marković, S.B., McCoy, W.D., Oches, E.A., Savić, S., Gaudenyi, T., Jovanović, M., Stevens,
523 T., Walther, R., Ivanišević, P., Galić, Z., 2005. Paleoclimate record in the upper
524 Pleistocene loess-palaeosol sequence at Petrovaradin brickyard (Vojvodina, Serbia).
525 *Geologica Carpathica* 56, 545–552.

- 526 Marković, S.B., Oches, E., Sümegi, P., Jovanović, M., Gaudenyi, T., 2006. An introduction
527 to the middle and upper Pleistocene loess-palaeosol sequence at Ruma brickyard,
528 Vojvodina, Serbia. *Quaternary International* 149, 80–86.
- 529 Marković, S.B., Oches, E.A., McCoy, W.D., Frechen, M., Gaudenyi, T., 2007. Malacological
530 and sedimentological evidence for “warm” glacial climate from the Irig loess
531 sequence, Vojvodina, Serbia. *Geochemistry Geophysics Geosystems* 8, Q09008.
- 532 Marković, S.B., Bokhorst, M., Vandenberghe, J., Oches, E.A., Zöller, L., McCoy, W.D.,
533 Gaudenyi, T., Jovanović, M., Hambach, U., Machalett, B., 2008. Late Pleistocene
534 loess- paleosol sequences in the Vojvodina region, north Serbia. *Journal of*
535 *Quaternary Science* 23, 73-84.
- 536 Marković, S.B., Hambach, U., Catto, N., Jovanović, M., Buggle, B., Machalett, B., Zöller, L.,
537 Glaser, B., Frechen, M., 2009. The middle and late Pleistocene loess-palaeosol
538 sequences at Batajanica, Vojvodina, Serbia. *Quaternary International* 198, 255-266.
- 539 Marković, S. B., Hambach, U., Stevens, T., Kukla, G. J., Heller, F., McCoy, W. D., Oches,
540 W.A., Buggle, B., Zöller, L., 2011. The last million years recorded at the Stari
541 Slankamen (Northern Serbia) loess-palaeosol sequence: revised chronostratigraphy
542 and long-term environmental trends. *Quaternary Science Reviews* 30, 1142–1154.
- 543 Marković, S., Hambach, U., Stevens, T., Jovanović, M., O’Hara-Dhand, K., Basarin, B., Lu,
544 H., Smalley, I., Buggle, B., Zech, M., Svirčev, Z., Sümegi, P., Milojković, N., Zöller,
545 L., 2012. Loess in the Vojvodina region (northern Serbia): an essential link between
546 European and Asian Pleistocene environments. *Netherlands Journal of Geosciences*
547 91, 173-188.
- 548 Marković, S.B., Korać, M., Mrđić, N., Buylaert, J.P., Thiel, C., McLaren, S.J., Stevens, T.,
549 Tomić, N., Petić, N., Jovanović, M., Vasiljević, D.A., Sümegi, P., Gavrilov, M.B.,
550 Obreht, I., 2014a. Palaeoenvironment and geoconservation of mammoths from the

- 551 Nosak loess-palaeosol sequence (Drmno, Northeastern Serbia): Initial results and
 552 perspectives. *Quaternary International* 334-335, 30-39.
- 553 Marković, S. B., Timar-Gabor, A., Stevens, T., Hambach, U., Popov, D., Tomić, N., Obreht,
 554 I., Jovanović, M., Lehmkuhl, F., Kels, H., Marković, R., Gavrilov, M. B., 2014b.
 555 Environmental dynamics and luminescence chronology from the Orlovat loess–
 556 palaeosol sequence (Vojvodina, northern Serbia). *Journal of Quaternary Science* 29,
 557 189-199.
- 558 Marković, S.B., Stevens, T., Kukla, G.J., Hambach, U., Fitzsimmons, K.E., Gibbard, P.,
 559 Buggle, B., Zech, M., Guo, Z., Qingzhen, H., Haibin, W., Dhand, O.K., Smalley, I.J.,
 560 Újvári, G., Sümegei, P., Timar-Gabor, A., Veres, D., Sirocko, F., Vasiljević, D.A., Jary,
 561 Z., Svensson, A., Jović, V., Lehmkuhl, F., Kovács, J., Svirčev, Z., 2015. Danube loess
 562 stratigraphy - towards a pan-European loess stratigraphic model. *Earth-Science*
 563 *Reviews* 148, 228-258.
- 564 Mejdahl, V., 1979. Thermoluminescence dating: beta-dose attenuation in quartz grains.
 565 *Archaeometry* 21, 61-72.
- 566 Mitrović, B., and Jovanović, G., 2000. Quaternary malacofauna of Topolovnik and Golubac
 567 (north-eastern Serbia). *Geologica Carpathica* 51, 3-6.
- 568 Moine, O., Antoine, P., Hatté, C., Landais, A., Mathieu, J.C., Prud'homme, C., and Rousseau,
 569 D., 2017. The impact of Last Glacial climate variability in west-European loess
 570 revealed by radiocarbon dating of fossil earthworm granules. *Proceedings of the*
 571 *National Academy of Sciences of the United States of America* 114, 6209–6214.
- 572 Munsell Colour, 1994. *Munsell Soil Colour Charts*. Macbeth Division of Kollmorgen
 573 Instruments: New Windsor.
- 574 Murray, A.S. and Wintle, A.G., 2000. Luminescence dating using an improved single-aliquot
 575 regenerative-dose protocol. *Radiation Measurements* 32, 57-73.

- 576 Murray, A.S. and Wintle, A.G., 2003. The single aliquot regenerative dose protocol: potential
577 for improvements in reliability. *Radiation Measurements* 37, 377-381.
- 578 Obreht, I., Zeeden, C., Schulte, P., Hambach, U., Eckmeier, E., Timar-Gabor, A., Lehmkuhl,
579 F., 2015. Aeolian dynamics at the Orlovat loess–paleosol sequence, northern Serbia,
580 based on detailed textural and geochemical evidence. *Aeolian Research* 18, 69–81.
- 581 Obreht, I., Zeeden, C., Hambach, U., Veres, D., Marković, S.B, Böskén, J., Svirčev, Z.,
582 Bačević, N., Gavrilov, M.B., Lehmkuhl, F., 2016. Tracing the influence of
583 Mediterranean climate on Southeastern Europe during the past 350,000 years.
584 *Scientific Reports* 6, 36334
- 585 Obreht, I., Zeeden, C., Hambach, U., Veres, D., Markovic, S.B., Lehmkuhl, F., 2019. A
586 critical reevaluation of palaeoclimate proxy records from loess in the Carpathian
587 Basin. *Earth- Science Reviews* 190, 498-520.
- 588 Papp, G., 2009. Simultaneous determination of terrain correction and local average
589 topographic density. *Acta Geodaetica and Geophysica Hungarica* 44, 191-202.
- 590 Perić, Z., Lagerbäck Adophi, E., Buylaert, J.-P, Stevens, T., Újvári, G., Marković, S.B,
591 Hambach, U., Fischer, P., Zeeden, C., Schmidt, C., Schulte, P., Huayu, L.,
592 Shuangwen1, Y., Lehmkuhl, F., Obreht, I., Veres, D., Thiel, C., Frechen, M., Jain, M., Vött,
593 A., Zöller, L., Gavrilov, M.B., 2019. Quartz OSL dating of late Quaternary Chinese and
594 Serbian loess: a cross Eurasian comparison of dating results and mass accumulation
595 rates. *Quaternary International* 502, 30-44.
- 596 Perić, Z.M., Marković, S.B., Sipos, G., Gavrilov, M.B., Thiel, C., Zeeden, C., Murray, A.S.,
597 2020. A post-IR IRSL chronology and dust mass accumulation rates of the Nosak
598 loess- paleosol sequence in north-eastern Serbia. *Boreas*, DOI:10.1111/bor.12459.
- 599 Prescott, J. R. and Hutton, J. T., 1994. Cosmic ray contributions to dose rates for
600 luminescence and ESR dating: Large depths and long-term time variations. *Radiation*
601 *Measurements* 23, 497-500.

- 602 Prud'Homme, C., Antoine, P., Moine, O., Turpin, E., Huguenard, L., Robert, V., Degeai, J.-P.,
603 2015. Earthworm calcite granules: a new tracker of millennial-timescale
604 environmental changes in Last Glacial loess deposits. *Journal of Quaternary Science*
605 30, 529-536.
- 606 Pyle, D.M., Ricketts, G.D., Margarib, V., van Andel, T.H, Sinitsyn, A.A., Praslov, N.D.,
607 Lisitsyn, S., 2006. Quaternary wide dispersal and deposition of distal tephra during the
608 Pleistocene „Campanian Ignimbrite/Y5“ eruption, Italy. *Quaternary Science Reviews*
609 25, 2713-2728.
- 610 Rees-Jones, J., 1995. Optical dating of young sediments using fine-grain quartz. *Ancient TL*
611 13, 9-13.
- 612 Smalley, I.J., O'Hara-Dhand, K., Wint, J., Machalett, B., Jary, Z., Jefferson, I.F., 2009. Rivers
613 and loess: the significance of long river transportation in the complex event-sequence
614 approach to loess deposit formation. *Quaternary International* 198, 7-18.
- 615 Sprafke, T., Fitzsimmons, K.E., Grützner, C., Elliot, A., Marquer, L., Nigmatova S., 2018.
616 Reevaluation of Late Pleistocene loess profiles at Remizovka (Kazakhstan) indicates
617 the significance of topography in evaluating terrestrial paleoclimate records.
618 *Quaternary Research* 89 674-690
- 619 Stevens, T., Marković, S.B., Zech, M., Hambach, U., Sümeği, P., 2011. Dust deposition and
620 climate in the Carpathian Basin over an independently dated last glacial-interglacial
621 cycle. *Quaternary Science Reviews* 30, 662-681.
- 622 Stevens, T., Buylaert, J.P., Lu, H., Thiel, C., Murray, A., Frechen, M., Yi, S., Zeng, L., 2016.
623 Mass accumulation rate and monsoon records from Xifeng, Chinese Loess Plateau,
624 based on a luminescence age model. *Journal of Quaternary Science* 31, 391-405.
- 625 Tecsa, V., Mason, J.A., Johnson, W.C., Miao, X., Radu, S., Magdas, D.A., Veres, D.,
626 Marković, S.B., Timar-Gabor, A., 2020. Latest Pleistocene to Holocene loess in the
627 central Great Plains: Optically stimulated luminescence dating and multi-proxy

- 628 analysis of the Enders loess section (Nebraska, USA). *Quaternary Science Reviews*,
 629 229, 106-130.
- 630 Timar-Gabor, A., Constantin, D., Marković, S. B., Jain, M., 2015. Extending the area of
 631 investigation of fine versus coarse quartz optical ages from the Lower Danube to the
 632 Carpathian Basin. *Quaternary International* 388, 168–176.
- 633 Timar-Gabor, A., Buylaert, J-P., Guralnik, B., Trandafir-Antohei, O., Constantin, D.,
 634 Anechitei-Deacu, V., Jain, M., Murray, A.S., Porat, N., Hao, Q., Wintle, A.G., 2017.
 635 On the importance of grain size in luminescence dating using quartz. *Radiation*
 636 *Measurements*, 106, 464-471.
- 637 Újvári, G., Kovács, J., Varga, G., Raucsik, B., Marković, S.B., 2010. Dust flux estimates for
 638 the Last Glacial Period in East Central Europe based on terrestrial records of loess
 639 deposits: a review. *Quaternary Science Reviews* 29, 3157-3166.
- 640 Újvári, G., Molnar, M., Novothny, A., Pall-Gergely, B., Kovács, J., Varhegyi, A., 2014. AMS
 641 ¹⁴C and OSL/IRSL dating of the Dunaszekcső loess sequence (Hungary): chronology
 642 for 20 to 150 ka and implications for establishing reliable age-depth models for the last
 643 40 ka. *Quaternary Science Reviews* 106, 140-154.
- 644 Újvári, G., Stevens, T., Molnar, M., Demény, A., Lambert, F., Varga, G., Jull, A.J.T., Páll-
 645 Gergely, B., Buylaert, J.P., Kovács, J., 2017. Coupled European and Greenland last
 646 glacial dust activity driven by North Atlantic climate. *Proceedings of the National*
 647 *Academy of Sciences of the United States of America* 114, E10622-E10631.
- 648 Újvári, G., Kele, S., Bernasconi, S.M., Haszpra, L., Novothny, Á., Bradák, B., 2019. Clumped
 649 isotope paleotemperatures from MIS 5 soil carbonates in southern Hungary.
 650 *Palaeogeography, Palaeoclimatology, Palaeoecology* 518, 72–81.
- 651 Vandenberghe, D., De Corte, F., Buylaert, J.-P., Kučera, J., Van den haute, P., 2008. On the
 652 internal radioactivity in quartz. *Radiation Measurements* 43, 771–775.

Veres, D., Lane, C.S., Timar-Gabor, A., Hambach, U., Constantin, D., Szakács, A., Fülling, A., Onac, B.P., 2013. The Campanian Ignimbrite/Y5 tephra layer – A regional stratigraphic marker for Isotope Stage 3 deposits in the Lower Danube region, Romania. *Quaternary International* 293, 22-33.

Zeeden, C., Dietze, M., Kreutzer, S., 2018. Discriminating luminescence age uncertainty composition for a robust Bayesian modelling. *Quaternary Geochronology* 43, 30-39. <https://doi.org/10.1016/j.quageo.2017.10.001>.

Table captions

Table 1. Summary of the measured equivalent doses along with the performance parameters of the SAR protocol for each grain-size of the 4 samples from the Kisiljevo section. *n* represents the number of aliquots averaged for D_e calculations. The errors represent 1 standard error.

Table 2. Summary of the OSL ages obtained on fine (4-11 μm) and coarse (63-90 μm) quartz and the radionuclide concentrations. The age uncertainties were determined following Aitken and Alldred (1972). The uncertainties associated with the luminescence and dosimetry data are random; the uncertainties mentioned with the optical ages are the overall uncertainties. The

systematic errors taken into account include: 2% beta source calibration, 3% conversion factors, 5% attenuation and etching factors, 3% gamma spectrometer calibration, 15% cosmic radiation, 25% water content. All uncertainties represent 1σ . Specific activities were measured on a planar detector by gamma spectrometry and the ages were determined considering 15% water content; beta attenuation and etching factors used for 63-90 μm quartz were 0.94 ± 0.050 whereas for 90-125 μm quartz were 0.92 ± 0.05 (Mejdahl, 1979); adopted alpha efficiency factor was 0.04 ± 0.02 for 4-11 μm quartz (Rees-Jones, 1995). The total dose rate consists of the contribution from the beta and gamma radiations for coarse grains as well as the contribution from alpha radiations in the case of fine grains. The contribution of cosmic radiation was taken into account and calculated according to Prescott and Hutton (1994). For coarse quartz grains an internal dose rate of $0.01 \pm 0.002 \text{ Gy ka}^{-1}$ was considered (Vandenbergh et al., 2008).

Figure captions

Fig. 1. Map of the loess distribution in north-eastern Serbia and the Vojvodina region showing the geographical position of the investigated section and other important loess sites. Also presented is the direction of the Košava wind. Distribution of loess, loess derivatives, aeolian sand and alluvium constructed according to Lehmkuhl et al. (2018). The LGM glacial extent and DEM (digital elevation model) shown according to Ehlers et al. (2011) and JAXA EORC (2016) respectively.

Fig. 2. The Kisiljevo loess-palaeosol sequence with the position of the luminescence samples and the supposed tephra layer. The cleaned and sampled surface is shaded in grey.

Fig. 3. Lithology of the Kisiljevo section, position of the luminescence samples (red circles) together with the preferred OSL ages and the magnetic susceptibility curve (for details see

text). Legend: Ah – Accumulated humus horizon; AC – Transitional horizon between initial pedogenesis and background sediment.

Fig 4. Representative sensitivity-corrected dose response curves constructed for sample ZAT 1/0.41 using one aliquot of (A) fine (4-11 μm) quartz grains and (B) coarse (63-90 μm) quartz grains. The sensitivity corrected natural signals are depicted as opened squares and dashed lines indicate the equivalent doses. The response to a zero dose is depicted as opened circles. Recycling and IR depletion points are represented as opened down triangles and opened up triangles, respectively. The insets show a typical decay curve of natural CW-OSL signal (open squares) in comparison to a regenerated signal (open circles) induced by a beta dose approximately equal with the equivalent dose and a calibration quartz signal (open triangles).

Fig. 5. Equivalent dose dependence on preheat temperatures for fine (4-11 μm) and coarse (63-90 μm) quartz fractions from sample ZAT 1/0.47. A cut-heat (test dose preheat) of 180°C was employed.

Fig. 6. Dose recovery test results for fine (closed circles) quartz from samples ZAT 2/0.395 and ZAT 1/0.47 and coarse (open circles) quartz from sample ZAT 1/0.47. The given irradiation doses were chosen to match the equivalent dose of each sample. The solid line indicates the ideal 1:1 dose recovery ratio while the dashed lines bracket a 10% variation from unity.

Fig. 7. OSL ages of the different grain size fractions used from the Kisiljevo loess site plotted against depth.

Fig. 8. Age modelling results and OSL ages from the Kisiljevo LPS. A) shows the results of the Bayesian modelling using the Bacon model of Blaauw and Christen (2011). The results are based on over 17 million Monte Carlo iterations and yield mean 95% confidence age ranges of 12790. B) shows the results of the ADmin age-depth modelling method by Zeeden

et al. (2018). The original data and uncertainty are plotted as closed circles with error bars: 1-sigma uncertainty as black line, and 2-sigma uncertainty as grey line.

Fig. 9. Dust Mass Accumulation Rate (MAR) as a function of age for the Kisiljevo LPS. A) MAR values obtained via the Bacon model and B) MAR values according to the model of Zeeden et al. (2018).

Table 1. Summary of the measured equivalent doses along with the performance parameters of the SAR protocol for each grain-size of the 4 samples from the Kisiljevo section. n represents the number of aliquots averaged for D_e calculations. The errors represent 1 standard error.

Sampling depth (cm)	Sample code	Grain size (μm)	n	D_e (Gy)	Recycling ratio	Recuperation (%)	IR depletion ratio
390	ZAT 2/0.395	4-11	13	22 ± 3	1.00 ± 0.001	0.12 ± 0.07	1.03 ± 0.02
		63-90	10	33 ± 1	0.96 ± 0.02	0.38 ± 0.12	0.97 ± 0.02
500	ZAT 1/0.23	4-11	7	84 ± 2	1.01 ± 0.002	-00.2 ± 0.09	0.96 ± 0.02
		63-90	8	81 ± 3	0.84 ± 0.13	0.10 ± 0.07	0.98 ± 0.03
700	ZAT 1/0.41	4-11	7	97 ± 3	0.99 ± 0.01	-00.7 ± 0.07	0.98 ± 0.02
		63-90	9	90 ± 7	0.97 ± 0.02	-0.04 ± 0.15	0.97 ± 0.02
790	ZAT 1/0.47	4-11	14	89 ± 2	0.96 ± 0.01	0.03 ± 0.08	0.98 ± 0.01
		63-90	18	90 ± 5	0.98 ± 0.01	-0.05 ± 0.58	0.97 ± 0.03

Table 2. Summary of the OSL ages obtained on fine (4-11 μm) and coarse (63-90 μm) quartz and the radionuclide concentrations. The age uncertainties were determined following Aitken and Alldred (1972). The uncertainties associated with the luminescence and dosimetry data are random; the uncertainties mentioned with the optical ages are the overall uncertainties. The systematic errors taken into account include: 2% beta source calibration, 3% conversion factors, 5% attenuation and etching factors, 3% gamma spectrometer calibration, 15% cosmic radiation, 25% water content. All uncertainties represent 1σ . Specific activities were measured on a planar detector by gamma spectrometry and the ages were determined considering 15% water content; beta attenuation and etching factors used for 63-90 μm quartz were 0.94 ± 0.050 whereas for 90-125 μm quartz were 0.92 ± 0.05 (Mejdahl, 1979); adopted alpha efficiency factor was 0.04 ± 0.02 for 4-11 μm quartz (Rees-Jones, 1995). The total dose rate consists of the contribution from the beta and gamma radiations for coarse grains as well as the contribution from alpha radiations in the case of fine grains. The contribution of cosmic radiation was taken into account and calculated according to Prescott and Hutton (1994). For coarse quartz grains an internal dose rate of $0.01 \pm 0.002 \text{ Gy ka}^{-1}$ was considered (Vandenberghe et al., 2008).

Sampling depth (cm)	Lab code	Grain size (μm)	^{232}Th (Bq kg^{-1})	^{226}Ra (Bq kg^{-1})	^{40}K (Bq kg^{-1})	Total random error (%)	Total systematic error (%)	Total dose rate (Gy ka^{-1})	Age (ka)
390	ZAT 2/0.395	4-11	39.1 ± 0.9	35.4 ± 0.6	481 ± 51	12.7	8.9	3.2 ± 0.15	7 ± 1
		63-90				5.7	6.9	2.7 ± 0.14	12 ± 1
500	ZAT 1/0.23	4-11	46.4 ± 0.7	41.1 ± 0.5	507 ± 57	5.3	9.2	3.6 ± 0.17	24 ± 3
		63-90				6.3	6.9	3.0 ± 0.16	27 ± 3
700	ZAT 1/0.41	4-11	45.9 ± 9.4	44.6 ± 0.6	484 ± 53	7.9	9.4	3.5 ± 0.25	27 ± 3
		63-90				10.3	6.9	2.9 ± 0.21	31 ± 4
790	ZAT 1/0.47	4-11	43.0 ± 0.5	39.6 ± 0.8	497 ± 54	5.4	9.1	3.4 ± 0.16	26 ± 3
		63-90				7.4	6.9	2.8 ± 0.15	32 ± 3

CONFLICT OF INTEREST

The authors declare no conflict of interest.

Sincerely

A handwritten signature in black ink, appearing to read 'Perić', followed by a long, horizontal, slightly wavy line that extends to the right.