Paleoceanography and Paleoclimatology

Supporting Information for

Biomarker and carbon isotopic evidence of marine incursions in the Himalayan Foreland Basin during its overfilled stage

Biswajit Roy¹, Sohom Roy¹, Kanva Goyal¹, Sambit Ghosh¹, Prasanta Sanyal¹

¹Indian Institute of Science Education and Research Kolkata, Mohanpur-741246, India

Content of this file

Figure S1 to S9

Table S1 to S6

Introduction

- The supporting files consist of nine figures and six tables.
- The supporting figures provide the process of estimation of ages, spectra of processed sample blanks and standards used. Some ternary and bivariate plots showing the OM sources and paleo-depositional conditions. Field photographs and some thin-section of the sandstone units are also provided as supporting files. Lastly, the lithostratigraphic correlation was done with other eastern foreland sections to identify the paleo-depositional condition of the HFB.
- The supporting tables provide an overview of the *n*-alkane and biomarker proxy ratios used to ascertain the organic sources and paleo-depositional conditions. A table has also been provided that show the biomarkers (hopane and steranes) that are identified in the Tista valley samples. Additionally, there are three consecutive tables that show temporal *n*-alkane, biomarker and $\delta^{13}\text{C}$ distribution in the sample.

Figure S1. Estimation of ages for the Tista valley sediments following similar lithological attributes of Muksar Khola (Nepal) east of the study area. In both sections, the beginning of thick amalgamated sand bodies characterising the Middle Siwaliks (which marks the upper boundary of the Lower Siwalik deposits) and the first appearance of the conglomerate beds characterising the commencement of the Upper Siwalik (the upper boundary of the Middle Siwalik deposits) are used as lower and upper tie-points, respectively, to constrain the age of the foreland sediments exposed in the Tista valley. The published magnetic polarity sequence (Ojha et al., 2009) ages was calibrated with the global magnetic polarity timescale (GPTS) of Gradstein et al. (2012). Note- Lithounits bearing leaf-macrofossils are marked against the litholog.

Figure S2. a) Representative GC-MS chromatograms of n-alkane external standard (C_8 - C_{40} SUPELCO) used to quantify Tista valley samples. GC-MRM representative chromatograms of : b) hopane (370→191) and c) sterane (372→217) sand blank processed along with the samples, d) 5α-Cholestane standard (Sigma-Aldrich).

Figure S3. a) Ternary plot showing relative abundance of short, mid and long-chain n -alkane concentrations; b) Plot of $\text{Pr}/n\text{-C}_{17}$ versus $\text{Ph}/n\text{-C}_{18}$ (fields marked after Peters et al., 2005), showing the organic matter sources and paleo-environment conditions of Tista valley sediments and Siwalik paleosols (Roy et al., 2020a)..

Figure S4. Covariance plot between a) $\delta^{13}\text{C}$ values of *n*-alkanes and CPI values; b) $\delta^{13}\text{C}$ values of *n*-alkanes and HMW *n*-alkane concentration; and c) Pr/Ph and gammacerane index.

Figure S5. Temporal variation in total, lower- and higher-molecular weight *n*-alkane abundance in the Tista valley sequence.

Figure S6. Stable carbon isotopic composition of *n*-alkanes (C_{15} - C_{33}) of the Tista valley samples, showing characteristic saw-tooth pattern in the HMW $\delta^{13}\text{C}$ values.

Figure S7. a) Ternary diagram showing the distribution of C_{27} , C_{28} and C_{29} regular steranes (measured on $m/z = 218$) and the paleo-depositional environment of the Tista valley samples (fields demarcated after Seifert & Moldowan, 1979). Relationships between b) $C_{27}/(C_{27} + C_{29})$ and Pr/Ph ratios, and c) C_{27}/C_{29} regular sterane and Pr/Ph ratios in Tista valley sediments. Vertical and horizontal lines inside the diagrams (b and c) represent boundaries separating depositional environments and source (terrigenous/marine) of organic matter (modified after Adegoke et al., 2014).

Figure S8. Thin section of a sandstone from lower part of the Tista valley sequence showing rounded framework grains and glauconite grains (marked with red arrows) in (a) plane and (b) cross polarised light. c) Bedding parallel preservation of wood blocks and d) scattered coal fragments within a sandstone unit of upper part of the section.

Table S1

Calculation of n-alkane and biomarker indices and ratios

CPI [*]	$0.5 \times \left[\frac{nC_{25} + nC_{27} + nC_{29} + nC_{31} + nC_{33}}{nC_{24} + nC_{26} + nC_{28} + nC_{30} + nC_{31}} + \frac{nC_{25} + nC_{27} + nC_{29} + nC_{31} + nC_{33}}{nC_{24} + nC_{26} + nC_{28} + nC_{30} + nC_{31}} \right]$
P _{aq} [#]	$\frac{nC_{23} + nC_{25}}{nC_{23} + nC_{25} + nC_{29} + nC_{31}}$
TAR ⁺	$\frac{nC_{27} + nC_{29} + nC_{31}}{nC_{15} + nC_{17} + nC_{19}}$
EOP [^]	$0.5 \times \left[\frac{nC_{16} + nC_{18}}{nC_{15} + nC_{17}} + \frac{nC_{16} + nC_{18}}{nC_{17} + nC_{19}} \right]$
Gammacerane index ^{**}	$\frac{C_{30} \text{ Gammacerane}}{C_{30} \text{ Gammacerane} + C_{30} \alpha\beta \text{ Hopane}} \times 10$
Oleanane index ^{##}	$\frac{C_{30} \text{ Oleanane}}{C_{30} \text{ Oleanane} + C_{30} \alpha\beta \text{ Hopane}}$
S/H Ratio ^{^^}	$\frac{(C_{27} - C_{29})\alpha\alpha\alpha + \alpha\beta\beta \text{ Steranes}}{(C_{29} - C_{32})\alpha\beta \text{ Hopanes}}$
Dia/RegSterane ⁺⁺	$\frac{(C_{27} - C_{29})\alpha\beta + \beta\alpha \text{ Diasteranes (S + R)}}{(C_{27} - C_{29})\alpha\alpha\alpha + \alpha\beta\beta \text{ Steranes}}$

*Bray & Evans, (1961); ** Peters & Moldowan, (1993); # Ficken et al. (2000); ## Dzou et al. (1999); + Bourbonniere & Meyers, 1996; ++McKirdy et al., 1983; ^ Cranwell et al., 1987; ^^ Moldowan et al., 1985

Table S2

List of biomarker compounds and their MRM-GC-MS precursor product relation

Biomarker compound list	Identifier	Precusor mass	Product mass
18 α (H)-22,29,30-Trisnorneohopane	Ts	370	191
17 α (H)-22,29,30-Trisnorhopane	Tm		
17 α ,21 β (H)-30-norhopane	C ₃₀ - norhopane	398	191
18 α (H)-30-norneohopane	C ₂₉ Ts		
17 α (H)-diahopane	C ₃₀ *	412	191
Oleanane	Oleanane		
17 α ,21 β (H)-hopane	C ₃₀ -Hopane		
17 β ,21 α (H)-hopane	C ₃₀ -moretane		
Gammacerane	Gammacerane		
17 α ,21 β (H)-29-homohopane 22S	C ₃₁ S-Hopane	426	191
17 α ,21 β (H)-29-homohopane 22R	C ₃₁ R-Hopane		
17 α ,21 β (H)-29-bishomohopane 22S	C ₃₂ S-Hopane	440	191
17 α ,21 β (H)-29-bishomohopane 22R	C ₃₂ R-Hopane		
13 β (H),17 α (H)-Diacholestane (20S)	C ₂₇ - Diasterane	372	217
13 β (H),17 α (H)-Diacholestane (20R)			
13 α (H),17 β (H)-Diacholestane (20S)			
13 α (H),17 β (H)-Diacholestane (20R)			
5 α (H),14 α (H),17 α (H)-Cholestane (20S)	C ₂₇ -Regular Sterane		
5 α (H),14 β (H),17 β (H)-Cholestane (20R)			
5 α (H),14 β (H),17 β (H)-Cholestane (20S)			
5 α (H),14 α (H),17 α (H)-Cholestane (20R)			
13 β (H),17 α (H)-Diaergostane (20S)	C ₂₈ - Diasterane	386	217
13 β (H),17 α (H)-Diaergostane (20R)			
13 α (H),17 β (H)-Diaergostane (20S)			
13 α (H),17 β (H)-Diaergostane (20R)			
5 α (H),14 α (H),17 α (H)-Ergostane (20S)	C ₂₈ -Regular Sterane		
5 α (H),14 β (H),17 β (H)-Ergostane (20R)			
5 α (H),14 β (H),17 β (H)-Ergostane (20S)			
5 α (H),14 α (H),17 α (H)-Ergostane (20R)			
13 β (H),17 α (H)-Diastigmastane (20S)	C ₂₉ - Diasterane	400	217
13 β (H),17 α (H)-Diastigmastane (20R)			
13 α (H),17 β (H)-Diastigmastane (20S)			
13 α (H),17 β (H)-Diastigmastane (20R)			
5 α (H),14 α (H),17 α (H)-Stigmastane (20S)	C ₂₉ -Regular Sterane		
5 α (H),14 β (H),17 β (H)-Stigmastane (20R)			

5 α (H),14 β (H),17 β (H)-Stigmastane (20S)			
5 α (H),14 α (H),17 α (H)-Stigmastane (20R)			
13 β (H),17 α (H)-C30 sterane (20S)			
13 β (H),17 α (H)-C30 sterane (20R)	C ₃₀ - Diasterane		
13 α (H),17 β (H)-C30 sterane (20S)			
13 α (H),17 β (H)-C30 sterane (20R)			
5 α (H),14 α (H),17 α (H)-C30 sterane (20S)		414	217
5 α (H),14 β (H),17 β (H)-C30 sterane (20R)	C ₃₀ -Regular Sterane		
5 α (H),14 β (H),17 β (H)-C30 sterane (20S)			
5 α (H),14 α (H),17 α (H)-C30 sterane (20R)			

Table S3 A: Variation in relative abundance (%) of *n*-alkanes in the Tista valley samples.

Sample	Strat. Height	15	16	17	Pr	18	Py	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	Short-chain (%)	Mid-chain (%)	Long-chain (%)	Short chain n-alkane conc. (ug/g)	Mid chain n-alkane conc. (ug/g)	Long chain n-alkane conc. (ug/g)	Total n-alkane conc. (ug/g)
SV 12	1400	1.6	7.3	2.9	7.2	12.0	1.9	0.9	0.9	3.0	0.2	3.7	2.7	4.2	2.9	4.8	2.7	5.9	1.4	9.0	4.3	6.7	1.0	13.0	25	10	56	0.23	0.10	0.52	0.94
SV 11	1125	2.1	9.4	3.2	3.4	4.8	2.7	3.7	7.9	5.5	2.4	3.3	0.4	5.0	4.9	5.7	5.3	5.8	3.9	10.0	5.2	5.4	0.0	0.0	23	19	51	0.02	0.02	0.05	0.10
SV10	972	19.4	22.5	0.1	0.0	3.6	1.2	15.7	4.6	6.3	0.3	0.8	1.7	2.7	3.1	3.3	3.2	2.8	2.5	2.4	1.5	1.4	0.8	0.0	61	14	24	0.06	0.01	0.02	0.10
SV 9	860	1.1	1.9	6.5	3.9	13.2	2.3	1.9	0.4	10.4	1.9	3.1	6.2	6.9	5.7	4.9	4.2	1.6	1.5	9.2	5.2	5.3	2.8	0.0	25	22	47	0.08	0.07	0.16	0.33
SV 8	710	3.0	4.9	4.2	11.6	5.6	4.4	3.4	2.9	3.1	5.2	4.1	3.3	5.2	5.2	6.8	4.9	5.6	0.9	7.9	2.8	5.1	0.0	0.0	21	19	44	0.04	0.04	0.09	0.20
SV 7	600	13.4	20.2	6.3	7.5	2.9	7.7	0.5	5.4	1.6	0.1	2.4	0.0	2.7	3.2	4.0	3.2	4.8	3.7	3.4	3.0	3.9	0.0	0.0	43	9	32	0.03	0.01	0.02	0.07
SV 6	500	0.7	11.5	1.5	1.9	9.4	5.1	6.6	1.5	17.0	2.1	0.9	0.8	1.6	2.6	3.4	5.0	5.4	6.3	5.1	5.3	6.2	0.0	0.0	30	22	41	0.18	0.14	0.25	0.62
SV 5	433	12.0	15.2	4.0	10.1	3.0	3.2	1.3	3.0	1.9	3.9	0.0	2.4	0.3	5.4	5.1	4.3	6.0	4.1	6.2	2.3	4.1	1.1	1.2	35	11	40	0.11	0.04	0.13	0.32
SV 4	302	22.6	0.2	15.5	6.6	35.0	5.8	1.5	0.5	0.1	0.3	0.4	0.4	0.8	0.5	0.9	0.7	1.2	1.2	1.8	1.5	2.4	0.0	0.0	75	2	11	0.09	0.00	0.01	0.12
SV 3	166	12.9	31.8	8.5	9.7	11.4	6.2	0.6	2.8	0.0	0.1	0.5	0.8	1.2	1.2	1.6	1.3	2.4	1.9	2.9	0.0	2.1	0.0	0.0	65	4	15	0.07	0.00	0.02	0.11
SV 2	73	3.1	4.5	4.8	6.1	4.1	2.6	0.0	0.0	3.8	0.0	0.2	5.4	6.1	6.3	7.7	6.2	14.4	5.5	8.9	3.1	4.4	1.4	1.2	17	9	65	0.07	0.04	0.27	0.42
SV 1	30	2.4	4.1	3.6	3.3	5.1	1.9	4.0	7.6	8.0	7.0	6.3	5.3	5.2	4.4	4.9	3.9	4.2	3.4	7.7	1.9	4.8	1.2	0.0	19	34	42	0.01	0.01	0.01	0.03

Table S3 B: Variation in concentration of individual *n*-alkanes (µg/g) in the Tista valley samples.

Sample	Strat. Height	15	16	17	Pr	18	Py	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35
SV 12	1400	0.015	0.069	0.027	0.067	0.113	0.018	0.008	0.008	0.029	0.002	0.035	0.025	0.039	0.027	0.045	0.025	0.056	0.013	0.085	0.040	0.063	0.009	0.122
SV 11	1125	0.002	0.009	0.003	0.003	0.005	0.003	0.004	0.008	0.006	0.002	0.003	0.000	0.005	0.005	0.006	0.005	0.006	0.004	0.010	0.005	0.005	0.000	0.000
SV10	972	0.019	0.022	0.000	0.000	0.004	0.001	0.016	0.005	0.006	0.000	0.001	0.002	0.003	0.003	0.003	0.003	0.003	0.002	0.002	0.002	0.001	0.001	0.000
SV 9	860	0.004	0.006	0.021	0.013	0.044	0.008	0.006	0.001	0.034	0.006	0.010	0.021	0.023	0.019	0.016	0.014	0.005	0.005	0.030	0.017	0.018	0.009	0.000
SV 8	710	0.006	0.010	0.008	0.023	0.011	0.009	0.007	0.006	0.006	0.010	0.008	0.007	0.010	0.010	0.014	0.010	0.011	0.002	0.016	0.006	0.010	0.000	0.000
SV 7	600	0.009	0.014	0.004	0.005	0.002	0.005	0.000	0.004	0.001	0.000	0.002	0.000	0.002	0.002	0.003	0.002	0.003	0.003	0.002	0.002	0.003	0.000	0.000
SV 6	500	0.004	0.071	0.009	0.012	0.058	0.032	0.041	0.010	0.105	0.013	0.006	0.005	0.010	0.016	0.021	0.031	0.033	0.039	0.032	0.033	0.039	0.000	0.000
SV 5	433	0.038	0.049	0.013	0.032	0.010	0.010	0.004	0.010	0.006	0.013	0.000	0.008	0.001	0.017	0.016	0.014	0.019	0.013	0.020	0.007	0.013	0.003	0.004
SV 4	302	0.027	0.000	0.019	0.008	0.042	0.007	0.002	0.001	0.000	0.000	0.000	0.000	0.001	0.001	0.001	0.001	0.001	0.001	0.002	0.002	0.003	0.000	0.000
SV 3	166	0.014	0.035	0.009	0.011	0.013	0.007	0.001	0.003	0.000	0.000	0.001	0.001	0.001	0.001	0.002	0.001	0.003	0.002	0.003	0.000	0.002	0.000	0.000
SV 2	73	0.013	0.019	0.020	0.026	0.017	0.011	0.000	0.000	0.016	0.000	0.001	0.023	0.025	0.026	0.032	0.026	0.061	0.023	0.038	0.013	0.019	0.006	0.005
SV 1	30	0.001	0.001	0.001	0.001	0.002	0.001	0.001	0.002	0.002	0.002	0.002	0.002	0.002	0.001	0.001	0.001	0.001	0.001	0.002	0.001	0.001	0.000	0.000

Table S4: Variation of proxy ratios and carbon isotopic composition of bulk and odd-HMW *n*-alkanes

Sample	Strat. Height	CPI	EOP	Paq	TAR	Pr/Ph	Pr/n C17	Ph/n C18	TOC (%)	$\delta^{13}\text{C}_{\text{bul k}}$ (‰)	stdv	$\delta^{13}\text{C}_{\text{n C27}}$ (‰)	stdv	$\delta^{13}\text{C}_{\text{n C29}}$ (‰)	stdv	$\delta^{13}\text{C}_{\text{n C31}}$ (‰)	stdv	$\delta^{13}\text{C}_{\text{n C33}}$ (‰)	stdv
SV 12	1400	2.3	4.7	0.3	3.6	3.9	2.5	0.2	0.8	-23.8	0.16	-31.3	0.6	-31.9	0.7	-29.7	0.1	-33.3	0
SV 11	1125	1.6	2.4	0.3	2.4	1.3	1.1	0.6	0.1	-23.6	0.07	-31.7	0.2	-32.7	0.4	-30.6	0.4	-35.5	0.1
SV10	972	1.1	1.5	0.4	0.2	0.0	0.3	0.3	0.05	-22.5	0.05	-30.9	0.1	-30.3	0.1	-31.3	0.3	-31.4	0.1
SV 9	860	1.3	1.9	0.5	1.7	1.7	0.6	0.2	0.16	-28.2	0.04	-30.1	0.4	-30.9	0.9	-30.6	0.7	-35.6	0.5
SV 8	710	2.0	1.4	0.4	1.9	2.6	2.7	0.8	0.04	-27.3	0.03	-31.4	0.6	-31.2	0.3	-39.4	0.5	-38.1	0.1
SV 7	600	1.4	2.3	0.4	0.6	1.0	1.2	2.7	0.04	-24.5	0.02	-34.2	0.5	-33.5	0.7	-32.5	0	-33.7	0.6
SV 6	500	1.1	6.1	0.2	1.6	0.4	1.3	0.5	0.28	-25.2	0.28	-32.8	0.5	-31.1	0.3	-32.1	0.1	-32.1	1
SV 5	433	1.2	2.3	0.0	1.0	3.1	2.5	1.1	0.1	-25.3	0.1	-31.1	0.3	-33.7	0.5	-33.8	0.3	-35.1	0.3
SV 4	302	1.7	1.5	0.3	0.1	1.1	0.4	0.2	0.09	-25.7	0.04	-33.5	0.2	-32.5	0.1	-35.4	0.2	-	-
SV 3	166	2.2	3.4	0.3	0.3	1.6	1.2	0.5	0.04	-24.6	0.04	-33.5	0.8	-32.3	0.4	-33.6	0.1	-33.6	1
SV 2	73	1.7	1.5	0.2	4.0	2.3	1.3	0.6	0.07	-26.1	0.07	-31.4	0.2	-33.9	0.3	-35.6	0.2	-35.2	0.1
SV 1	30	1.6	1.4	0.5	1.7	1.7	0.9	0.4	0.02	-25.1	0.02	-27.9	0.1	-29.7	1.2	-33.5	0.7	-31.6	0.7

$\mu\text{g/g}$ = micrograms of compound per gram of dry sediment; Pr: pristane; Ph: phytane

Table S5: Variation in $\delta^{13}\text{C}$ values among different homologues of *n*-alkanes in the Tista valley samples.

Sample	Strat. Height	Age (Ma)	15	16	17	Pr	18	Ph	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33
SV12	1400	3.3	-27.5	-27.2	-30.2	-29.5	-28.6	-31.9	-27.4	-27.5	-27.7	-27.0	-27.8	-28.8	-29.1	-28.6	-31.3	-30.7	-31.9	-29.5	-29.7	-28.4	-33.3
SV11	1125	5.0	-28.0	-28.4	-28.6	-	-29.8	-31.1	-31.8	-31.4	-30.7	-31.6	-30.5	-32.1	-31.8	-30.5	-31.7	-30.5	-32.7	-30.7	-30.6	-29.4	-35.5
SV10	972	6.0	-27.5	-28.0	-	-28.7	-29.9	-	-33.1		-31.3	-	-31.3	-30.7	-31.2	-30.9	-30.9	-28.9	-30.3	-29.9	-31.3	-30.8	-31.4
SV9	860	6.5	-26.8	-28.8	-27.4	-28.4	-33.1	-31.9	-30.9	-30.9	-30.0	-30.7	-29.8	-29.7	-30.2	-28.1	-30.1	-28.0	-30.9	-30.0	-30.6	-29.5	-35.6
SV8	710	7.3	-	-28.8	-32.3	-	-32.2	-33.2	-27.8	-31.6	-33.6	-31.6	-32.8	-33.5	-32.1	-30.9	-31.4	-29.4	-31.2	-30.7	-39.4	-35.5	-38.1
SV7	600	8.0	-28.0	-27.9	-27.8	-27.6	-28.8	-28.3	-28.5	-	-29.0	-29.6	-	-30.2	-35.0	-34.0	-34.2	-33.7	-33.5	-31.6	-32.5	-30.5	-33.7
SV6	500	8.5	-28.2	-28.3	-27.7	-	-29.3	-31.9	-	-28.7	-	-29.7	-	-31.2	-34.1	-32.7	-32.8	-30.9	-31.1	-30.0	-32.1	-31.8	-32.1
SV5	433	9.0	-29.6	-27.6	-30.2	-30.9	-33.8	-31.4	-31.6	-	-32.9	-30.7	-30.7	-30.2	-31.9	-30.8	-31.1	-29.9	-33.7	-30.5	-33.8	-32.7	-35.1
SV4	302	9.6	-26.6	-28.3	-28.1	-	-27.9	-34.4	-28.5	-28.2	-32.5	-38.3	-32.0	-31.0	-31.9	-30.0	-33.5	-30.5	-32.5	-29.9	-35.4	-33.8	-
SV3	166	10.1	-27.0	-27.8	-27.9	-30.4	-30.2	-34.5	-	-	-28.2	-29.0	-28.8	-28.1	-35.6	-34.0	-33.5	-30.6	-32.3	-31.7	-33.6	-31.9	-33.6
SV2	73	10.7	-30.9	-28.7	-28.6	-30.0	-37.1	-38.1	-29.5	-30.9	-32.1	-31.4	-31.8	-29.7	-31.2	-29.9	-31.4	-28.5	-33.9	-31.8	-35.6	-33.7	-35.2
SV1	30	11.0	-28.1	-27.8	-29.8	-29.0	-29.0	-31.1	-29.5	-28.8	-28.1	-27.8	-27.5	-27.4	-28.0	-27.0	-27.9	-25.5	-29.7	-27.0	-33.5	-31.6	-31.6

Pr: pristane; Ph: phytane

Table S6. Sample information, source organic matter and depositional environment parameters from saturated hydrocarbons of Tista valley sediments

Sample	Strat. Height	Age (Ma)	Hopane Ts/Tm	Hopane Ts/(Ts+Tm)	Hopane C ₂₉ /C ₃₀	Hopane C ₃₁ R/C ₃₀	Hopane C ₃₁ R-S/(S+R)	Gammacerane Index	Oleanane Index	Sterane Dia/Reg	Sterane C ₂₇ /C ₂₉	Sterane C ₂₇ /(C ₂₇ +C ₂₉)	S/H Ratio
SV 12	1400	3.3	1.96	0.66	0.47	0.15	0.51	0.10	0.17	0.94	0.12	0.10	1.26
SV 11	1125	5.0	1.54	0.61	1.40	0.36	0.35	0.36	0.09	0.42	0.12	0.11	1.08
SV 10	972	6.0	2.41	0.71	1.12	0.32	0.56	0.64	0.17	0.71	0.45	0.31	1.73
SV 9	860	6.5	0.71	0.41	0.73	0.33	0.38	0.14	0.13	0.48	0.30	0.23	0.85
SV 8	710	7.3	1.94	0.66	0.49	0.15	0.46	0.11	0.17	1.35	0.21	0.18	0.32
SV 7	600	8.0	2.77	0.73	0.68	0.30	0.54	0.72	0.13	0.64	0.31	0.24	0.61
SV 6	500	8.5	2.92	0.75	1.30	0.40	0.53	0.43	0.16	1.18	0.61	0.38	0.94
SV 5	433	9.0	2.25	0.69	1.04	0.26	0.57	0.17	0.18	0.80	0.39	0.28	0.60
SV 4	302	9.6	1.67	0.63	1.33	0.33	0.52	0.45	0.14	1.47	0.95	0.49	1.15
SV 3	166	10.1	2.81	0.74	1.68	0.26	0.52	0.21	0.03	1.84	1.85	0.65	1.10
SV 2	73	10.7	2.74	0.73	1.06	0.21	0.54	0.43	0.06	0.77	0.27	0.21	0.58
SV 1	30	11.0	5.15	0.84	1.17	0.25	0.47	0.34	0.08	0.61	0.27	0.21	0.92

References

- Bourbonniere, R. A., & Meyers, P. A. (1996). Sedimentary geolipid records of historical changes in the watersheds and productivities of Lakes Ontario and Erie. *Limnology and Oceanography*, 41(2), 352-359.
- Bray, E. E., & Evans, E. D. (1961). Distribution of n-paraffins as a clue to recognition of source beds. *Geochimica et Cosmochimica Acta*, 22(1), 2-15.
- Chirouze, F., Dupont-Nivet, G., Huyghe, P., van der Beek, P., Chakraborti, T., Bernet, M., & Erens, V. (2012). Magnetostratigraphy of the Neogene Siwalik Group in the far eastern Himalaya: Kameng section, Arunachal Pradesh, India. *Journal of Asian Earth Sciences*, 44, 117-135.
- Coutand, I., Barrier, L., Govin, G., Grujic, D., Hoorn, C., Dupont-Nivet, G., & Najman, Y. (2016). Late Miocene-Pleistocene evolution of India-Eurasia convergence partitioning between the Bhutan Himalaya and the Shillong Plateau: New evidences from foreland basin deposits along the Dungsam Chu section, eastern Bhutan. *Tectonics*, 35(12), 2963-2994.
- Cranwell, P. A., Eglinton, G., & Robinson, N. (1987). Lipids of aquatic organisms as potential contributors to lacustrine sediments—II. *Organic Geochemistry*, 11(6), 513-527.
- Dzou, L. I., Holba, A. G., Ramón, J. C., Moldowan, J. M., & Zinniker, D. (1999). Application of new diterpane biomarkers to source, biodegradation and mixing effects on Central Llanos Basin oils, Colombia. *Organic Geochemistry*, 30(7), 515-534.
- Ficken, K. J., Li, B., Swain, D. L., & Eglinton, G. (2000). An n-alkane proxy for the sedimentary input of submerged/floating freshwater aquatic macrophytes. *Organic Geochemistry*, 31(7-8), 745-749.
- Gradstein, F. M., Ogg, J. G., Schmitz, M. B., & Ogg, G. M. (Eds.). (2012). *The geologic time scale 2012*. Elsevier.
- Hirschmiller, J., Grujic, D., Bookhagen, B., Coutand, I., Huyghe, P., Mugnier, J. L., & Ojha, T. (2014). What controls the growth of the Himalayan foreland fold-and-thrust belt?. *Geology*, 42(3), 247-250.
- McKirdy, D. M., Aldridge, A. K., & Ypma, P. J. M. (1983). A geochemical comparison of some crude oils from pre-Ordovician carbonate rocks. In *Advances in Organic Geochemistry 1981*(pp. 99-107). Wiley Chichester.
- Moldowan, J. M., Seifert, W. K., & Gallegos, E. J. (1985). Relationship between petroleum composition and depositional environment of petroleum source rocks. *AAPG bulletin*, 69(8), 1255-1268.
- Ojha, T. P., Butler, R. F., DeCelles, P. G., & Quade, J. (2009). Magnetic polarity stratigraphy of the Neogene foreland basin deposits of Nepal. *Basin Research*, 21(1), 61-90.
- Peters, K. E., & Moldowan, J. M. (1993). The biomarker guide. Prentice Hall. *Englewood Cliffs, NJ*.