



CODEN [USA]: IAJPBB

ISSN : 2349-7750

INDO AMERICAN JOURNAL OF PHARMACEUTICAL SCIENCES

SJIF Impact Factor: 7.187

Available online at: <http://www.iajps.com>

Research Article

A DESCRIPTIVE STUDY ON THE PELVIC MASS IN WOMENS IN TERMS OF PARITY AND AGE

¹Dr Taimoor Hafeez, ²Dr Iqra Idrees,

¹Shaikh Zayed Hospital Lahore, ²Gujranwala Medical College Gujranwala.

Article Received: November 2020 Accepted: December 2020 Published: January 2021

Abstract:

Objective: The aim of this study was to Analyze the pelvic mass patients in terms of parity, age, clinical presentation. **material And Methods:** The design of this study was a descriptive study. This study was carried out at shaikh Zayed Hospital Lahore and the duration of this study was from January 2020 to December 2020. We included all the patients of pelvic mass without any discrimination of parity and age, the disease was diagnosed through clinical assessment, history diagnosis and also through USG. Confirmation of pelvic mass type was made through histopathology and surgery.

Results: We included in our research 110 patients diagnosed with pelvic mass; among these patients 61 were (55.45%) in the age group of 30 – 50 years and one patient were observed as under twenty and two were above seventy years of age. Maximum of the women were in the category of parous. Through clinical presentations we observed that 42 patients faced lower pain of the abdomen (38.18%) and 38 menstrual disturbances patients were also reported (34.54%). First clinical assessment was observed in 84 patients (76.36%) further confirmation was made through USG. In all the patients we included 104 patients (94.54%) with the help of USG. Gentile tract tumor was observed in 68 patients (61.82%); whereas, four patients were diagnosed with non-gynecological mass. Size of mass was observed (above 20 cm) in 10 patients (9.1%); however, fewer common observations were made about the malignant lesions below ten centimeters in size.

Conclusion: Every patient of pelvic mass without any discrimination of parity and age was treated and investigated in detail. Diagnosis of the non-gynecological masses was also made; therefore, competent surgeons were assigned the task to treat the patients of pelvic masses.

Keywords: Pelvic Mass, Diagnosed, Palpable, Genital Organs, Tumor, Sonography.

Corresponding author:

Dr. Taimoor Hafeez,

Shaikh Zayed Hospital Lahore.

QR code



Please cite this article in press Taimoor Hafeez et al, A Descriptive Study On The Pelvic Mass In Womens In Terms Of Parity And Age., Indo Am. J. P. Sci, 2021; 08(1).

INTRODUCTION:

In the practice of gynecology, the incidence of pelvic pain is common and repeated, which is measurable and palpable arising from the presence of the extra-genital and extra genital organs in pelvis. Above 289,000 ladies are reported in USA about the adnexal mass accumulation through clinical and physical examination. In the pre-operative examination, it becomes a complex challenge. Detailed physical assessment and history about the pelvic assessment assists the diagnosis [1]. Obese patients having the inflammatory disease of pelvic face painful examination and palpation is also difficult. To plan and diagnose the treatment of such patients, method of diagnosis is sonography (transvaginal and trans-abdominal real-time imaging). Excellent results can be produced through transvaginal imaging about the pelvic organs; whereas, beyond the range masses are difficult to find out and determine, there are chances that diagnosis will miss those masses. Tissue contrast can best be achieved through MRI scans which permits the visualization of the posterior region about the false and true pelvis and it can also discriminate blood or fluid presence in the ovarian cyst. Uterus can also be imaged through MRI scan if USG cannot observe it in order to revoke any chance of misdiagnosis about the absence of congenital in uterus. Reasonable diagnosis can be made through sonographic and clinical features for the interpretation of the results. Sometimes, no clear information about the nature of mass is available till the performance of laparotomy [2].

These days, frequent practice has been observed for the determination adnexal mass through laparoscopy. Laparoscopy when compared to laparotomy has a plus feature of the less rate of morbidity, shorter hospital stays, lesser de novo formation of adhesion, decreased postoperative pain, faster recovery, reduced cure cost and better cosmetic results. However, there is a need of the careful pre-operative assessment for a successful and appropriate laparoscopy use to remove adnexal mass without compromising any of the clinical results. Risk factor identification has been enforced by numerous authors in order to careful use of laparoscopy for the management of the masses that may benign. The importance of the laparoscopic removal avoidance of ovarian cancer is because unintentional spillage could make the prognosis even worse and / or it may also mandate chemotherapy administration to Stage – I patients those who may then not be receiving it. Genital masses are adnexal and uterine [3]. These uterine masses are pregnancy, congenital irregularities, adenomyosis, pyometra, sarcoma, trophoblastic disease and leiomyoma. It may

also be tubal, ovarian or vaginal. Other related genital masses are pelvic lithopedion's and abscess. In the category of the extra-genital masses there are urinary bladder, appendicular mass, pelvic kidney, diverticula, retro-peritoneal tumor, bowel tumor, abdominal wall lesions, rare pelvic castle-man disease and retro-peritoneal fibrosis that also has an involvement of the pelvic para-cervical lymph nodes. After clinical examination, history, radiological imaging methods, laboratory investigations and surgically invasive techniques (laparotomy and laparoscopy), histopathology report helps in the definite diagnosis [4]. Pelvic mass excision through histologic assessment remains as the only available and reliable method of diagnosis establishment. Analysis of pelvic mass patients in terms of parity, age, clinical presentation, operative procedure and pathology was as per the mass type in the tertiary hospitals of Sindh was the aim of this research.

MATERIAL AND METHODS:

110 patients diagnosed with pelvic mass; among these patients 61 were (55.45%) in the age group of 30 – 50 years and one patient were observed as under twenty and two were above seventy years of age. This study was carried out at shaikh Zayed Hospital Lahore and the duration of this study was from January 2020 to December 2020. OPD admitted the patients after the diagnosis of pelvic mass which is also included in the research. Evaluation of the symptoms was made through USG and results were recorded about the size, type, consistency, ascitic fluid presence and metastasis evidence in para-aortic lymph nodes and liver. Another provisional diagnosis was also made after observation of a detailed history through abdominal, pelvic and physical assessment. Pelvic mass was observed in number of women through clinical investigations and confirmation was made through USG at a frequency of (3.5 MHz) with the help of linear probe. Operative processes include simple cystectomy, bilateral sapling oophorectomy, debulking surgery or total abdominal hysterectomy.

Analysis of the histopathological reports was carried out for the identification of the mass and its nature, we also followed our patients. Statistical data analysis was made through SPSS – 11.

RESULTS:

We included in our research 110 patients diagnosed with pelvic mass; among these patients 61 were (55.45%) in the age group of 30 – 50 years and one patient were observed as under twenty and two were above seventy years of age as shown in Table – I. Maximum of the women were in the category of

parous as shown in Table – II. Through clinical presentations we observed that 42 patients faced lower pain of the abdomen (38.18%) and 38 menstrual disturbances patients were also reported (34.54%) as shown in Table – III. First clinical assessment was observed in 84 patients (76.36%) further confirmation was made through USG. In all the patients we included 104 patients (94.54%) with the help of USG. Gentile tract tumor was observed in 68 patients (61.82%);

whereas, four patients were diagnosed with non-gynecological mass. Size of mass was observed (above 20 cm) in 10 patients (9.1%); however, fewer common observations were made about the malignant lesions below ten centimeters in size. Surgical operation was carried out in 95 patients (86.36%); these patients also included 43 conservatives (39.1%) and 52 radicals (47.27%) surgery patients. In the light of medical advice 5 patients (4.54%) were left.

Table I: Age and Parity Distribution of Patients (n = 110)

	Details	Number	Percentage
Age (in years)	≤ 20	1	0.91
	21 – 29	33	30
	30 – 39	33	30
	40 – 49	25	22.72
	50 – 59	12	10.91
	60 – 69	4	3.63
	≥ 70	2	1.82
Parity	0	35	31.82
	3 – Jan	25	22.72
	6 – Apr	24	21.82
	6 +	26	23.64

Table II: Clinical Presentation and Diagnosis of Pelvic Mass (n = 110)

	Detail	Number	Frequency
Clinical presentation	Mass alone	16	14.54
	Lower abdominal pain	42	38.18
	Mass with pain	14	12.73
	Menstrual disturbance	38	34.54
Diagnosis	By clinical examination	84	76.36
	By ultrasonography	104	94.54
	At surgery	6	5.45

Table III: Pattern of Masses Among Patients (n = 110)

Mass pattern	No. of Patients n (%)	Benign n (%)	Malignant n (%)
		30(27.27) 24(21.82)	11(10) 02(1.82)
Genital tract tumors	41(37.27)	-	-
Ovarian	26(23.6)	-	-
Uterine	16(14.54)	-	-
Cervix	14(12.72)	-	-
Tubo-ovarian mass Ectopic tubal gestation	01(0.91)	-	-
Hydrosalpinx	01(0.91)	-	-
Haematosalpinx	02(1.82)	-	-
Pyosalpinx	02(1.82)	-	-
Pyometra	01(0.91)	-	-
Haematometra	01(0.91)	-	-
Pelvic abscess Non-Gynaecological Masses			
Avulsed spleen Hydatid cyst of spleen	01(0.91)	-	-
Appendicular abscess	01(0.91)	-	-
Ruptured bladder	01(0.91)	-	-
	01(0.91)	-	-
	01(0.91)	-	-

Table IV: Size of Mass at The Time of Diagnosis (n = 110)

Size (in centimeters)	Frequency (%)
5-10	(60)
11-15	(18.18)
16-20	(12.73)
> 20	(9.1)

DISCUSSION:

Maximum number of patients were in the category of productive group of age. Ovarian tumor was the repeated benign of mass; whereas, second common cause was uterine leiomyoma was the second most common cause [5]. Both extremes of life were not common for pelvic mass. Contrarily, we also observed that in the age group of thirteen years hematometra and haematocolpus was commonly found in the presentation of pelvic mass. Malignancy risk was increased with the advancement of the age; we also observed that above fifty years of patients common was ovarian tumor [6]. Mallick also reports same observation as he states that ovarian cancer increased incidence in the age of forty-five and above (60% – 70%) [7]. In the consideration of the relation of parity in women we observed parous women in majority. Uterine fibroids and ovarian tumors are associated to the nulliparity risk in women. Our research also observed that there was an increased malignancy incidence of ovarian tumor in nullipara women [8]. However, another cause was common in the nullipara women known as leiomyoma, we also observed its presence in grand-multiparous women. Among grand-multipara women having fibroid uterus, in the observation of the last-born age was in the range of 10 – 20 years [9]. According to Alam in the age of 40 – 60 years there was a common presentation of the ovarian tumors, in terms of old and young respectively we observed 65 years and 15 years. An early aged pregnancy and late menopause were associated to the ovary trauma caused by the recurrent ovulation. USG diagnostic value about the pelvic mass location is proved. Diagnosis was made through transabdominal USG among 104 patients in the total research sample.

According to Qureshi, transvaginal USG is superior in sixty-three percent of the patients, equal in twenty-seven percent and inferior in ten percent of patients in comparison to the trans-abdominal sonography. To detect the large amount of pelvic mass less importance is given to the trans-vaginal ultrasound; in order to measure and monitor ovarian follicle, endometrial carcinoma, polycystic ovarian syndrome and for the suspected ectopic pregnancy.

Surgery was carried out in 95 patients (86.36%). Radical surgery was carried out in the patients of bilateral salping oophorectomy and total abdominal hysterectomy. Conservative surgery was carried out in 43 patients (39.1%) as myomectomy, salping oophorectomy, cystectomy and biopsy was also carried out. Pelvic mass assessment detected through routine USG or in the patient of acute symptoms, which needs few methodical steps [10]. Lesion primary sight detection is very important.

Transvaginal USG is an interactive and dynamic assessment along with the echo structure analysis, and pelvic mass 'elasticity', which also allows the site-specific pain evaluation in various areas of the pelvic and mass movement evaluation in connection with the nearby structures. However, during research, our facility lacked in transvaginal USG. The variation between malignant and benign ovarian tumor is considered as a challenge in the clinical assessment. Color Doppler USG combined with tumor markers may also help in the improvement of the method accuracy. We found that 30 patients (27.27%) among the 41 women had benign tumor and ovarian mass and 11 malignant ovarian tumor patients were (10%). This was confirmed through surgery and histopathological outcomes [11].

In four out of 110 patients, during surgery a non-gynecological mass was observed. Therefore, for the definite diagnosis we need laparotomy. Gynecologists and surgeons dealt with the patient of non-gynecological masses. Splenectomy was carried out in two patients, bladder repair was carried out in one patient and also referred to the urologist for expert advice and appendectomy in one patient.

CONCLUSION:

Every patient of pelvic mass without any discrimination of parity and age was treated and investigated in detail. Diagnosis of the non-gynecological masses was also made; therefore, competent surgeons were assigned the task to treat the patients of pelvic masses.

REFERENCES:

1. Moroni, R.M., et al., Presentation and treatment of uterine leiomyoma in adolescence: a systematic review. *BMC women's health*, 2015. 15(1): p. 4.
2. Parra-Herran, C., et al., Inflammatory myofibroblastic tumor of the uterus: clinical and pathologic review of 10 patients including a subset with aggressive clinical course. *The American journal of surgical pathology*, 2015. 39(2): p. 157-168.
3. Kho, K.A., et al., Risk of occult uterine sarcoma in women undergoing hysterectomy for benign indications. *Obstetrics & Gynecology*, 2016. 127(3): p. 468-473.
4. Oduyebo, T., et al., The value of re-exploration in patients with inadvertently morcellated uterine sarcoma. *Gynecologic oncology*, 2014. 132(2): p. 360-365.
5. Brunham, R.C., S.L. Gottlieb, and J. Paavonen, Pelvic inflammatory disease. *New England*

- Journal of Medicine, 2015. 372(21): p. 2039-2048.
6. Kelleher, C.M. and A.M. Goldstein, Adnexal masses in children and adolescents. *Clinical obstetrics and gynecology*, 2015. 58(1): p. 76-92.
 7. Sharma, J.B., Current diagnosis and management of female genital tuberculosis. *The Journal of Obstetrics and Gynecology of India*, 2015. 65(6): p. 362-371.
 8. Acién, P. and M. Acién, The presentation and management of complex female genital malformations. *Human reproduction update*, 2015. 22(1): p. 48-69.
 9. Roitman, P.D., et al., Is needle biopsy clinically useful in preoperative grading of central chondrosarcoma of the pelvis and long bones? *Clinical Orthopaedics and Related Research*, 2017. 475(3): p. 808-814.
 10. Valour, F., et al., Actinomycosis: etiology, clinical features, diagnosis, treatment, and management. *Infection and drug resistance*, 2014. 7: p. 183.
 11. Mishra, V.V., et al., Prevalence; characteristics and management of endometriosis amongst infertile women: a one year retrospective study. *Journal of clinical and diagnostic research: JCDR*, 2015. 9(6): p. QC01.