Role of Benzylic Deprotonation in Ni-Catalyzed Benzylic Dehydrogenation

Pengpeng Zhang, Rachel L. Cantrell, and Timothy R. Newhouse*

Department of Chemistry, Yale University
225 Prospect St., New Haven, Connecticut 06520-8107

*E-mail: timothy.newhouse@yale.edu

Table of Contents

General Experimental.....	SI-2
Procedure and Characterization Data for Base Screen.....	SI-3
Procedure and Characterization Data for Temperature-Dependent Studies.....	SI-7
Procedure and Characterization Data for Time-Dependent Studies	SI-12
Procedure and Characterization Data for Substrate Scope.....	SI-15
References	SI-45

General Experimental

General Experimental Procedures: All reactions were carried out under an inert nitrogen atmosphere with dry solvents under anhydrous conditions unless otherwise stated. All reactions were capped with a rubber septum, or Teflon-coated silicon microwave cap unless otherwise stated. Stainless steel cannula or syringe was used to transfer solvent, and air- and moisture sensitive liquid reagents. Reactions were monitored by thin-layer chromatography (TLC) carried out on 0.25 mm Merck silica gel plates (60F-254) using UV light as the visualizing agent and potassium permanganate, an acidic solution of *p*-anisaldehyde, phosphomolybdic acid, or I₂ on SiO₂ as developing agents. Flash column chromatography employed SiliaFlash® P60 (40-60 µm, 230-400 mesh) silica gel purchased from SiliCycle, Inc.

Materials: All reaction solvents were purified using a Seca solvent purification system by Glass contour. TMEDA, Et₃N were distilled over CaH₂. Zn(TMP)₂ (0.5 M in toluene), PMe₃ (1.0 M in THF), TBAI and *n*-BuLi (2.5 M in hexanes) were purchased from Sigma-Aldrich. NiBr₂(dme) was purchased from Strem. All other reagents were used as received without further purification, unless otherwise stated.

Instrumentation: All new compounds were characterized by means of ¹H NMR, ¹³C NMR, ¹⁹F NMR, FT-IR (thin film), and HR-MS. Copies of the ¹H- and ¹³C-NMR spectra can be found at the end of each experimental procedure. NMR spectra were recorded using a Varian 400 MHz NMR spectrometer, Varian 500 MHz NMR spectrometer, or a Varian 600 MHz NMR spectrometer. All ¹H-NMR data are reported in δ units, parts per million (ppm), and were calibrated relative to the signals for residual chloroform (7.26 ppm) in deuteriochloroform (CDCl₃). ¹H NMR for starting materials and after deuteration were acquired with the same protocol (instrument, D1, number of scans, etc.) in order to ensure cancellation of error due to variable T1 relaxation times. All ¹³C-NMR data are reported in ppm relative to CDCl₃ (77.16 ppm) or Acetone-*d*₆ (206.26 ppm) and were obtained with ¹H decoupling unless otherwise stated. The following abbreviations or combinations thereof were used to explain the multiplicities: s = singlet, d = doublet, t = triplet, q = quartet, abq = ab quartet, br = broad, m = multiplet, and a = apparent. All IR spectra were taken on an FT-IR/Raman Thermo Nicolet 6700. High resolution mass spectra (HR-MS) were recorded on a Waters Xevo Qtof mass spectrometer using ESI-TOF (electrospray ionization-time of flight). Gas chromatography mass spectra (GC-MS) were recorded on an Agilent Technologies 6890N Network Gas Chromatograph System with an Agilent Technologies 5973N Mass Selective Detector. Optical rotation data was obtained using a Perkin-Elmer 341 polarimeter.

Procedure and Characterization Data for Base Screen

Entry	Base	Deuteration (%)	Conversion (%) ^a
1	LiOH	nd	13
2	CS ₂ CO ₃	nd	8
3	K ₃ PO ₄	nd	12
4	LiOtBu	nd	52
5	<i>n</i> -BuLi	6	85
6	LDA	7	78
7	LiTMP	1	72
8	LiCyan	11	78
9	LHMDS	15	85
10	Zn(HMDS) ₂	nd	53
11	Zn(TMP) ₂	3	34
12	Zn(TMP)Cl•LiCl	1	49
13	Mg(TMP)Cl•LiCl	nd	70

[a] Yield of deuterated and conversion were determined by ¹H NMR analysis using 1,3,5-trimethoxybenzene as an internal standard. [b] Bases in entry 12 were made via the procedure by Zhao et. al.¹

General Procedure A (entry 1-3):

To a flame-dried microwave vial equipped with a magnetic stir bar was added solid bases (0.075mmol, 1.5 equiv). The reaction vessel was evacuated and backfilled with N₂ (this process was repeated three times). To the reaction vessel was added a solution of starting material (0.05mmol, 1 equiv.) in anhydrous 1,4-dioxane (0.1 M, 0.5 mL). The reaction was allowed to stir at room temperature for 30 minutes before being placed in an 85 °C oil bath, where it was stirred for 4 hours.

Immediately after removal from heat, the reaction was quenched with D₂O (0.5 mL) and stirred for 5 minutes to cool down to room temperature. A solution of internal standard 1,3,5-trimethoxybenzene in EtOAc was added before extraction with EtOAc (4 x 1.5 mL), and the combined organic extracts were dried with Na₂SO₄ and filtered through a short silica plug and washed with EtOAc (5 mL) and concentrated under reduced pressure by rotary evaporation. Then 0.2 mL of CDCl₃ was added and the excess solvent was removed under vacuum. Finally, the deuteration and degradation percentage was determined by crude ¹H NMR based on the internal standard.

General Procedure B (entry 4-13):

A flame-dried microwave vial equipped with a magnetic stir bar was evacuated and backfilled with N₂ (this process was repeated three times). To the reaction vessel was added a solution of starting material (0.05 mmol, 1 equiv.) in anhydrous 1,4-dioxane (0.1 M, 1 mL), followed by the solution of commercially available or fresh-made base (0.075 mmol, 1.5 equiv). The reaction was stirred at room temperature for 30 mins and then moved to an 85 °C oil bath for 4 hours.

Immediately after removal from heat, the reaction was quenched with D₂O (0.5 mL) and stirred for 5 minutes to cool down to room temperature. A solution of internal standard 1,3,5-trimethoxybenzene in EtOAc was added before extraction with EtOAc (4 x 1.5 mL), and the combined organic extracts were dried with Na₂SO₄ and filtered through a short silica plug and washed with EtOAc (5 mL) and concentrated under reduced pressure by rotary evaporation. Then 0.2 mL of CDCl₃ was added and the excess solvent was removed under vacuum. Finally, the deuteration and degradation percentage was determined by crude ¹H NMR based on the internal standard.

Note: 2 equiv. of internal standard was added in entries 1-3, and 11.

2-phenethyl-5-(trifluoromethyl)pyridine (**1**)

Compound **1** was prepared according to known procedure,² and the characterization data match those previously reported in the literature.

¹H NMR (400 MHz, CDCl₃): δ 8.85 (s, 1H), 7.79 (dd, *J* = 8.2, 2.4 Hz, 1H), 7.32 – 7.27 (m, 2H), 7.24 – 7.16 (m, 4H), 3.21 – 3.16 (m, 2H), 3.13 – 3.07 (m, 2H)

¹³C NMR (101 MHz, CDCl₃): δ 165.4, 146.4 (q, *J*_{C-F} = 4.1 Hz), 141.0, 133.4 (q, *J*_{C-F} = 3.5 Hz), 128.6, 128.5, 126.6, 124.4 (q, *J*_{C-F} = 33.0 Hz), 123.9 (q, *J*_{C-F} = 272.1 Hz), 122.8, 40.2, 35.7

¹⁹F NMR (376 MHz, CDCl₃): δ –62.2 (s)

ESI-HRMS (m/z): [M+H]⁺ calc'd for C₁₄H₁₃F₃N⁺: 252.0995; found: 252.1000

Procedure and Characterization Data for Temperature-Dependent Studies

[a] Yield of dehydrogenation was determined by ¹H NMR analysis using 1,3,5-trimethoxybenzene as an internal standard. Ox = 2-bromo-5-methylthiophene

General Procedure C: (condition A)

A flame-dried microwave vial equipped with a magnetic stir bar was evacuated and backfilled with N₂ (this process was repeated three times). To the reaction vessel was added a solution of starting material **2** (0.1 mmol, 1 equiv.) in anhydrous 1,4-dioxane (0.1 M, 1 mL), followed by the solution of commercial Zn(TMP)₂ (0.3 mL, 0.5 M in toluene, 0.15 mmol, 1.5 equiv). The reaction was stirred at different temperature for 2 h as the table shows.

Immediately after removal from heat, the reaction was quenched with D₂O (0.5 mL) and stirred for 5 minutes to cool down to room temperature. A solution of internal standard 1,3,5-trimethoxybenzene in EtOAc was added (0.33 mmol unless otherwise stated) before extraction with EtOAc (4 x 1.5 mL), and the combined organic extracts were dried with Na₂SO₄ and filtered through a short silica plug and washed with EtOAc (5 mL) and concentrated under reduced pressure by rotary evaporation. Then 0.2 mL of CDCl₃

was added and the excess solvent was removed under vacuum. Finally, the deuteration and degradation percentage was determined by crude ^1H NMR based on the internal standard.

General Procedure D: (condition B)²

To a flame-dried microwave vial equipped with a magnetic stir bar was added starting material **2** (0.20 mmol, 1.0 equiv). The reaction vessel was sealed, evacuated and backfilled with N_2 (this process was repeated three times). To the reaction vessel was added anhydrous 1,4-dioxane (1.5 mL) and commercial $\text{Zn}(\text{TMP})_2$ (0.48 mL, 0.5 M in toluene, 0.3 mmol, 1.5 equiv) at room temperature. Then to this mixture was added 0.5 mL of a stock solution containing $\text{NiBr}_2(\text{dme})$ (6.2 mg, 0.02 mmol, 10 mol %), PMe_3 (0.06 mL, 0.06 mmol, 30 mol%), and 2-bromo-5-methylthiophene (27.4 μL , 0.24 mmol, 1.2 equiv) in 1,4-dioxane. The reaction vessel was placed into a pre-heated oil bath for 2 h (at different temperature as table shows).

The reaction mixture was cooled to ambient temperature and quenched by the addition of sat. aq. NH_4Cl (5 mL). The reaction mixture was diluted with EtOAc (5 mL) and the organic phase was separated. The aqueous phase was extracted with EtOAc (3×5 mL) and the combined organic extracts were washed with brine (5 mL), dried over anhydrous Na_2SO_4 , filtered, and concentrated under reduced pressure by rotary evaporation. Yield was determined by crude ^1H NMR based on the internal standard.

2-phenethylpyridine (2)

Compound **2** was prepared according to known procedure,² and the characterization data match those previously reported in the literature.

¹H NMR (400 MHz, CDCl₃) δ 8.57 (d, *J* = 4.1 Hz, 1H), 7.55 (td, *J* = 7.7, 1.8 Hz, 1H), 7.31 – 7.24 (m, 2H), 7.23 – 7.18 (m, 3H), 7.06 – 7.13 (m, 2H), 3.13 – 3.04 (m, 4H).

The deuteration is determined by the compare the integration change of benzylic protons at 3.13 – 3.04 (m, 4H) with compound **2** by setting integration of 7.55 ppm as 1.00.

Procedure and Characterization Data for Time-Dependent Studies

[a] Yield of dehydrogenation was determined by ¹H-NMR analysis using 1,3,5-trimethoxybenzene as an internal standard. Ox = 2-bromo-5-methylthiophene

2-phenethylpyrazine (3)

3

Compound **3** was prepared according to known procedure,² and the characterization data match those previously reported in the literature.

¹H NMR (400 MHz, CDCl₃): δ 8.52 (s, 1H), 8.40 (d, *J* = 2.5 Hz, 1H), 8.36 (d, *J* = 1.5 Hz, 1H), 7.30 – 7.24 (m, 2H), 7.21 – 7.17 (m, 3H), 3.15 – 3.05 (m, 4H)

¹³C NMR (101 MHz, CDCl₃): δ 156.9, 144.8, 144.2, 142.5, 140.9, 128.6, 128.5, 126.4, 37.3, 35.5

ESI-HRMS (*m/z*): [M+H]⁺ calc'd for C₁₂H₁₃N⁺: 185.1073; found: 185.1079

3

Procedure and Characterization Data for Substrate Scope

[a] Deuteration percentage was determined by crude ¹H NMR. [b] Isolated Yield of Ni-catalyzed dehydrogenation was shown in parentheses; Ox = 2-bromo-5-methylthiophene for condition B. [c] compounds **6** and **13** gave obvious degradation.

General Procedure E for deprotonation of substrate scope (condition A):

A flame-dried microwave vial equipped with a magnetic stir bar was evacuated and backfilled with N₂ (this process was repeated three times). To the reaction vessel was added a solution of starting material (0.1 mmol, 1 equiv.) in anhydrous 1,4-dioxane (0.1 M, 1 mL), followed by the commercial solution of Zn(TMP)₂ (0.3 mL, 0.5 M in toluene, 0.15 mmol, 1.5 equiv). The reaction was stirred at 85 °C for 4 hours.

Immediately after removal from heat, the reaction was quenched with D₂O (0.5 mL) and stirred for 5 minutes to cool down to room temperature. A solution of internal standard 1,3,5-trimethoxybenzene in EtOAc was added (0.33 mmol unless otherwise stated) before extraction with EtOAc (4 x 1.5 mL), and the combined organic extracts were dried with Na₂SO₄ and filtered through a short silica plug and washed with EtOAc (5 mL) and concentrated under reduced pressure by rotary evaporation. Then 0.2 mL of CDCl₃

was added and the excess solvent was removed under vacuum. Finally, the deuteration and degradation percentage was determined by crude ^1H NMR based on the internal standard.

General Procedure F for dehydrogenation (condition B)²:

To a flame-dried microwave vial equipped with a magnetic stir bar was added starting material **1** (0.20 mmol, 1.0 equiv) and TBAI (148 mg, 0.40 mmol, 2.0 equiv). The reaction vessel was sealed, evacuated and backfilled with N_2 (this process was repeated three times). To the reaction vessel was added anhydrous 1,4-dioxane (1.5 mL) and commercial Zn(TMP)_2 (0.48 mL, 0.5 M in toluene, 0.24 mmol, 1.2 equiv) at room temperature. Then to this mixture was added 0.5 mL of a stock solution containing $\text{NiBr}_2(\text{dme})$ (6.2 mg, 0.02 mmol, 10 mol %), PMe_3 (0.06 mL, 0.06 mmol, 30 mol%), and 2-bromo-5-methylthiophene (27.4 μL , 0.24 mmol, 1.2 equiv) in 1,4-dioxane. The reaction vessel was placed into a pre-heated 85 °C oil bath, and stirred until complete consumption of starting material as determined by TLC analysis. (Note: more reproducible results can be obtained using the sequential addition procedure.)

The reaction mixture was cooled to ambient temperature and quenched by the addition of sat. aq. NH_4Cl (5 mL). The reaction mixture was diluted with EtOAc (5 mL) and the organic phase was separated. The aqueous phase was extracted with EtOAc (3×5 mL) and the combined organic extracts were washed with brine (5 mL), dried over anhydrous Na_2SO_4 , filtered, and concentrated under reduced pressure by rotary evaporation. The crude mixture was purified by flash column chromatography on silica gel.

5-methoxy-2-phenethylpyridine (4)

Compound **4** was prepared according to known procedure,² and the characterization data match those previously reported in the literature.

^1H NMR (400 MHz, CDCl_3): δ 8.27 (d, J = 3.0 Hz, 1H), 7.29 – 7.25 (m, 2H), 7.22 – 7.16 (m, 3H), 7.09 (dd, J = 8.5, 3.0 Hz, 1H), 6.99 (d, J = 8.5 Hz, 1H), 3.83 (s, 3H), 3.06 – 3.01 (m, 4H)

^{13}C NMR (101 MHz, CDCl_3): δ 154.0, 153.4, 141.8, 136.7, 128.6, 128.4, 126.0, 123.1, 121.2, 55.7, 39.3, 36.4

ESI-HRMS (m/z): $[\text{M}+\text{H}]^+$ calc'd for $\text{C}_{14}\text{H}_{16}\text{NO}^+$: 214.1226; found: 214.1232

5-fluoro-2-phenethylpyridine (**5**)

Compound **5** was prepared according to known procedure,² and the characterization data match those previously reported in the literature.

¹H NMR (400 MHz, CDCl₃): δ 8.43 (d, J = 3.0 Hz, 1H), 7.31 – 7.25 (m, 3H), 7.22 – 7.15 (m, 3H), 7.05 (dd, J = 8.6, 4.4 Hz, 1H), 3.13 – 3.03 (m, 4H)

¹³C NMR (101 MHz, CDCl₃): δ 158.2 (d, J_{C-F} = 253.4 Hz), 157.2 (d, J_{C-F} = 3.9 Hz), 141.4, 137.4 (d, J_{C-F} = 23.1 Hz), 128.6, 128.5, 126.1, 123.7 (d, J_{C-F} = 3.9 Hz), 123.1 (d, J_{C-F} = 18.1 Hz), 39.4, 36.1

¹⁹F NMR (376 MHz, CDCl₃): δ –131.2 (q, J = 3.8 Hz)

ESI-HRMS (m/z): [M+H]⁺ calc'd for C₁₃H₁₃FN⁺: 202.1027; found: 202.1032

6-phenethylnicotinonitrile (**6**)

Compound **6** was prepared according to known procedure,² and the characterization data match those previously reported in the literature.

¹H NMR (400 MHz, CDCl₃): δ 8.84 (d, *J* = 1.7 Hz, 1H), 7.81 (dd, *J* = 8.1, 2.2 Hz, 1H), 7.29 (dd, *J* = 8.0, 6.5 Hz, 2H), 7.28 – 7.17 (m, 4H), 3.20 – 3.16 (m, 2H), 3.10 – 3.05 (m, 2H)

¹³C NMR (101 MHz, CDCl₃): δ 165.9, 152.3, 140.6, 139.3, 128.6, 128.5, 126.4, 123.2, 117.0, 107.5, 40.4, 35.4

ESI-HRMS (*m/z*): [M+H]⁺ calc'd for C₁₄H₁₃N₂⁺: 209.1073; found: 209.1079

2-phenethyl-4-(trifluoromethyl)pyridine (7)

Compound **7** was prepared according to known procedure,² and the characterization data match those previously reported in the literature.

¹H NMR (400 MHz, CDCl₃): δ 8.85 (s, 1H), 7.79 (dd, *J* = 8.2, 2.4 Hz, 1H), 7.32 – 7.27 (m, 2H), 7.24 – 7.16 (m, 4H), 3.21 – 3.16 (m, 2H), 3.13 – 3.07 (m, 2H)

¹³C NMR (101 MHz, CDCl₃): δ 165.4, 146.4 (q, *J*_{C-F} = 4.1 Hz), 141.0, 133.40 (q, *J*_{C-F} = 3.5 Hz), 128.6, 128.5, 126.6, 124.4 (q, *J*_{C-F} = 33.0 Hz), 123.9 (q, *J*_{C-F} = 272.1 Hz), 122.8, 40.2, 35.7

¹⁹F NMR (376 MHz, CDCl₃): δ –62.2 (s)

ESI-HRMS (m/z): [M+H]⁺ calc'd for C₁₄H₁₃F₃N⁺: 252.0995; found: 252.1000

2-(cyclohexylmethyl)pyridine (**8**)

Compound **8** was prepared according to known procedure,² and the characterization data match those previously reported in the literature.

¹H NMR (400 MHz, CDCl₃): δ 8.52 (d, *J* = 4.8 Hz, 1H), 7.57 – 7.53 (m, 1H), 7.09 – 7.05 (m, 2H), 2.64 (d, *J* = 7.1 Hz, 2H), 1.81 – 1.72 (m, 1H), 1.71 – 1.58 (m, 5H), 1.27 – 1.09 (m, 3H), 1.06 – 0.93 (m, 2H)

¹³C NMR (101 MHz, CDCl₃): δ 161.5, 149.4, 136.0, 123.7, 120.9, 46.5, 38.8, 33.3, 26.6, 26.4

ESI-HRMS (*m/z*): [M+H]⁺ calc'd for C₁₂H₁₈N⁺: 176.1434; found: 176.1439

2-(5-methoxypentyl)pyridine (**9**)

Compound **9** was prepared according to known procedure,² and the characterization data match those previously reported in the literature.

¹H NMR (400 MHz, CDCl₃): δ 8.50 (d, *J* = 4.0 Hz, 1H), 7.58 – 7.53 (m, 1H), 7.14 – 7.05 (m, 2H), 3.35 (t, *J* = 6.6 Hz, 2H), 3.30 (s, 3H), 2.88 – 2.73 (m, 2H), 1.78 – 1.70 (m, 2H), 1.63 – 1.56 (m, 2H), 1.44 – 1.36 (m, 2H)

¹³C NMR (101 MHz, CDCl₃): δ 162.4, 149.3, 136.3, 122.8, 121.0, 72.9, 58.6, 38.5, 29.8, 29.6, 26.1

ESI-HRMS (*m/z*): [M+H]⁺ calc'd for C₁₁H₁₈NO⁺: 180.1383; found: 180.1388

1,2,3,4-tetrahydroacridine (**10**)

Compound **10** was prepared according to known procedure,³ and the characterization data match those previously reported in the literature.

¹H NMR (400 MHz, CDCl₃) δ 7.97 (d, J = 8.5 Hz, 1H), 7.76 (s, 1H), 7.67 (dd, J = 8.2, 1.5 Hz, 1H), 7.59 (t, J = 7.7 Hz, 1H), 7.46 – 7.36 (m, 1H), 3.12 (t, J = 6.5 Hz, 2H), 2.95 (t, J = 6.4 Hz, 2H), 2.07 – 1.92 (m, 2H), 1.92 – 1.82 (m, 2H).

¹³C NMR (101 MHz, CDCl₃) δ 159.4, 146.8, 135.1, 131.1, 128.6, 128.4, 127.3, 127.0, 125.6, 33.7, 29.4, 23.4, 23.0.

GC-MS (m/z): [M+H]⁺ calc'd for C₁₃H₁₃N⁺: 183.1; found: 183.1

2-phenethylquinoline (**11**)

Compound **11** was prepared according to known procedure,² and the characterization data match those previously reported in the literature.

¹H NMR (400 MHz, CDCl₃): δ 8.10 – 8.04 (m, 2H), 7.79 (d, *J* = 8.1 Hz, 1H), 7.73 – 7.69 (m, 1H), 7.52 – 7.48 (m, 1H), 7.31 – 7.18 (m, 6H), 3.33 – 3.28 (m, 2H), 3.20 – 3.15 (m, 2H)

¹³C NMR (101 MHz, CDCl₃): δ 161.9, 148.1, 141.7, 136.4, 129.5, 129.0, 128.7, 128.5, 127.7, 126.9, 126.1, 125.9, 121.7, 41.1, 36.1

ESI-HRMS (*m/z*): [M+H]⁺ calc'd for C₁₇H₁₆N⁺: 234.1277; found: 234.1283

3-(5-(((tert-butyldimethylsilyl)oxy)pentyl)-6-phenylpyridazine (12)

Compound **12** was prepared according to known procedure,² and the characterization data match those previously reported in the literature.

¹H NMR (400 MHz, CDCl₃): δ 8.08 (d, *J* = 6.7 Hz, 2H), 7.77 (d, *J* = 8.7 Hz, 1H), 7.54 – 7.45 (m, 3H), 7.37 (d, *J* = 8.7 Hz, 1H), 3.60 (t, *J* = 6.4 Hz, 2H), 3.02 (t, *J* = 6.4 Hz, 2H), 1.90 – 1.80 (m, 2H), 1.56 – 1.47 (m, 2H), 1.47 – 1.35 (m, 4H), 0.89 (s, 9H), 0.04 (s, 6H)

¹³C NMR (101 MHz, CDCl₃): δ 162.3, 157.3, 136.5, 129.7, 128.9, 126.9, 126.6, 123.9, 63.1, 35.9, 32.7, 29.5, 29.0, 26.0, 25.6, 18.3, –5.3

ESI-HRMS (*m/z*): [M+H]⁺ calc'd for C₂₂H₃₅N₂OSi⁺: 371.2513; found: 371.2519

2-(butyl)-4,6-dimethoxy-1,3,5-triazine (13)

Compound **13** was prepared according to known procedure,² and the characterization data match those previously reported in the literature.

¹H NMR (400 MHz, CDCl₃): δ 3.89 (s, 6H), 2.59 (t, *J* = 7.4 Hz, 2H), 1.63 (dt, *J* = 13.1, 7.6 Hz, 2H), 1.32 – 1.19 (m, 2H), 0.79 (t, *J* = 7.4 Hz, 3H)

¹³C NMR (101 MHz, CDCl₃): δ 183.5, 172.3, 54.8, 38.2, 29.3, 22.2, 13.7

ESI-HRMS (*m/z*): [M+H]⁺ calc'd for C₉H₁₆N₃O₂⁺: 198.1237; found: 198.1243

General Experimental Procedure G for substrate preparation⁴

To a flame-dried microwave vial equipped with a magnetic stir bar was added magnesium (264 mg, 11 mol, 1.1 equiv), a catalytic amount of I_2 (5 mg), and THF (5 mL, 2.0 M). The resulting mixture was heated to 80 °C, then removed the oil bath and alkyl bromide (10 mol, 1.0 equiv) was added via syringe dropwise over 10 minutes to maintain a gentle reflux. The reaction vessel was then stirred for 30 minutes at room temperature.

A flame-dried flask under N_2 was charged with aryl halide (5.0 mmol, 1.0 equiv), $\text{Fe}(\text{acac})_3$ (88 mg, 0.25 mmol 5 mol%), N-methylpyrrolidone (3.0 mL), and THF (30 mL, 0.15 M). A solution of alkyl magnesium bromide/chloride (3.0 mL, 2.0 M in THF, 6.00 mmol, 1.2 equiv) is added via syringe to the resulting red solution at 0 °C, causing an immediate color change to dark brown or black. The resulting mixture was stirred at room temperature for 30 minutes, and the reaction was diluted with Et_2O and is carefully quenched by the slow addition of aq. 1.0 M HCl (10 mL). The reaction mixture was diluted with water (20 mL) and EtOAc (20 mL) and the organic phase was separated. The aqueous phase was extracted with EtOAc (3×25 mL) and the combined organic extracts were washed with brine (15 mL), dried over anhydrous Na_2SO_4 , filtered, and concentrated under reduced pressure by rotary evaporation. The crude mixture was purified by flash column chromatography on silica gel.

2-methoxy-6-(3-phenylpropyl)pyridine (**14**)

Compound **14** was prepared from 2-chloro-6-methoxypyridine and (3-bromopropyl)benzene according to procedure G.

^1H NMR (400 MHz, CDCl_3) δ 7.47 (t, $J = 8.1, 7.7$ Hz, 1H), 7.38 – 7.25 (m, 2H), 7.25 – 7.11 (m, 3H), 6.71 (d, $J = 7.2$ Hz, 1H), 6.56 (d, $J = 8.2$ Hz, 1H), 3.94 (s, 3H), 2.82 – 2.59 (m, 4H), 2.22 – 1.87 (m, 2H).

^{13}C NMR (101 MHz, CDCl_3) δ 166.4, 160.2, 142.4, 139.7, 131.4, 126.9, 115.7, 108.7, 77.8, 77.5, 77.2, 53.6, 37.7, 35.9, 30.3.

GC-MS (m/z): $[\text{M}+\text{H}]^+$ calc'd for $\text{C}_{15}\text{H}_{17}\text{NO}^+$: 227.1; found: 227.1

4-(2-phenylethyl-1-d)-3-(trifluoromethyl)pyridine (**15**)

Compound **15** was prepared from 4-chloro-3-(trifluoromethyl)pyridine and (2-bromoethyl)benzene according to procedure G.

¹H NMR (400 MHz, CDCl₃) δ 8.84 (s, 1H), 8.63 (d, *J* = 5.1 Hz, 1H), 7.35 – 7.26 (m, 2H), 7.25 – 7.13 (m, 4H), 3.16 – 3.04 (m, 2H), 2.99 – 2.81 (m, 2H).

¹³C NMR (101 MHz, CDCl₃) δ 152.9, 149.7 (d, *J* = 1.6 Hz), 147.3 (q, *J* = 6.3 Hz), 140.4, 128.8, 128.6, 126.7, 125.5, 125.0 (q, *J* = 30.3 Hz), 124.3 (q, *J* = 275.7 Hz), 36.9, 34.4.

GC-MS (m/z): [M+H]⁺ calc'd for C₁₄H₁₂F₃N⁺: 251.1; found: 251.2

5,6,7,8-tetrahydroisoquinoline (16) (commercially available)

References

1. Zhao, Y.; Chen, Y.; Newhouse, T. R. Allyl-Palladium Catalyzed α,β -Dehydrogenation of Carboxylic Acids via Enediolates. *Angew. Chem. Int. Ed.* **2017**, *56*, 13122.
2. Zhang, P.; Huang, D.; Newhouse, T. R. Aryl-Nickel-Catalyzed Benzylic Dehydrogenation of Electron-Deficient Heteroarenes. *J. Am. Chem. Soc.* **2020**, *142*, 1757.
3. Mierde, H. V.; Van Der Voort, P.; De Vos, D.; Verpoort, F. A Ruthenium-Catalyzed Approach to the Friedländer Quinoline Synthesis. *Eur. J. Org. Chem.* **2008**, 1625.
4. Fürstner, A.; Leitner, A.; Méndez, M.; Krause, H. Iron-Catalyzed Cross-Coupling Reactions. *J. Am. Chem. Soc.* **2002**, *124*, 13856.