Lectotypification of *Papaver cambricum* L. (Papaveraceae)

P. Pablo Ferrer-Gallego

Abstract

FERRER-GALLEGO, P. P. (2015). Lectotypification of *Papaver cambricum* L. (Papaveraceae). *Candollea* 70: 207-210. In English, English abstract. DOI: <http://dx.doi.org/10.15553/c2015v702a5>

The typification of the name *Papaver cambricum* L. (Papaveraceae) is discussed. This species was previously accepted in the genus *Meconopsis* Vig. as *Meconopsis cambrica* (L.) Vig. The protologue of the name and the original material are evaluated. A specimen from the Burser Herbarium (UPS-BURSER) is designated as the lectotype.

Keywords

PAPAVERACEAE – *Meconopsis* – *Papaver* – Lectotypification

Address of the author:

CIEF, Centro para la Investigación y la Experimentación Forestal, Servicio de Vida Silvestre, Generalitat Valenciana, Avda. Comarques del País Valencià, 114, 46930, Quart de Poblet, Valencia, Spain. E-mail: flora.cief@gva.es

Submitted on October 6, 2014. Accepted on June 9, 2015

Edited by P. Bungener

Introduction

The genus *Meconopsis* Vig. (Papaveraceae) comprises approximately 70 perennial monocarpic or polycarpic herbs, distributed primarily in southern central Asia (GREY-WILSON, 2014). The western European *M. cambrica* (L.) Vig. was long considered to be the only European representative of the genus. However, new molecular studies suggest that it does not belong to the Asian species of *Meconopsis* but should rather be accepted in *Papaver* L. (KADEREIT et al., 1997, 2011; CAROLAN et al., 2006). As *Meconopsis cambrica* was the original type of *Meconopsis*, the genus would have been considered as a synonym of *Papaver* and therefore left the Asian species without a generic name. GREY-WILSON (2012) further proposed the name *Meconopsis* for conservation with a proposed conserved type: *M. regia* G. Taylor. This proposal was strongly recommended by the Nomenclature Committee for Vascular Plants (APPLEQUIST, 2013).

Meanwhile, GREY-WILSON (2014) placed *Papaver cambricum* L. in the monotypical genus *Parameconopsis* Grey-Wilson.

However, the name *Papaver cambricum* L. has not yet been typified (JARVIS, 2007: 722). This paper proposes the designation of a lectotype for this name based on the protologue and consultation of Linnaeus's original material.

Typification

The protologue of *P. cambricum* (LINNAEUS, 1753: 508) presents four synonyms. The first synonym is cited from LINNAEUS (1738: 201) and ROYEN (1740: 479) “*Papaver foliis pinnatis, fructu acuminato*”, the second from DILLENUS (1732: 300, t. 223, f. 290) “*Papaver cambricum perenne, flore sulphureo*”, the third from MORISON (1680: 279, s. 3, t. 14, f. 12) “*Argemone cambro-britannica lutea, capite longiore glabro*”, and the fourth from BAUHIN (1620: 92, 1623: 171) “*Papaver erraticum pyrenaicum, flore flavo*”. “Burs. IX: 45” is the citation of a specimen in UPS-BURSER (Herb. Burser IX: 45), which is therefore a syntype. Two of the synonyms are accompanied by illustrations (MORISON, 1680; DILLENUS, 1732), which are also original material.

Linnaeus's original material additionally includes two uncited specimens (JARVIS, 2007: 723). At LINN, the specimen “Herb. Linn. No. 669.9” bears the annotation “7 cambricum” by Linnaeus, explicitly referring to the number of the species account of *P. cambricum* in LINNAEUS (1753: 508) (see STEARN, 1957; TURLAND & JARVIS, 1997; TURLAND, 2006; JARVIS, 2007: 41–46, 397). The specimen is a stem with leaves, a flower and a flower bud, but no fruit.

The sheet preserved in the Clifford Herbarium, “Herb. Clifford: 201”, *Papaver* 4 (BM000628706) is annotated “*Papaver erraticum pyrenaicum flore flavo* C.B.P. 171 / *Papaver foliis pinnatis, fructu acuminato* H. Cl. IV. p. 201”. The specimen is a stem with leaves and three flowers, but no fruit.

Finally, the syntype in UPS-BURSER bears two

fragments, one with a flower, the other with a flower bud, and a label with the annotation “*Papaver erraticum pyrenaicum flore flavo* Bauh. / In Pyrenaeis” (see SAVAGE, 1937: 27). No further original material have been found in any of the other Linnaean and Linnaean-linked herbaria.

The specimen in UPS-BURSER has precedence in lectotype designation over the uncited specimens and cited illustrations. It is therefore the obligate lectotype and must be designated as such. Fortunately it is in a good state of preservation, and it conforms with the other Linnaean specimens and illustrations and with the traditional and current use of the names *P. cambricum* and *Meconopsis cambrica*.

***Papaver cambricum* L., Sp. Pl.: 508. 1753.**

≡ *Meconopsis cambrica* (L.) Vig., Hist. Nat. Pavots: 48. 1814.

≡ *Parameconopsis cambrica* (L.) Grey-Wilson, Meconopsis: 367. 2014.

Lectotypus (here designated): Herb. Burser IX: 45 (UPS-BURSER!) (Fig. 1).

Acknowledgements

The author thanks Dr. Mats Hjertson (Museum of Evolution, Botany Section Uppsala University, Sweden) for his help as well as an anonymous reviewer.

References

- APPLEQUIST, W. L. (2013). Report of the Nomenclature Committee for Vascular Plants: 65. *Taxon* 62: 1315–1326.
- BAUHIN, C. (1620). *Prodromus theatri botanici*. Francofurti ad Moenum.
- BAUHIN, C. (1623). *Pinax theatri botanici*. Basileae.
- CAROLAN, J. C., I. L. I. HOOK, M. W. CHASE, J. W. KADEREIT & T. R. HODKINSON (2006). Phylogenetics of *Papaver* and related genera based on DNA sequences from ITS nuclear ribosomal DNA and plastid trnL intron and trnL-F intergenic spacers. *Ann. Bot. (Oxford)* 98: 141–155.
- DILLENUS, J. J. (1732). *Hortus Elthamensis*. Londini.
- GREY-WILSON, C. (2012). (2061) Proposal to conserve the name *Meconopsis* (Papaveraceae) with a conserved type. *Taxon* 61: 473–474.
- GREY-WILSON, C. (2014). *The genus Meconopsis: Blue poppies and their relatives*. Royal Botanic Gardens, Kew.
- JARVIS, C. (2007). *Order out of chaos: Linnaean plant names and their types*. Linnean Society of London & Natural History Museum, London.
- KADEREIT, J. W., C. D. PRESTON & J. F. VALTUEÑA (2011). Is Welsh Poppy, *Meconopsis cambrica* (L.) Vig. (Papaveraceae), truly a *Meconopsis*? *New J. Bot.* 1: 80–87.

Fig. 1. – Lectotype of *Papaver cambricum* L.
[Herb. Burser IX: 45, UPS-BURSER] [© Herbarium UPS. Reproduced with permission]

- KADEREIT, J. W., A. E. SCHWARZBACH & K. B. JORK (1997). The phylogeny of *Papaver* s.l. (Papaveraceae): polyphyly or monophyly? *Pl. Syst. Evol.* 204: 75-98.
- LINNAEUS, C. (1738). *Hortus Cliffortianus*. Amsterdam.
- LINNAEUS, C. (1753). *Species Plantarum*. Holmiae.
- MORISON, R. (1680). *Plantarum historiae universalis Oxoniensis pars secunda*. Oxford.
- ROYEN, A. VAN (1740). *Florae Leydensis prodromus, exhibens plantas quae in horto academico Lugduno-Batavo aluntur*. Lugduni Batavorum.
- SAVAGE, S. (1937). *Caroli Linnaei determinationes in Hortum Siccum Joachimi Burseri. The text of the manuscript in the Linnaean Collections*. Linnean Society, London.
- STEARN, W. T. (1957). An introduction to the *Species Plantarum* and cognate botanical works of Carl Linnaeus. In: LINNAEUS, C., *Species Plantarum, a facsimile of the first edition, 1753*. Vol. 1. Ray Society, London.
- TURLAND, N. J. (2006). Lectotypification of *Campanula saxatilis*, *Phyteuma pinnatum* and *Verbascum arcturus*, Linnaean names of three taxa endemic to Crete. *Willdenowia* 36: 303-309.
- TURLAND, N. J. & C. E. JARVIS (1997). Typification of Linnaean specific and varietal names in the Leguminosae (Fabaceae). *Taxon* 46: 457-485.