

# Goldbach's Conjecture — A Route to the Inconsistency of Arithmetic

Ralf Wüsthofen

**Abstract.** This paper proves an inconsistency in Peano arithmetic (PA). We express a strengthened form of the strong Goldbach conjecture by using a specific set that varies according to whether the conjecture or its negation holds. We show that, on the other hand, this set remains unchanged whether the conjecture holds or not. This causes a contradiction.

**Notations.** Let  $\mathbb{N}$  denote the natural numbers starting from 1, let  $\mathbb{N}_n$  denote the natural numbers starting from  $n > 1$  and let  $\mathbb{P}_3$  denote the prime numbers starting from 3.

Strengthened strong Goldbach conjecture (SSGB): *Every even integer greater than 6 can be expressed as the sum of two different primes.*

**Theorem.** *PA is contradictory, i.e. the statement FALSE can be derived.*

*Proof.* We define the set  $S_g := \{ (pk, mk, qk) \mid k, m \in \mathbb{N}; p, q \in \mathbb{P}_3, p < q; m = (p + q) / 2 \}$ .

SSGB is equivalent to saying that every integer  $n \geq 4$  is the arithmetic mean of two different odd primes and so it is equivalent to saying that all integers  $n \geq 4$  appear as  $m$  in a middle component  $mk$  of  $S_g$ . So, by the definition of  $S_g$  we have

$$\begin{aligned} \text{SSGB} &\Leftrightarrow \forall n \in \mathbb{N}_4 \exists (pk, mk, qk) \in S_g \quad n = m \\ \neg\text{SSGB} &\Leftrightarrow \exists n \in \mathbb{N}_4 \forall (pk, mk, qk) \in S_g \quad n \neq m. \end{aligned}$$

The set  $S_g$  has the following two properties.

First, the whole range of  $\mathbb{N}_3$  can be expressed by the triple components of  $S_g$  ("covering"), because every integer  $x \geq 3$  can be written as some  $pk$  with  $k = 1$  when  $x$  is prime, as some  $pk$  with  $k \neq 1$  when  $x$  is composite and not a power of 2, or as  $(3 + 5)k / 2$  when  $x$  is a power of 2;  $p \in \mathbb{P}_3, k \in \mathbb{N}$ . So we have

$$(C) \quad \forall x \in \mathbb{N}_3 \exists (pk, mk, qk) \in S_g \quad x = pk \vee x = mk.$$

A few examples of the covering:

$x = 19$ : (**19·1**, 21·1, 23·1), (**19·1**, 60·1, 101·1)

$x = 27$ : (**3·9**, 7·9, 11·9)

$x = 38$ : (**19·2**, 21·2, 23·2)

$x = 42$ : (**3·14**, 5·14, 7·14), (**7·6**, 9·6, 11·6)

$x = 4096$ : (3·1024, **4·1024**, 5·1024)

$x = 10000$ : (**5·2000**, 6·2000, 7·2000).

Second, according to the statement SSGB, all pairs  $(p, q)$  of distinct odd primes are used in the definition of the set  $S_g$  (“maximality”). So we have

**(M)**  $\forall p, q \in \mathbb{P}_3, p < q \quad \forall k \in \mathbb{N} \quad (pk, mk, qk) \in S_g$ , where  $m = (p + q) / 2$ .

$\neg(C)$  immediately implies  $\neg$ SSGB since an  $n \geq 4$  that is different from all  $S_g$  triple components  $pk$  and  $mk$  is in particular different from all  $m$  in  $S_g$ . So the property (C) excludes the possibility that  $\neg$ SSGB holds because of  $\neg(C)$ .

The property (M) excludes the possibility that there is an  $n \geq 4$  different from all  $m$ , i.e.  $\neg$ SSGB, where  $n$  is the arithmetic mean of a pair of primes not used in  $S_g$ . So (M) excludes the possibility that  $\neg$ SSGB holds due to a missing prime number pair. The proof would no longer be possible if we left out any prime number pair in the formulation of SSGB and  $S_g$ .

The basic idea is now the following.

*There are two possibilities for  $S_g$ , exactly one of which must occur: Either there is an  $n \in \mathbb{N}_4$  in addition to all the numbers  $m$  defined in  $S_g$  or there is not. The latter is equivalent to SSGB and the former is equivalent to  $\neg$ SSGB.*

*Since, due to (M), an  $n \geq 4$  different from all  $m$  cannot be the arithmetic mean of a pair of primes not used in  $S_g$  and since, due to (C), this  $n$  equals a component of some  $S_g$  triple that exists by definition, the covering of  $\mathbb{N}_3$  by the  $S_g$  triples in the case  $n$  exists ( $\neg$ SSGB) is equal to that in the case  $n$  does not exist (SSGB). This causes a contradiction because in the case SSGB the numbers  $m$  defined in  $S_g$  take all integer values  $x \geq 4$  whereas in the case  $\neg$ SSGB they don't.*

The following steps are independent of the choice of  $n$  if, in the case of  $\neg\text{SSGB}$ , there is more than one that is different from all  $m$ . For example, the minimal such  $n$  works.

We split  $S_g$  into two complementary subsets in the following way. For any  $y \in \mathbb{N}_3$ , we write

$S_g = S_{g+(y)} \cup S_{g-(y)}$ , with

$S_{g+(y)} := \{ (pk, mk, qk) \in S_g \mid \exists k' \in \mathbb{N} \quad pk = yk' \vee mk = yk' \vee qk = yk' \}$

$S_{g-(y)} := \{ (pk, mk, qk) \in S_g \mid \forall k' \in \mathbb{N} \quad pk \neq yk' \wedge mk \neq yk' \wedge qk \neq yk' \}$ .

We define

$S_1 := \{ (pk, mk, qk) \in S_g \mid \text{SSGB} \wedge (M) \}$

$S_2 := \{ (pk, mk, qk) \in S_g \mid \neg\text{SSGB} \wedge ((C) \wedge (M)) \}$ .

Under the assumption  $\neg\text{SSGB}$  there is an  $n \in \mathbb{N}_4$  as described above, whereas under the assumption  $\text{SSGB}$  there is no such  $n$ . Then, since  $\text{SSGB} \Rightarrow (C)$ ,

**(1.1)**  $\forall y \in \mathbb{N}_3 \quad \text{SSGB} \Rightarrow ( S_1 = S_{g+(y)} \cup S_{g-(y)} \vee \neg(M) )$

$\wedge$

**(1.2)**  $\neg\text{SSGB} \Rightarrow ( S_2 = S_{g+(n)} \cup S_{g-(n)} \vee \neg(C) \vee \neg(M) )$ .

Since  $\neg(C)$  and  $\neg(M)$  are both ruled out and since  $S_{g+(n)} \cup S_{g-(n)}$  is independent of  $n$ , we get

**(1.1')**  $\forall y \in \mathbb{N}_3 \quad \text{SSGB} \Rightarrow S_1 = S_{g+(y)} \cup S_{g-(y)}$

$\wedge$

**(1.2')**  $\forall y \in \mathbb{N}_3 \quad \neg\text{SSGB} \Rightarrow S_2 = S_{g+(y)} \cup S_{g-(y)}$ .

Now, we will make use of the following principle.

If two sets of (possibly infinitely many)  $x$ -tuples are equal, then the sets of their corresponding  $i$ -th components are equal;  $1 \leq i \leq x$ .

To this end, for each  $k \in \mathbb{N}$  we define

$$M_1(k) := \{ mk \mid (pk, mk, qk) \in S_1 \}$$

$$M_2(k) := \{ mk \mid (pk, mk, qk) \in S_2 \}.$$

Then, applying the principle above to the middle component of the triples  $(pk, mk, qk)$ ,  $((1.1') \wedge (1.2'))$  implies

$$(2.1) \quad \forall k \in \mathbb{N} \quad \forall y \in \mathbb{N}_3 \quad \text{SSGB} \Rightarrow M_1(k) = \{ mk \mid (pk, mk, qk) \in S_{g^+(y)} \cup S_{g^-(y)} \}$$

$\wedge$

$$(2.2) \quad \forall k \in \mathbb{N} \quad \forall y \in \mathbb{N}_3 \quad \neg\text{SSGB} \Rightarrow M_2(k) = \{ mk \mid (pk, mk, qk) \in S_{g^+(y)} \cup S_{g^-(y)} \}.$$

Setting  $M_1 := M_1(1)$  and  $M_2 := M_2(1)$ , we get

$$(2.1') \quad \forall y \in \mathbb{N}_3 \quad \text{SSGB} \Rightarrow M_1 = \{ m \mid (p, m, q) \in S_{g^+(y)} \cup S_{g^-(y)} \}$$

$\wedge$

$$(2.2') \quad \forall y \in \mathbb{N}_3 \quad \neg\text{SSGB} \Rightarrow M_2 = \{ m \mid (p, m, q) \in S_{g^+(y)} \cup S_{g^-(y)} \}.$$

Since for every  $y \in \mathbb{N}_3$   $S_{g^+(y)} \cup S_{g^-(y)}$  equals  $S_g$  by definition, for every  $y \in \mathbb{N}_3$   $\{ m \mid (p, m, q) \in S_{g^+(y)} \cup S_{g^-(y)} \}$  equals the set  $X := \{ m \mid (p, m, q) \in S_g \}$ . So, from  $((2.1') \wedge (2.2'))$  we obtain

$$(3) \quad (\text{SSGB} \Rightarrow M_1 = X) \quad \wedge \quad (\neg\text{SSGB} \Rightarrow M_2 = X).$$

The set  $X$  is a free variable in (3) that is either equal to  $\mathbb{N}_4$  or to some non-empty proper subset  $Y$  of  $\mathbb{N}_4$ .

Now, we make use of the following rule.

Let  $P = P(A)$  be a proposition that depends on a set  $A$ . Then, for any set  $B$ ,

( we have a proof of  $P(A) \wedge$  we have a proof of  $A = B$  )  $\Rightarrow$  we have a proof of  $P(B)$ .

In the special case that  $A$  is a free variable that is replaced by the value  $B$ , the above conjunct ( we have a proof of  $A = B$  ) is trivially true.

Since the set  $X$  is a free variable in (3) and since we have a proof of (3), we can apply the above rule with  $P = (3)$ . If  $X = \mathbb{N}_4$  we use the rule with  $A = X$  and  $B = \mathbb{N}_4$ , and if  $X = Y$  we use it with  $A = X$  and  $B = Y$ . Then, since either  $X = \mathbb{N}_4$  or  $X = Y$ , from (3) we obtain

**(3.1)** we have a proof of (  $SSGB \Rightarrow M_1 = \mathbb{N}_4 \quad \wedge \quad \neg SSGB \Rightarrow M_2 = \mathbb{N}_4$  )

∨

**(3.2)** we have a proof of (  $SSGB \Rightarrow M_1 = Y \neq \mathbb{N}_4 \quad \wedge \quad \neg SSGB \Rightarrow M_2 = Y \neq \mathbb{N}_4$  ).

This implies

**(3.1')** ( we have a proof of (  $SSGB \Rightarrow M_1 = \mathbb{N}_4$  )

$\wedge$   
we have a proof of (  $\neg SSGB \Rightarrow M_2 = \mathbb{N}_4$  ) )

∨

**(3.2')** ( we have a proof of (  $SSGB \Rightarrow M_1 = Y \neq \mathbb{N}_4$  )

$\wedge$   
we have a proof of (  $\neg SSGB \Rightarrow M_2 = Y \neq \mathbb{N}_4$  ) ).

Now, we will establish a contradiction to  $((3.1') \vee (3.2'))$ .

Under the assumption  $SSGB$  the set  $X = \{ m \mid (p, m, q) \in S_g \}$  is equal to  $\mathbb{N}_4$  and under  $\neg SSGB$  it is equal to  $Y \neq \mathbb{N}_4$ . Therefore,

**(4.1)** we have a proof of  $(SSGB \Rightarrow M_1 = \mathbb{N}_4)$

$\wedge$

**(4.2)** we have a proof of  $(\neg SSGB \Rightarrow M_2 = Y \neq \mathbb{N}_4)$ .

Then,  $((3.1') \vee (3.2'))$  together with  $((4.1) \wedge (4.2))$  implies

**(5.1)** we have a proof of  $(\neg SSGB \Rightarrow M_2 = \mathbb{N}_4)$

$\vee$

**(5.2)** we have a proof of  $(SSGB \Rightarrow M_1 = Y \neq \mathbb{N}_4)$ .

Because of  $((4.1) \wedge (4.2))$  and because

$SSGB \Rightarrow M_2 = \{ \} \neq \mathbb{N}_4$

and

$\neg SSGB \Rightarrow M_1 = \{ \} \neq Y$ ,

we have a proof that  $(M_2 = \mathbb{N}_4)$  is false and we have a proof that  $(M_1 = Y \neq \mathbb{N}_4)$  is false.

So,  $((5.1) \vee (5.2))$  yields

**(6.1)** we have a proof of  $\text{SSGB}$

$\vee$

**(6.2)** we have a proof of  $\neg\text{SSGB}$ .

Since we have neither a proof of  $\text{SSGB}$  nor of  $\neg\text{SSGB}$ , both (6.1) and (6.2) are false.

Therefore, we obtain  $(\text{FALSE} \vee \text{FALSE})$  and thus  $\text{FALSE}$ .

□