



Antimicrobial and Anti-inflammatory Effects of Two Different Marine Red Algae Species Collected from Quseir, the Red Sea, Egypt

Ibraheem B. M. Ibraheem¹, Basma E. Abd Elaziz¹, Abeer Moawad²,
Hossam M. Hassan^{2,3*}, Waleed A. Mohamed⁴ and Neveen Abdel-Raouf¹

¹Department of Botany and Microbiology, Faculty of Science, Beni-Suef University, 62514, Egypt.

²Department of Pharmacognosy, Faculty of Pharmacy, Beni-Suef University, 62514, Egypt.

³Department of Pharmacognosy, Faculty of Pharmacy, Nahda University, BeniSuef, 62514, Egypt.

⁴Department of Biochemistry, Kasr El-Eini Teaching Hospitals, Cairo University, Egypt.

Authors' contributions

This work was carried out in collaboration between all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJOB/2017/32964

Editor(s):

(1) Xing Li, Division of Biomedical Statistics and Informatics, Department of Health Sciences Research, Mayo Clinic College of Medicine, USA.

Reviewers:

(1) Irshad Ahmad, King Fahd University of Petroleum and Minerals (KFUPM), 31261 Dhahran, Saudi Arabia.

(2) Kusmiati, Research Center For Biotechnology-Indonesian Institute of Sciences (Lipi), Indonesia.

(3) Marisa Alves Nogueira Diaz, Federal University of Vicosa, Brazil.

Complete Peer review History: <http://www.sciencedomain.org/review-history/19064>

Original Research Article

Received 25th March 2017

Accepted 26th April 2017

Published 15th May 2017

ABSTRACT

Aims: This study aimed to analyze the bioactivities of two different species of marine red algae crude extracts; *Acanthophora spicifera* (M.Vahl) Borgese and *Digenea simplex* (Wulfen) C.Agardh.

Place and Duration of Study: Sample: Quseir city, Red Sea, Egypt at (April, 2015), extraction and biological study; pharmacognosy department, summer 2015, Kasr El-Eini Teaching Hospitals, Cairo University, Egypt.

Methodology: The crude extract screened against some human pathogenic bacteria including ten Gram positive, ten Gram negative strains and ten unicellular and filamentous fungi. The antimicrobial activity was done through the agar well diffusion method. The anti-inflammatory effect of the crude extract on inflamed liver cells was evaluated by measuring SOD, MDA, catalase, GSH, Il-6 and TNF-alpha in the serum of treated Wister albino rats compared to positive and negative control groups.

*Corresponding author: E-mail: abuh20050@yahoo.com;

Results: The antimicrobial study showed that both *Acanthophora spicifera* and *Digenea simplex* extracts showed the highest zone of inhibition against *Streptococcus agalactiae* with clear zones (22.5±0.58 mm and 23.1±0.58 mm) as Gram positive bacteria compared to Ampicillin as positive control. But with Gram negative bacteria *Acanthophora spicifera* showed highest zone of inhibition against *Pseudomonas aeruginosa* (22.3±2.1 mm) and *Digenea simplex* showed highest zone of inhibition against *Serratia marcescens* (24.1±0.58 mm) compared to Gentamycin as positive control. Finally, the effect of both *Acanthophora spicifera* and *Digenea simplex* against fungi showed highest zone of inhibition (23.8±0.63 mm and 22.4±2.1 mm) against *Geotricum candidum*. On the other hand results revealed that all methanol extracts had equally potent anti-inflammatory effects on inflamed liver rats as it observed significant decreased in the levels of SOD, Catalase and GSH. Also observed that significant increase in the inflamed liver state of MDA, IL-6 and TNF-Alpha After treatment of inflamed liver rats with crude extracts observed that the difference in the parameters levels relatively equal to the normal control results.

Conclusion: *Acanthophora spicifera* (M.Vahl) Borgese and *Digenea simplex* (Wulfen) C.Agardh might be a good source of anti-inflammatory and antibacterial compounds activity.

Keywords: Marine red macro-algae; antimicrobial; anti-inflammation.

1. INTRODUCTION

Marine macro algae are one of nature's most biologically active resources, as they hold a wealth of bioactive compounds. Many compounds isolated from marine macroalgae have demonstrated various biological activities, such as antibacterial activity, antioxidant potential, anti-inflammatory properties, anticoagulant activities, antiviral activities, apoptotic activities, and prebiotic activity [1-6]. Marine macro algae are commonly classified into three main groups based on their pigmentation; Phaeophyta (brown algae), Chlorophyta (green algae), and Rhodophyta (red algae). Macroalgae was considered as a rich source of dietary fiber, minerals, lipids, proteins, omega-3 fatty acids, essential amino acids, polysaccharides, and vitamins A, B, C, and E. Various bioactive compounds from marine organisms have been experimentally tested to comprehensively study the biological effects of recently developed drugs [7]. Studies on the bioactivities of marine algae have revealed numerous health-promoting effects, including anti-oxidative, anti-inflammatory, antimicrobial, and anti-cancer effects. Seaweeds are an excellent source of components such as polysaccharides, tannins, flavonoids, phenolic acids, bromophenols, and carotenoids have exhibits different biological activities, relaying on their solubility and polarity, different solvents shows different antimicrobial activity. Chemical compounds should be extracted from different seaweeds in order to optimize their antibacterial and antifungal activity by selecting the best solvent system [8].

Oxidative stress plays important role in endothelial dysfunction because the molecular disintegration in reactive oxygen species (ROS) metabolites which plays a main reason in the pathogenesis of it by a loss of nitric oxide (NO) bioavailability [9], lung disease [10], gastrointestinal dysfunction [11], and atherosclerosis [12], all of which involve inflammatory reactions. Many marine natural products that contain antioxidants were known to have anti-inflammatory effects [13-15]. This study was carried out to figure the antimicrobial and anti-inflammatory effects of 2 different algal species. Finally marine algae considered as the main of the food chain in the areas overlooking the ocean and enhance about more than two-thirds of the world's populations. In addition to its importance, algae are responsible for about half of global photosynthetic activity [16].

2. MATERIALS AND METHODS

2.1 Collection and Identification of Marine Macroalgae

The fresh algal species were collected from the inter-tidal region of Quseir city (Fig. 1) which located on the west coastal area of Red Sea shore between longitude 34° 17'E and latitude 26° 06'N during the spring year 2015. Collected sample was immediately brought to the laboratory in new plastic bags containing pond water to prevent evaporation. Algal material was washed thoroughly with tap water and distilled water to remove extraneous materials and shade-dried for 5 days and oven dried at 60°C



Fig. 1. Map show the studied area [Quseir city, Red Sea, Egypt]

until constant weight was obtained, then was grind into fine powder using electric mixer and stored at 4°C for future use. Algal species were identified according to [17-20].

2.2 Extraction of Selected Algal Species

Powdered marine algae (500 g, each) were extracted with 70% methanol [21], because some of the active components in the crude extracts are polar and thus dissolved readily in the methanol (polar solvent), by percolation till exhaustion. The alcoholic extract in each case was evaporated under reduced pressure to obtain a semisolid residue.

2.3 Antimicrobial Activity (the Microorganisms Used are from the RCMB Collection)

The hydro alcoholic extracts of the selected species were investigated for their antimicrobial activities using Agar well diffusion and Muller Henton against gram positive and gram negative bacteria and used Sab. dextrose agar against some fungi. All tested microorganisms were kindly supplied from Biotechnology Research

Center, Al-Azhar University (for boys), Cairo, Egypt.

2.3.1 Gram positive bacteria (the microorganisms used are from the RCMB collection)

Staphylococcus aureus (RCMB 010027), *Staphylococcus epidermidis* (RCMB 010024), *Streptococcus sanguis*(RCMB 01001 71-3), *Streptococcus pyogenes* (RCMB 01001 74-2), *Streptococcus agalactiae*(RCMB 01001 73-2), *Bacillus subtilis* (RCMB 01001 69-3), *Enterococcus faecalis* (RCMB 01001 54-2), *Corynebacterium diphtheriae* (RCMB 01001 26-7), *Micrococcus luteus* (RCMB 01001 76-9), *Methicillin-resistant Staphylococcus aureus MRSA* (RCMB 01001 94-5).

2.3.2 Gram negative bacteria (the microorganisms used are from the RCMB collection)

Escherichia coli (RCMB 01002 52-6), *Proteus mirabilis* (RCMB 01002 54-2), *Acinetobacter baumannii* (RCMB 01002 82-9), *Klebsiella pneumonia* (RCMB 01002 23-5), *Pseudomonas aeruginosa* (RCMB 01002 43-5), *Serratia*

plymuthica (RCMB 01002 75-3), *Serratia marcescens* (RCMB 01002 75b-8), *Salmonella tophi* (RCMB 01002 15-4), *Enterobacter cloacae* (RCMB 01002 64-5), *Shigella dysenteriae* (RCMB 01002 41-8).

2.3.3 Unicellular fungi and filamentous fungi (the microorganisms used are from the RCMB collection)

Aspergillus fumigatus (RCMB 02568), *Syncephalastrum racemosum* (RCMB 05922), *Geotricum candidum* (RCMB 05097), *Candida albicans* (RCMB 05036), *Aspergillus niger* (RCMB 02724), *Cryptococcus neoformans* (RCMB 05642), *Candida tropicalis* (RCMB 05239), *Penicillium expansum* (RCMB 01924), *Microsporium canis* (RCMB 0834), *Trichophyton mentagrophytes* (RCMB 0925).

2.3.4 Methods of measuring antimicrobial activity

2.3.4.1 Well-diffusion method for anti-bacterial activity: (Mueller Henton)

The solution of 50 mg/ml of each sample in DMSO (Dimethyl sulphoxide) was prepared for testing against bacteria. Centrifuged pellets of bacteria from 24 h old culture containing approximately 104-106 CFU/ml (Colony forming unit per ml) were spread on the surface of Nutrient agar (type tone 1%, Yeast extract 0.5%, agar 1%, 100 ml of distilled water, PH 7.0) which autoclaved under 121°C for at least 20 min. Wells were created in medium with the help of a sterile metallic bores and then cooled down to 45°C. The activity was determined by measuring the diameter of the inhibition zone (in mm). 100 µl of the tested samples (100 mg/ml) were loaded into the wells of the plates. All samples were prepared in DMSO. DMSO was loaded as control. The plates were kept for incubation at 37°C for 24 h and then the plates were examined for the formation of zone of inhibition. Each inhibition zone was measured three times by caliper to get an average value. The test was performed three times for each bacterium culture. Ampicillin and Gentamycin were used as antibacterial standard drugs [22].

2.3.4.2 Well-diffusion method for anti-fungal activity: (Sabourad dextrose)

The antifungal activity was investigated by agar well diffusion method by the following procedure: Sabourad dextrose agar plates: A homogenous

mixture of glucose-peptone-agar (40: 10: 15) was sterilized by autoclaving at 121°C for 20 min. The sterilized solution (25 ml) was poured in each sterilized petridish in laminar flow and left for 20 min to form the solidified sabourad dextrose agar plate. These plates were inverted and kept at 30°C in incubator to remove the moisture and check for any contamination. Antifungal assay: Fungal strain was grown in 5 ml Sabourad dextrose broth (glucose: peptone; 40: 10) for 3-4 days to achieve 10⁵ CFU/ml cells. The fungal culture (0.1 ml) was spread out uniformly on the Sabourad dextrose agar plates. Now small wells of size (4 mm×20 mm) were cut into the plates with the help of well cutter and bottom of the wells were sealed with 0.8% soft agar to prevent the flow of test sample at the bottom of the well. 100 µl of the tested samples (10mg/ml) were loaded into wells of the plates. All samples was prepared in dimethyl sulfoxide (DMSO), DMSO was loaded as control. The plates were kept for incubation at 30°C for 3-4 days and then the plates were examined for the formation of zone of inhibition. Each inhibition zone was measured three times by caliper to get an average value. The test was performed three times for each fungus. Amphotericin B was used as antifungal standard drug [22].

2.4 Anti-inflammatory Activity

Animal: Wister albino rats weighing between (140-200 g) were obtained from animal house of National Research Center, Dokki – Giza, Egypt. They were housed in well protected polypropylene cages and maintained in standard laboratory conditions in an air conditioned area at 25±2°C in 10:14 hr light dark cycle and allowed to take standard laboratory feed and tap water [23]. The study was approved by Institutional Animal Ethics Committee of National Research Center that the animal doesn't suffering at any stage of experiment and maintained in accordance with the Guide for the Care and Use of Laboratory Animals. The inflamed liver was induced in rats according to the described method [24]. The animals were divided into three groups consisted of six animals; Normal control (A), Diseased or inflamed Liver cancer (positive control receiving saline) (B), Inflamed liver rats were treated by injection with methanol extract of the first extract (*Acanthopora spicifera*) (C) and the second extract (*Digenea simplex*) (D). The methanol extract (0.1 g of each sample, separately) mixed or dissolved with 1 ml of phosphate buffer and every day injected 100 µm to each rat for 15

days. After 15 days, rats were sacrificed and blood samples were collected by puncture the sublingual vein in clean and dry test tube. Allow clotting for 10 minutes before centrifuging at 3000 rpm for serum separation. The separated serum was stored at -80°C for further determinations of the following tests: SOD, MDA, catalase, GSH, Il-6 and TNF-alpha. The anti-inflammatory activity was observed on Wister albino rats after injection by the extracts of the marine algal species (Table 4) by measuring the parameters in the normal and diseased states.

3. RESULTS AND DISCUSSION

3.1 Collection and Identification

Isolated algal species: Two algal samples of red algae *Acanthophora spicifera* (M.Vahl) Borgese (Fig. 2) and *Digenea simplex*(Wulfen) C.Agardh (Fig. 3) were collected from Red Sea, Egypt (Quseir city).




Fig. 2. *Acanthophora spicifera*




Fig. 3. *Digenea simplex*

3.2 Antimicrobial Activity

Results summarized in the Tables (1-3) showed the antimicrobial screening of the crude extracts of two macro algae against some human pathogenic bacteria as well as some unicellular and filamentous fungi. Both *A. spicifera* and *D.*

simplex extracts showed similar potent inhibitory growth activities against three Gram +ve bacteria [*Streptococcus agalactiae*, *Streptococcus pyogenes* and *Streptococcus sanguis*] with zone of inhibition ranging from $[23.1\pm 0.58$ to 20.6 ± 0.63 mm] and showed moderate activities with [*Corynebacterium diphtheriae*, *Bacillus subtilis* and *Staphylococcus aureus*] with inhibition zones ranging from $[20.1\pm 1.5$ to 16.3 ± 2.1 mm]. Also the crude extracts were found to be more active than the positive control Ampicillin, (22.3 ± 1.5 mm), against *Streptococcus agalactiae* which showing inhibition zone $[22.5\pm 0.58$ mm with *A. spicifera* and 23.1 ± 0.58 mm with *D. simplex*]. *Staphylococcus epidermidis*, *Enterococcus faecalis*, *Micrococcus luteus* and *Methicillin-resistant Staphylococcus aureus* are resistance to the activity of the algae compared to the positive control Ampicillin and Vancomycin, this was on the contrary of previously published activity of *A.spicifera* against MRSA [25]. On the other hand the two algal species were showing activities against seven Gram -ve bacteria [*Escherichia coli*, *Proteus mirabilis*, *Klebsiella pneumonia*, *Pseudomonas aeruginosa*, *Serratia plymuthica*, *Serratia marcescens* and *Enterobacter cloacae*] with zones of inhibition ranging from $[17.2\pm 1.5$ to 24.1 ± 0.58 mm] similar to previously reported [26]. Both algal extracts showed higher selective activity against *Pseudomonas aeruginosa* with zone of inhibition $[22.3\pm 2.1$ mm with *A. spicifera* and 22.9 ± 0.63 mm with *D. simplex*] compared to the positive control Gentamycine (20.6 ± 1.5 mm) and against *Serratia marcescens* with zone of inhibition $[22.3\pm 1.5$ mm with *A. spicifera* and 24.1 ± 0.58 mm with *D. simplex*] compared to the same positive control with (20.4 ± 0.58 mm of inhibition zone). *Acinetobacter baumannii*, *Salmonella typhi* and *Shigella dysenteriae* are resistance to the effect of the selected algae compared to the positive control Gentamycin. Previous results were different from the reported one that showed the crude extract of *Digenea simplex* as a weak growth inhibition zones ranging from 6 to 8 mm against *Staphylococcus aureus*, *Enterococcus faecalis* and *Pseudomonas aeruginosa* [27]. Discussing the antifungal activity, the algal extracts were found to be active on the selected species of filamentous fungi against six fungi [*Aspergillus fumigatus*, *Syncephalastrum racemosum*, *Geotricum candidum*, *Aspergillus niger*, *Candida tropicalis* and *Micro sporumcanis*] with zones of inhibition ranging from $[18.6\pm 0.63$ to 23.8 ± 0.63 mm]. *Geotricum candidum* and *Aspergillus niger* are the most sensitive fungal

Table 1. Antibacterial activity against gram + bacteria from red algae extracts (zone of inhibition in mm)

Sample tested microorganisms	<i>Acanthophora spicifera</i>	<i>Digenea simplex</i>	St.
Gram positive bacteria			Ampicillin
<i>Staphylococcus aureus</i> (RCMB 010027)	16.3±2.1 mm	17.1±2.1 mm	22±1.0 mm
<i>Staphylococcus epidermidis</i> (RCMB 010024)	NA	NA	23±1.0 mm
<i>Streptococcus sanguis</i> (RCMB 01001 71-3)	20.6±0.63 mm	21.2±0.63 mm	21.7±1.5 mm
<i>Streptococcus pyogenes</i> (RCMB 01001 74-2)	21.4±1.2 mm	21.6±1.2 mm	22.7±1.5 mm
<i>Streptococcus agalactiae</i> (RCMB 01001 73-2)	22.5±0.58 mm	23.1±0.58 mm	22.3±1.5 mm
<i>Bacillus subtilis</i> (RCMB 01001 69-3)	18.3±0.63 mm	19.3±1.2 mm	25.3±1.5 mm
<i>Enterococcus faecalis</i> (RCMB 01001 54-2)	NA	NA	19.3±0.58 mm
<i>Corynebacterium diphtheria</i> (RCMB 01001 26-7)	19.3±1.2 mm	20.1±1.5 mm	20±1.0 mm
<i>Micrococcus luteus</i> (RCMB 01001 76-9)	NA	NA	19.6±1.5 mm
Methicillin-resistant microorganisms			Vancomycine
<i>Methicillin-resistant Staphylococcus aureus</i> MRSA (RCMB 01001 94-5)	NA	NA	21.6±2.1 mm

RCMB: Regional Center of Mycology and Biotechnology Antimicrobial unit test organism.

NA: No activity

Table 2. Antibacterial activity against gram - bacteria from red algae extracts (zone of inhibition in mm)

Sample tested microorganisms	<i>Acanthophora spicifera</i>	<i>Digenea simplex</i>	St.
Gram negative bacteria			Gentamycine
<i>Escherichia coli</i> (RCMB 01002 52-6)	18.9 ± 1.2 mm	19.1± 1.2 mm	20.3± 0.85 mm
<i>Proteus mirabilis</i> (RCMB 01002 54-2)	17.2± 1.5 mm	17.8± 1.5 mm	21.2± 1.2 mm
<i>Acinetobacter baumannii</i> (RCMB 01002 82-9)	NA	NA	23.4± 1.2 mm
<i>Klebsiella pneumonia</i> (RCMB 01002 23-5)	21.4 ± 1.2 mm	21.6 ± 0.72 mm	27.2± 2.1 mm
<i>Pseudomonas aeruginosa</i> (RCMB 01002 43-5)	22.3 ± 2.1 mm	22.9± 0.63 mm	20.6± 1.5 mm
<i>Serratia plymuthica</i> (RCMB 01002 75-3)	20.2 ± 0.72 mm	20.8 ± 2.1 mm	22.3± 0.58 mm
<i>Serratia marcescens</i> (RCMB 01002 75b-8)	22.3 ± 1.5 mm	24.1 ± 0.58 mm	20.4± 0.58 mm
<i>Salmonella typhi</i> (RCMB 01002 15-4)	NA	NA	21.1± 0.72 mm
<i>Enterobacter cloacae</i> (RCMB 01002 64-5)	21.3 ± 0.58 mm	21.8± 1.5 mm	22.4± 2.1 mm
<i>Shigella dysenteriae</i> (RCMB 01002 41-8)	NA	NA	21.3± 1.5 mm

RCMB: Regional Center of Mycology and Biotechnology Antimicrobial unit test organism.

NA: No activity

species against both *A. spicifera* with clear zones [23.8±0.63 and 20.6±0.58 mm] and *D. simplex* with clear zones [22.4±2.1 and 21.2±0.63 mm] compared to the positive control Amphotericin B with zone of inhibition [20.3±1.5 and 20.3±0.58

mm]. *Candida albicans*, *Cryptococcus neoformans*, *Penicillium expansum* and *Trichophyton mentagrophytes* were resistance to the selected algae compared to the positive control Amphotericin B, on the other hand this

result was differ from previous reported where the inhibition zone (6 mm) was recorded in methanol extract against *C. albicans* [28].

The presence of flavonoids [29,30,31,32], terpenoides [33], saponins [34] and the strong presence of tanins [35] in the extract showed significant antimicrobial activity.

3.3 Anti-inflammatory Activity

In this study table and Fig. 4. represent the levels of SOD, MDA, Catalase, GSH, IL-6 and TNF-Alpha in the liver tissue homogenates of the normal and treated groups. Each parameter showed significant different degree in the disease state compared to the normal 'control'

Table 3. Antibacterial activity against fungi from red algae extracts (zone of inhibition in mm)

Sample tested microorganisms	<i>Acanthophora spicifera</i>	<i>Digenea simplex</i>	St.
Fungi			Amphotericin B
<i>Aspergillus fumigatus</i> (RCMB 02568)	21.3±1.2 mm	22.4±0.58 mm	25.7±1.5 mm
<i>Syncephalastrum racemosum</i> (RCMB 05922)	19.3±1.2 mm	20.3±1.2 mm	24.3±1.2 mm
<i>Geotricum candidum</i> (RCMB 05097)	23.8±0.63 mm	22.4±2.1 mm	20.3±1.5 mm
<i>Candida albicans</i> (RCMB 05036)	NA	NA	21.3±1.5 mm
<i>Aspergillus niger</i> (RCMB 02724)	20.6±0.58 mm	21.2±0.63 mm	20.3±0.58 mm
<i>Cryptococcus neoformans</i> (RCMB 05642)	NA	NA	21±1.0 mm
<i>Candida trobicalis</i> (RCMB 05239)	18.6±0.63 mm	20.3±1.2 mm	23.7±2.0 mm
<i>Penicillium expansum</i> (RCMB 01924)	NA	NA	21.7±2.0 mm
<i>Microsporium canis</i> (RCMB 0834)	20.3±1.2 mm	21.9±0.58 mm	23.3±1.5 mm
<i>Trichophyton mentagrophytes</i> (RCMB 0925)	NA	NA	21.3±1.5 mm

Data in table 1,2,3 are represented as mean zone of inhibition ± standard deviation.
RCMB: Regional Center of Mycology and Biotechnology Antimicrobial unit test organism.
NA: No activity




Fig. 4. Anti-inflammatory activity of *Acanthophora spicifera* and *Digenea simplex* on inflamed liver Wister Albino rats

Table 4. Anti-inflammatory activity of the studied algal species

Parameter group	SOD (U/ml)	MAD (nmol/ml)	Catalase (U/ml)	GSH (μ mol/ml)	IL-6 (Pg/ml)	TNF-Alpha (Pg/ml)
Group A	72.03 \pm 6.49	20.31 \pm 0.90	75.68 \pm 2.6	58.85 \pm 3.6	15.51 \pm 1.04	23.89 \pm 1.07
Group B	5.93 \pm 0.31	133.56 \pm 5.10	6.96 \pm 0.31	5.03 \pm 0.29	144.1 \pm 5.3	257.56 \pm 11.85
Group C (<i>Acanthophora spicifera</i>)	69.18 \pm 0.51	16.61 \pm 0.30	70.46 \pm 0.40	56.01 \pm 0.58	18.35 \pm 1.03	29.08 \pm 1.04
Group D (<i>Digenea simplex</i>)	70.18 \pm 0.73	18.25 \pm 0.29	72.3 \pm 0.36	56.63 \pm 0.33	19.63 \pm 1.02	26.58 \pm 1.95
F-Prob	P<0.0001	P<0.0001	P<0.0001	P<0.0001	P<0.0001	P<0.0001
LSD at5%	9.68	7.67	3.98	5.43	8.26	15.16
LSD at1%	13.20	10.46	5.43	7.41	11.27	22.32

Data in Table 4 expressed as Mean \pm S.E.M = Mean values \pm Standard error of means of six experiments. Number of animals in each group is six, Statistical evaluation of results was performed using the one-way ANOVA and least significant difference was determined at the level of LSD at 5% and LSD at 1%.

Test drugs: significant from normal control, P < 0.0001, SOD: Superoxide Dismutase, MAD: Malonaldehyde, GSH: Glutathione, IL-6: Interleukin-6, TNF- α : Tumor necrosis factor, Group A: Normal control, Group B: Diseased or inflamed Liver cancer (positive control receiving saline), Group C and D: Inflamed liver rats were treated by injection with methanol extract of the first drug (*Acanthophora spicifera*) and the second drug (*Digenea simplex*)

state. As it observed significant decreased in the levels of SOD from (72.03 \pm 6.49 to 5.93 \pm 0.31 U/ml), Catalase from (75.68 \pm 2.6 to 6.96 \pm 0.31 U/ml) and GSH from (58.85 \pm 3.6 to 5.03 \pm 0.29 μ mol/ml). Also observed that significant increase in the inflamed liver state of MDA from (20.31 \pm 0.90 to 133.56 \pm 5.10 nmol/ml), IL-6 from (15.51 \pm 1.04 to 144.1 \pm 5.3 Pg/ml) and TNF-Alpha from (23.89 \pm 1.07 to 257.56 \pm 11.85 Pg/ml). After treatment of inflamed liver rats with crude extracts using a dose of 500 mg/kg of each drug, observed that the difference in the parameters levels relatively equal to the normal control results.

4. CONCLUSION

The bioactive components from seaweeds crude extract have antimicrobial activity on some human pathogenic bacterial species and unicellular and filamentous fungi and anti-inflammatory effect which might protect the human health against some oxidative stress which attack DNA, proteins and membrane lipids and induced cellular damage.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Haroun BM, Sharaf AM, Ibraheem B. Evaluation of natural productions in some common Egyptian marine algae. J. Union. Arab Biol. B Botany. 1995;2:137-153.
2. O' Sullivan L, Murphy B, McLoughlin P, Duggan P, Lawlor PG, Hughes H, Gardiner GE. Prebiotics from marine macroalgae for human and animal health applications. Marine Drugs. 2010;8(7): 2038–2064.
3. Ibraheem BMI, Neveen AR, Mohamed SAH, Khaled EY. Antimicrobial and antiviral activities against newcastle disease virus (NDV) from marine algae isolated from qusier and marsaalam seashore (red sea), Egypt. African Journal of Biotechnology. 2012;11:8332-8340.
4. Elsayed KN, Radwan MM, Hassan SH, Abdelhameed MS, Ibraheem IB, Ross SA. Phytochemical and biological studies on some Egyptian seaweeds. Nat. Prod. Commun. 2012;7:1209–1210.
5. Al-Saif SS, Abdel-Raouf N, El-Wazanani HA, Aref IA. Antibacterial substances from marine algae isolated from Jeddah coast of Red sea, Saudi Arabia. Saudi Journal of Biological Sciences. 2014;21:57-64.
6. Abdel-Raouf N, Al-Enazi NM, Al-Homaidan AA, Ibraheem IBM, Al-Othman MR, Hatamleh. Antibacterial β -amylin isolated from *Lauren ciamicrocladia*. Arabian Journal of Chemistry. 2015;8(1):32–37.
7. Arif JM, Al-Hazzani AA, Kunhi M, Al-Khodairy F. Novel marine compounds: Anticancer or genotoxic? J Biomed Biotechnol. 2004;(2):93–98.

8. Hediat MH, Salama, NajatMarraiki. Antimicrobial activity and phytochemical analyses of *Polygonum aviculare* L. (*Polygonaceae*), naturally growing in Egypt. Saudi Journal of Biological Sciences. 2010;17:57-63.
9. Schramm A, Matusik P, Osmenda G, Guzik TJ. Targeting NADPH oxidases in vascular pharmacology. Vasc Pharmacol. 2012;56(5-6):216-231.
10. Rosanna DP, Salvatore C. Reactive oxygen species, inflammation, and lung diseases. Curr Pharm Des. 2012;18(26):3889-3900.
11. Kim YJ, Kim EH, Hahm KB. Oxidative stress in inflammation-based gastrointestinal tract diseases: Challenges and opportunities. J Gastroenterol Hepatol. 2012;27(6):1004-1010.
12. Hulsmans M, Van Dooren E, Holvoet P. Mitochondrial reactive oxygen species and risk of atherosclerosis. Curr Atheroscler Rep. 2012;14(3):264-276.
13. Abad MJ, Bedoya LM, Bermejo P. Natural marine anti-inflammatory products. Mini Rev Med Chem. 2008;8(8):740-754.
14. Wang W, Wang SX, Guan HS. The antiviral activities and mechanisms of marine polysaccharides: An overview. Mar Drugs. 2012;10(12):2795-2816.
15. D'Orazio N, Gammona MA, Gemello E, De Girolamo M, Cusenza S, Riccioni G. Marine bioactives: Pharmacological properties and potential applications against inflammatory diseases. Mar Drugs. 2012;10(4):812-833.
16. Ribeiro, et al. Marine algae as bioproducts for the prospection of new drugs. EC Pharmacology and Toxicology. 2017;3(3):64-65.
17. Aleem AA. Contributions to the study of the marine algae of the Red Sea. III- Marine algae from Obhor, in the vicinity of Jeddah, Saudi Arabia. Bull. Fac. Sci. KAU J. 1978;2:99-118.
18. Bold HC. "Introduction to the algae" structure and reproduction. Prentice. Hall. Inc. New-Jersey. 1978;07632.
19. Aleem AA. The marine algae of Alexandria. Egypt. Egyptian Books House; 1993.
20. Coppejans E, Leliaert F, Dargent O, Gunasekara K, Clerck O. Srilanka seaweeds. Methodologies and field guide to the dominant species. University of Ruhuna, Dept. of Botany. Matora, Srilanka. 2009;1-265.
21. Soad M, Mohy El-Din, Amani MD, El-Ahwany. Science direct bioactivity and phytochemical constituents of marine red seaweeds (*Jania rubens*, *Corallina mediterranea* and *Pterocladia capillacea*). Journal of Taibah University for Science. 2016;10:471-484.
22. Rahman A, Choudhary M, Thomsen W. Bioassay techniques for drug development. Hardwood Academic Publishers, The Netherlands. 2011;16.
23. International A. Official methods of analysis of AOAC International, AOAC International; 2005.
24. Keegan D. Foundations of distance education. Psychology Press; 1996.
25. Tapiero H, Tew KD, Naguyen G, Math_e G. Polyphenols: Do they play a role in the prevention of human pathologies? Review. Biomed. Pharmacother. 2002;56:200-207.
26. Wang J, Zhang Q, Zhang Z. Antioxidant activity of sulfated polysaccharide fractions extracted from *Laminaria japonica*. Int J Biol Macromols. 2008;42:127-132.
27. Frllich I, Riederer P. Free radical mechanisms in dementia of Alzheimer type and the potential for antioxidative treatment. Drug Research. 1995;45:443-c449.
28. Butterfield DA, Castenga A, Pocernich CB, Drake J, Scapagnini G, Calabrese CV. Nutritional approaches to combat oxidative stress in Alzheimer's disease. Journal of Nutritional Biochemistry. 2002;13:444-461.
29. Lin Y, Shi R, Wang X, Shen HM. Luteolin, a flavonoid with potential for cancer prevention and therapy. Curr. Can. Drug Targ. 2008;8:634-46.
30. Lopez-Lazaro M. Distribution and biological activities of the flavonoid luteolin. Mini Rev. Med. Chem. 2009;9:31-59.
31. Yoshida T, Konishi M, Horinaka M, Yasuda T, Goda AE, Taniguchi H, Yano K, Wakada M, Sakai T. Kaempferol sensitizes colon cancer cells to TRAIL-induced apoptosis. Biochem. Biophys. Res. Comm. 2008;375:129-133.
32. Amaral S, Mira L, Nogueira JM, da Silva AP, Florencio MH. Plant extracts with anti-inflammatory properties--a new approach

- for characterization of their bioactive compounds and establishment of structure-antioxidant activity relationships. *Bioorg. Med. Chem.* 2009;17(5):1876-1883.
33. Singh B, Singh S. Antimicrobial activity of terpenoids from *Trichodesma amplexicaule* Roth. *Phyto. Res.* 2003; 17(7):814-816.
34. Mandal P, Babu SSP, Mandal NC. Antimicrobial activity of saponins from *Acacia auriculiformis*. *Fitoterapia.* 2005;76(5):462-465.
35. Kaur GJ, Arora DS. Antibacterial and phytochemical screening of *Anethum graveolens*, *Foeniculum vulgare* and *Trachyspermum ammi*. *BMC Compl. Altern. Med.* 2009;9:30.

© 2017 Ibraheem et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://sciencedomain.org/review-history/19064>