



FUZZY OPTIMIZATION MODELING IN THE ANALYSIS OF HUMAN HEALTH CARE BASED ON LINEAR PROGRAMMING PROBLEM

A. Venkatesh*, G. Sivakumar* & S. Ramesh Babu**

* Assistant Professor, Department of Mathematics, A.V.V.M Sri Pushpam College, Poondi, Thanjavur, Tamilnadu

** Research Scholar, Department of Mathematics, A.V.V.M Sri Pushpam College, Poondi, Thanjavur, Tamilnadu

Cite This Article: A. Venkatesh, G. Sivakumar & S. Ramesh Babu, "Fuzzy Optimization Modeling in the Analysis of Human Health Care Based on Linear Programming Problem", International Journal of Applied and Advanced Scientific Research, Volume 2, Issue 2, Page Number 82-90, 2017.

Abstract:

Communicable diseases are the diseases which can be spread from one person to the other. It can also spread from infected animals. The transfer of the infection can occur through air, water, surfaces which are contaminated or through the direct contact. Moreover, *Communicable Diseases* are common among people nowadays days. In this paper, Fuzzy Optimization Model is developed based on Linear Programming Problem to control the communicable diseases with minimum curing time and dosage by distributing the various treatments to the disease population in order to reduce the human productivity loss. Finally, to demonstrate the feasibility of the proposed optimization model, an analytic technique is given for some four combinable diseases with four types of treatments.

Key Words: Fuzzy Optimization, Fuzzy Linear Programming Problem & Communicable Diseases

1. Introduction:

Optimization is the act of achieving the best possible result under given circumstances. In design, construction, maintenance etc, professionals have to take decisions. The goal of all such decisions is either to minimize effort or to maximize benefit. The effort or the benefit can be usually expressed as a function of certain design variables. Hence, optimization is the process of finding the conditions given the maximum or the minimum value of the function. In the last three decades many optimization techniques have been invented and successfully applied to optimizations problems in Computer Sciences, Information Technology, Engineering, Chemistry, Biology, Biochemistry, Medicine, Economics etc. Different modeling techniques are developed to meet the requirements of different types of optimization problems. Major categories of modeling approaches are classical optimization techniques, linear programming, non-linear programming, geometric programming, dynamic programming, integer programming etc. Transportation problem refers to a special class of linear programming problem. Since ancient day, the transportation problem is placing an important role in optimization problems to keep the balance in economic world. Transportation problem was developed in earlier days with single objective and with assumption that supply, demand and cost parameters are exactly known, where as in real life situations, the transportation problem with multi-objective and the parameters are not defined precisely. The necessity to optimize more than one objective or goal while satisfying the physical limitations led to the development of multi-objective programming methods. Fuzzy multi-objective programming satisfies all the requirements of real life situation to optimize the effort and benefit.

In recent years there has been a dramatic increase in the application to optimize techniques to the study the medical problems and the delivery of health care. To indicate the wide spread scope of the subject, some special typical applications in medical discipline are as follows: In 2008, Caetano et al [9] discussed optimal medication in HIV seropositive patient treatment using fuzzy cost function, Crina Grosan et al [3] developed Multi-Criteria programming model for medical diagnosis and treatment. In 2009, Denton et al [4] developed the optimization model to optimize the start time of strain therapy for patients with diabetes, Lee et al [7] have discussed about modeling and optimizing the public health infrastructure for emergency response. In 2010, Zhao Jingwei [16] discussed Fuzzy Multi-Objective Routing Inventory Problem in recycling infectious medical waste. In 2012, Mason et al [10] developed the revised optimization model to optimizing strain treatment decisions for diabetes patients in the presence of uncertain future adherence. In 2013, Persi Pamela et al [11] proposed a fuzzy optimization technique for the prediction of coronary Heart Disease using Decision Tree. Recently, in 2014, Lee et al [8] developed optimization modeling for emergency department workflow. This paper proposes Fuzzy Multi Objective Linear Programming Model to minimize the overall curing time of a disease population and medicinal who have to be cured by the various treatments. The main aim of this paper is to develop the multi objective optimization model based on Fuzzy Multi Objective Linear Programming technique to minimize the human productivity loss by distributing the various treatments to the different disease population so as to minimize the overall curing time and to minimize the overall dosage.

Preliminaries:

In this section, some basic definitions, generalized triangular fuzzy number, generalized trapezoidal fuzzy numbers and defuzzification, are presented.

Definition 2.1: Let R be the set of all real numbers. We assume a fuzzy number \tilde{A} that can be expressed for all $x \in R$ in the form

$$\mu_{\tilde{A}}(x) = \begin{cases} \mu_{\tilde{A}_L}(x) & a \leq x \leq b \\ w & b \leq x \leq c \\ \mu_{\tilde{A}_R}(x) & c \leq x \leq d \\ 0 & \text{otherwise} \end{cases}$$

Where $0 \leq w \leq 1$ is a constant, a, b, c, d are real numbers, such that $a < b \leq c < d$, $\mu_{\tilde{A}_L}(x): [a, b] \rightarrow [0, w]$, $\mu_{\tilde{A}_R}(x): [c, d] \rightarrow [0, w]$ are two strictly monotonic and continuous functions from R to the close interval $[0, w]$.

Since $\mu_{\tilde{A}_L}(x)$ is continuous and strictly increasing, the inverse function of $\mu_{\tilde{A}_L}(x)$ exists. Similarly $\mu_{\tilde{A}_R}(x)$ is continuous and strictly decreasing, the inverse function of $\mu_{\tilde{A}_R}(x)$ also exist. The inverse functions of $\mu_{\tilde{A}_L}(x)$ and $\mu_{\tilde{A}_R}(x)$ can be denoted by $\mu_{\tilde{A}_L}^{-1}(x)$ and $\mu_{\tilde{A}_R}^{-1}(x)$, respectively. $\mu_{\tilde{A}_L}^{-1}(x)$ and $\mu_{\tilde{A}_R}^{-1}(x)$ are

continuous on $[0, w]$ that means both $\int_0^w \mu_{\tilde{A}_L}^{-1}(x) dx$ and $\int_0^w \mu_{\tilde{A}_R}^{-1}(x) dx$ exist.

Definition 2.2: A fuzzy number $\tilde{A} = (a, b, c, d; w)$ is said to be generalized trapezoidal fuzzy number if its membership function is given by

$$\mu_{\tilde{A}}(x) = \begin{cases} w \left(\frac{x-a}{b-a} \right) & a \leq x \leq b \\ w & b \leq x \leq c \\ w \left(\frac{x-d}{c-d} \right) & c \leq x \leq d \\ 0 & \text{otherwise} \end{cases}$$

Definition 2.3: A fuzzy number $\tilde{A} = (a, b, c; w)$ is said to be generalized triangular fuzzy number if its membership function is given by

$$\mu_{\tilde{A}}(x) = \begin{cases} 0 & x \leq a \\ w \left(\frac{x-a}{b-a} \right) & a \leq x \leq b \\ w \left(\frac{c-x}{c-b} \right) & b \leq x \leq c \\ 0 & x > c \end{cases}$$

2. Defuzzification:




Figure 1: Centroid of Centroids

The process of converting the fuzzy output to a crisp value is said to be defuzzification. A number of defuzzification techniques are known, including centre-of-area, centre of gravity, and mean of maximums. A common and useful defuzzification technique is center of gravity. This technique was developed by Sugeno in 1985. This is the most commonly used technique and is very accurate. In 2011 and 2012, Phani Bushan Rao and others [12, 14, 15] have proposed a centroid formula for defuzzification using circumcenter of centroids, orthocenter of centroids, and centroid of centroids. In 2014, Hari Ganesh & Jayakumar [6] have proposed a centroid by using radius of gyration of centroid for ranking of fuzzy numbers. Herewith, a new centroid is proposed for defuzzification based on centroid of centroid which is presented as follows:

We define the centroid $G(\bar{x}_0, \bar{y}_0)$ of the triangle with vertices G_1, G_2 and G_3 of the generalized trapezoidal fuzzy number $\tilde{A} = (a, b, c, d; w)$ as

$$G(\bar{x}_0, \bar{y}_0) = \left(\frac{4a + 5b + 5c + 4d}{18}, \frac{5w}{9} \right) \quad (1)$$

Its Ranking Function is defined as

$$R(\tilde{A}) = \frac{4a + 5b + 5c + 4d}{18} \quad (2)$$

As a special case, for triangular fuzzy number $\tilde{A} = (a, b, d; w)$, i.e., cb the centroid of Centroids is given by

$$G(\bar{x}_0, \bar{y}_0) = \left(\frac{2a + 5b + 2d}{9}, \frac{5w}{9} \right) \quad (3)$$

Its Ranking Function is defined as

$$R(\tilde{A}) = \frac{2a + 5b + 2d}{9} \quad (4)$$

3. Mathematical Formulation of Fuzzy Multi-objective Transportation Model:

In this section, mathematical formulation of fuzzy multi-objective transportation model with fuzzy cost and fuzzy time is presented. Generally the fuzzy transportation problem is to transport various amounts of a single homogeneous commodity that are initially stored at various sources, to different destinations in such a way that the total fuzzy transportation cost is a minimum. Let there be m sources, i^{th} source possessing fuzzy supply units of a certain product, n destinations (n may or may not be equal to m) with destination j requiring fuzzy demand units. Cost of shipping of an item from each of m sources to each of the n destinations are known either directly or indirectly in terms of mileage, shipping hours, etc. If the objective of a transportation problem is to minimize fuzzy cost, and fuzzy time, then this type of fuzzy problem is treated as a fuzzy multi-objective transportation problem.

Mathematically, the fuzzy multi-objective transportation problem can be stated as:

$$\text{Minimize } \tilde{z}_k = \sum_{i=1}^m \sum_{j=1}^n (\tilde{p}_{ij}^k) x_{ij}$$

$$\text{Subject to } \sum_{j=1}^n x_{ij} \cong \tilde{a}_i \quad i = 1, 2, \dots, m$$

$$\sum_{i=1}^m x_{ij} \cong \tilde{b}_j \quad j = 1, 2, \dots, n$$

Where $\tilde{z}_k = \{\tilde{z}_1, \tilde{z}_2, \dots, \tilde{z}_k\}$

If the objective function \tilde{z}_1 denotes the fuzzy cost function,

$$\tilde{z}_1 = \sum_{i=1}^m \sum_{j=1}^n \tilde{c}_{ij} x_{ij}$$

If the objective function \tilde{z}_2 denotes the fuzzy cost function,

$$\tilde{z}_1 = \sum_{i=1}^m \sum_{j=1}^n \tilde{t}_{ij} x_{ij}$$

Then it is a bi-objective fuzzy transportation problem, which is represented by using weights of objectives to consider the priorities of the objective as follows:

$$\tilde{Z}_1 = w_1 \sum_{i=1}^m \sum_{j=1}^n \tilde{c}_{ij} x_{ij} + w_2 \sum_{i=1}^m \sum_{j=1}^n \tilde{t}_{ij} x_{ij}$$

Subject to

$$\sum_{j=1}^n x_{ij} \cong \tilde{a}_i \quad i = 1, 2, \dots, m$$

$$\sum_{i=1}^m x_{ij} \cong \tilde{b}_j \quad j = 1, 2, \dots, n$$

$$x_{ij} \geq 0 \quad i = 1, 2, \dots, m; j = 1, 2, \dots, n$$

And $w_1 + w_2 = 1$

Where

$\tilde{c}_{ij} = (\alpha_{ij}, \beta_{ij}, \gamma_{ij})$: Fuzzy cost from i^{th} source to j^{th} destination

$\tilde{t}_{ij} = (\alpha_{ij}, \beta_{ij}, \gamma_{ij})$: Fuzzy time from i^{th} source to j^{th} destination

$\tilde{a}_i = (\alpha_{ij}, \beta_{ij}, \gamma_{ij})$: Fuzzy supply from i^{th} source to j^{th} destination

$\tilde{b}_j = (\alpha_{ij}, \beta_{ij}, \gamma_{ij})$: Fuzzy demand from i^{th} source to j^{th} destination

All denotes $\tilde{c}_{ij}, \tilde{t}_{ij}, \tilde{a}_i, \tilde{b}_j$ a non-negative triangular fuzzy numbers.

$\sum_{i=1}^m \sum_{j=1}^n \tilde{c}_{ij} x_{ij}$: Total fuzzy cost for shipping from i^{th} source to j^{th} destination.

$\sum_{i=1}^m \sum_{j=1}^n \tilde{t}_{ij} x_{ij}$: Total fuzzy time for shipping from i^{th} source to j^{th} destination.

The same fuzzy transportation problem with two objectives may be represented in the following form of $m \times n$ fuzzy matrix (Table 1) where each cell having a fuzzy cost, and fuzzy time.

Table 1: Fuzzy Multi Objective Transportation Model with Fuzzy Cost and Fuzzy Time

Source / Destination	1	2	j	n	Supply
1	$\tilde{c}_{11}; \tilde{t}_{11}$	$\tilde{c}_{12}; \tilde{t}_{12}$	$\tilde{c}_{1j}; \tilde{t}_{1j}$	$\tilde{c}_{1n}; \tilde{t}_{1n}$	\tilde{a}_1
2	$\tilde{c}_{21}; \tilde{t}_{21}$	$\tilde{c}_{22}; \tilde{t}_{22}$	$\tilde{c}_{2j}; \tilde{t}_{2j}$	$\tilde{c}_{2n}; \tilde{t}_{2n}$	\tilde{a}_2
⋮			⋮		⋮		⋮
i	$\tilde{c}_{i1}; \tilde{t}_{i1}$	$\tilde{c}_{i2}; \tilde{t}_{i2}$	$\tilde{c}_{ij}; \tilde{t}_{ij}$	$\tilde{c}_{in}; \tilde{t}_{in}$	\tilde{a}_i
⋮			⋮		⋮		⋮
m	$\tilde{c}_{m1}; \tilde{t}_{m1}$	$\tilde{c}_{m2}; \tilde{t}_{m2}$	$\tilde{c}_{mj}; \tilde{t}_{mj}$	$\tilde{c}_{mn}; \tilde{t}_{mn}$	\tilde{a}_m
Demand	\tilde{b}_1	\tilde{b}_2		\tilde{b}_j		\tilde{b}_n	

4. Proposed Fuzzy Model for Planning Treatment of Communicable Diseases:

In this section, a Fuzzy Multi Objective Linear Programming Model is proposed based on multi objective fuzzy transportation model for computing minimum curing time and treatment dosage of a disease population affected by various communicable diseases in order to minimize the human productivity loss. This model is concerned with finding the overall minimum curing time and treatment dosage of a disease population affected by various communicable diseases which are to be cured by various treatments in a region.

The data of the model include

- ✓ The size of patients affected by each disease to be taken the treatment and the total availability of various treatments in a particular region.
- ✓ The unit curing time and treatment dosage (i.e. curing time and treatment dosage per patient) of the disease.

The objective is to determine how the various treatments may be distributed to the different disease population so as to minimize the overall curing time and to minimize the treatment dosage. Therefore, the decision variables are:

x_{ij} = the affordability of the j^{th} treatment to the i^{th} disease, where $i = 1, 2, \dots, m$ and $j = 1, 2, \dots, n$. That is a set of $m \times n$ variables.

In order to minimize the curing time and dosage, the following problem must be solved

The Objective Function:

Consider the size of patients to be taken the treatment i who have been affected by disease j . For any i and any j , the unit curing time is t_{ij} , unit dosage time d_{ij} , affordability of the treatment to the disease x_{ij} . Since we assume that the time and dosage functions are linear, the total curing time and total dosage is given by $t_{ij}x_{ij}$ and $d_{ij}x_{ij}$ respectively. Summing over all i and all j now yields the overall curing time and dosage for all disease – treatment combinations. That is, our objective functions are

$$\text{Minimize } \tilde{Z} = \sum_{i=1}^m \sum_{j=1}^n \tilde{t}_{ij} x_{ij}$$

$$\text{Minimize } \tilde{Z} = \sum_{i=1}^m \sum_{j=1}^n \tilde{d}_{ij} x_{ij}$$

Then it is a two objective transportation using the weights of the objectives which consider the priorities of the objective.

$$\tilde{Z} = w_1 \sum_{i=1}^m \sum_{j=1}^n \tilde{t}_{ij} x_{ij} + w_2 \sum_{i=1}^m \sum_{j=1}^n \tilde{d}_{ij} x_{ij}$$

The Constraints:

Consider treatment i , total affordability of this treatment for the various given diseases in the region is the sum $x_{i1} + x_{i2} + \dots + x_{in}$. Since the availability of this treatment for various diseases in the region is a_i , the affordability of this treatment for the various given diseases cannot exceed a_i .

$$\text{(i.e.) } \sum_{j=1}^n x_{ij} \leq a_i \text{ for } i = 1, 2, \dots, m$$

Consider disease j , the total affordability of various given treatments for this disease in the region is the sum $x_{1j} + x_{2j} + \dots + x_{mj}$. Since the total size of patients affected by this disease to be taken the treatment is b_j , the total affordability of various treatments should not be less than b_j .

$$\text{(i.e.) } \sum_{i=1}^m x_{ij} \geq b_j \text{ for } j = 1, 2, \dots, n$$

Where $x_{ij} \geq 0$ for all i and j

The above implies that the total availability of various given treatments for various given diseases $\sum_{i=1}^m a_i$ is greater than or equal to the total number of patients affected by the various given diseases $\sum_{j=1}^n b_j$.

When the total availability of various given treatments is equal to the total number of patients affected by the various given diseases (i.e. $\sum_{i=1}^m a_i = \sum_{j=1}^n b_j$) then the model is said to be balanced. In a balanced model, each of the constraints is an equation:

$$\sum_{j=1}^n x_{ij} = a_i \text{ for } i = 1, 2, \dots, m$$

$$\sum_{i=1}^m x_{ij} = b_j \text{ for } j = 1, 2, \dots, n$$

A model in which total availability of various given treatments and total number of patients affected by the various given diseases are unequal is called unbalanced. It is always possible to balance an unbalanced problem.

The fuzzy problem, in which the curing time t_{ij} , treatment dosage d_{ij} , total availability of treatment a_i and total number of patients to be taken the treatment b_j quantities are fuzzy quantities, can be formulated as follows:

$$\text{Minimize } \tilde{Z} = w_1 \sum_{i=1}^m \sum_{j=1}^n \tilde{t}_{ij} x_{ij} + w_2 \sum_{i=1}^m \sum_{j=1}^n \tilde{d}_{ij} x_{ij}$$

$$\text{Subject to } \sum_{j=1}^n x_{ij} \leq \tilde{a}_i \text{ for } i = 1, 2, \dots, m$$

$$\text{And } \sum_{i=1}^m x_{ij} \geq \tilde{b}_j \text{ for } j = 1, 2, \dots, n$$

Where $x_{ij} \geq 0$ for all i and j

This fuzzy problem is explicitly represented by the following table.

Table 2: Fuzzy Model for Optimization of Time and dosage of Treatment of Diseases

Treatments / Diseases	D ₁	D ₂	D _j	D _n	Supply (availability of treatment T _i)
T ₁	$\tilde{t}_{11}; \tilde{d}_{11}$	$\tilde{t}_{12}; \tilde{d}_{12}$	$\tilde{t}_{1j}; \tilde{d}_{1j}$	$\tilde{t}_{1n}; \tilde{d}_{1n}$	\tilde{a}_1
T ₂	$\tilde{t}_{21}; \tilde{d}_{21}$	$\tilde{t}_{22}; \tilde{d}_{22}$	$\tilde{t}_{2j}; \tilde{d}_{2j}$	$\tilde{t}_{2n}; \tilde{d}_{2n}$	\tilde{a}_2
⋮			⋮		⋮		⋮
T _i	$\tilde{t}_{i1}; \tilde{d}_{i1}$	$\tilde{t}_{i2}; \tilde{d}_{i2}$	$\tilde{t}_{ij}; \tilde{d}_{ij}$	$\tilde{t}_{in}; \tilde{d}_{in}$	\tilde{a}_i
⋮			⋮		⋮		⋮
T _m	$\tilde{t}_{m1}; \tilde{d}_{m1}$	$\tilde{t}_{m2}; \tilde{d}_{m2}$	$\tilde{t}_{mj}; \tilde{d}_{mj}$	$\tilde{t}_{mn}; \tilde{d}_{mn}$	\tilde{a}_m
Demand (no. of patients affected by the disease D _i to be taken the treatment)	\tilde{b}_1	\tilde{b}_2	\tilde{b}_j	\tilde{b}_n	

5. Example:

Communicable diseases are diseases that are as a result of the causative organism spreading from one person to another. They are among the major causes of illnesses in many countries. These diseases affect people of all ages but more so children due to their exposure to environmental conditions that support the spread. Communicable diseases are preventable base on interventions placed on various levels of transmission of the disease. Health Departments have an important role to play in the control of these diseases by applying effective and efficient management, prevention and control measures. In Thanjavur Region, the availability of various treatments like Allopathy (T₁), Ayurvedic (T₂), Homeopathy (T₃) and Unani (T₄) for all type of diseases are (50000, 52000, 55000), (31000, 34000, 37000), (10500, 12500, 14500), and (5500, 7500, 9500) respectively. Moreover, the size of patients affected by the communicable diseases in winter season like Dengue (D₁), Malaria (D₂) and Tuberculosis (D₃) are (21500, 22500, 25500), (14250, 17250, 19500), and (10250, 12450, 15500) respectively. Curing time and treatment dosage for all above said treatment - disease combination per patient are as follows:

Table 3: Curing Time & dosage per Patient

Treatment	Disease	Curing Time per Patient (in days)	Dosage per Patient (in gms) (per course)
Allopathy	Dengue	(31, 35, 37)	(5, 6, 7)
	Malaria	(20, 22, 25)	(8, 10, 12)
	Tuberculosis	(250, 270, 300)	(450, 500, 550)
Ayurvedic	Dengue	(19, 20, 24)	(100, 300, 500)

	Malaria	(29, 30, 34)	(500, 550, 600)
	Tuberculosis	(90, 120, 130)	(1000, 1500, 2000)
Homeopathy	Dengue	(70, 90, 110)	(0.5, 0.7, 1.0)
	Malaria	(55, 75, 95)	(1, 1.5, 2)
	Tuberculosis	(325, 345, 355)	(40, 45, 50)
Unani	Dengue	(100, 120, 130)	(200, 300, 400)
	Malaria	(60, 90, 120)	(400, 470, 500)
	Tuberculosis	(400, 420, 450)	(1000, 1250, 1500)

The data collected from the Department of Medical and Rural Health Services at Thanjavur District.

Let us consider a optimization problem with rows representing treatments Allopathy (T_1), Ayurvedic (T_2), Homeopathy (T_3) and Unani (T_4) and column representing communicable diseases Dengue (D_1), Malaria (D_2) and Tuberculosis (D_3) which are affected in the winter season at Thanjavur Region.

Table 4: Unbalanced Table with Fuzzy Curing Time and Fuzzy Dosage

Treatments / Diseases	Dengue (D_1)	Malaria (D_2)	Tuberculosis (D_3)	Supply (availability of treatment T_i)
Allopathy (T_1)	(31,35,37) (5, 6, 7)	(20,22,25) (8, 10, 12)	(250,270,300) (450, 500, 550)	(50000, 52000, 55000)
Ayurvedic (T_2)	(19,20,24) (100, 300, 500)	(29,30,34) (500, 550, 600)	(90,120,130) (1000, 1500, 2000)	(31000, 34000, 37000)
Homeopathy (T_3)	(70,90,110) (0.5, 0.7, 1.0)	(55,75,95) (1, 1.5, 2)	(325,345,355) (40, 45, 50)	(10500, 12500, 14500)
Unani (T_4)	(100,120,130) (200, 300, 400)	(60,90,120) (400, 470, 500)	(400,420,450) (1000, 1250, 1500)	(5500, 7500, 9500)
Demand (no. of patients affected by the disease D_i to be taken the treatment)	(21500, 22500, 25500)	(14250, 17250, 19500)	(10250, 12450, 15500)	

Using the ranking function in equation (4), the values of $R(\tilde{c}_{ij})$, $R(\tilde{t}_{ij})$, $R(\tilde{a}_i)$ and $R(\tilde{b}_j)$ for all i and j are calculated and given in the following table.

Table 5: Unbalanced Table with Curing Time and Dosage

Treatments / Diseases	Dengue (D_1)	Malaria (D_2)	Tuberculosis (D_3)	Supply (availability of treatment T_i)
Allopathy (T_1)	35 6	22 10	272 500	52222
Ayurvedic (T_2)	21 300	31 550	116 1500	34000
Homeopathy (T_3)	90 0.7	75 1.5	343 45	12500
Unani (T_4)	118 300	90 461	422 1250	7500
Demand (no. of patients affected by the disease D_i to be taken the treatment)	22944	17083	12639	

The above problem is unbalanced. For make it as a balanced one, the dummy column is introduced as follows:

Table 6: Balanced Table with Curing Time and Dosage

Treatments / Diseases	Dengue (D_1)	Malaria (D_2)	Tuberculosis (D_3)	(D_4)	Supply (availability of treatment T_i)
Allopathy (T_1)	35 6	22 10	272 500	0 0	52222
Ayurvedic (T_2)	21 300	31 550	116 1500	0 0	34000
Homeopathy (T_3)	90 0.7	75 1.5	343 45	0 0	12500
Unani (T_4)	118 300	90 461	422 1250	0 0	7500
Demand (no. of patients affected by the disease D_i to be taken the treatment)	22944	17083	12639	53556	106222

The above crisp multi-objective transportation problem is converted into the following crisp linear programming problem by using step 6.

$$\begin{aligned} \text{Minimize} \quad & (23.4)x_{11} + (17.2)x_{12} + (363.2)x_{13} + (0)x_{14} + \\ & (132.6)x_{21} + (238.6)x_{22} + (669.6)x_{23} + (0)x_{24} + \\ & (54.28)x_{31} + (45.6)x_{32} + (223.8)x_{33} + (0)x_{34} + \\ & (190)x_{41} + (238.4)x_{42} + (753.2)x_{43} + (0)x_{44}. \\ \text{Subject to:} \quad & x_{11} + x_{21} + x_{31} + x_{41} = 22944 & x_{12} + x_{22} + x_{32} + x_{42} = 17083 \\ & x_{13} + x_{23} + x_{33} + x_{43} = 12639 & x_{14} + x_{24} + x_{34} + x_{44} = 53556 \\ & x_{11} + x_{12} + x_{13} + x_{14} = 52222 & x_{21} + x_{22} + x_{23} + x_{24} = 34000 \\ & x_{31} + x_{32} + x_{33} + x_{34} = 12500 & x_{41} + x_{42} + x_{43} + x_{44} = 7500 \end{aligned}$$

Using TORA software, the crisp linear programming problem is solved to find the optimum solution which is as follows:

$$x_{14}=12056, x_{11}=22944, x_{16}=7500, x_{13}=139, x_{33}=12500, x_{24}=34000, x_{12}=17083.$$

The overall minimum curing time and dosage respectively are 5504174 days and 940494.

6. Conclusion:

In multi-objective optimization problems, the problem of minimizing the total cost and time of transportation is studied as multi-objective transportation problem since long time and is well known in which time – cost minimization problem has been studied by many authors. In 1977, Bhatia [1] and others have given a procedure for time minimization in transportation problem. In 1981, Prakash [13] introduced transportation problem with objectives to minimize total cost and duration of transportation. Recently, in 2010, Chakraborty Ananya [2] has proposed a method for the minimization of transportation cost as well as time of transportation when the demand, supply and transportation cost are fuzzy. In 2014, Hari Ganesh & Jayakumar [5] proposed a fuzzy linear programming model for minimizing agricultural production cost. Hence, in this paper, Fuzzy Multi Objective Linear Programming Model has been developed in order to distribute the various treatments to the different disease population so as to minimize the overall curing time and to minimize the dosage by employing the supply, demand, time and dosage parameters as triangular fuzzy numbers. As minimizing the overall curing time and dosage, the human productivity loss may be minimized.

7. References:

1. Bhatia, H.L., et al. (1977), “A procedure for time minimizing transportation problem”, Indian Journal of Pure and Applied Mathematics, 8, pp. 920-929.
2. Chakraborty Ananya and Chakraborty M., (2010), Cost-time Minimization in a Transportation Problem with Fuzzy Parameters: A Case Study, Journal of Transportation Systems Engineering and Information Technology, 10(6), pp. 53–63.
3. Crina Grosan et al., (2008), Multi Criteria Programming in Medical Diagnosis and Treatment, Applied Soft Computing, 8(4), pp. 1407 – 1417.
4. Denton, B.T., (2009), Optimizing the Start time of Statin Therapy for Patients with Diabetes. Med Decis Making, 29(3), pp.351 – 367.
5. Hari Ganesh, A. and Jayakumar, S., (2014), A Fuzzy Linear Programming Model for Optimizing Agricultural Production Cost, International Journal of Applied Mathematical Sciences, 7(1), pp. 59-70.
6. Hari Ganesh, A. and Jayakumar, S., (2014), Ranking of Fuzzy Numbers Using Radius of Gyration of Centroids, International Journal of Basic and Applied Sciences, 2014 Vol. 3, No. 1, pp 17 – 22.
7. Lee, E.K. et al., (2009) Modeling and optimizing the public health infrastructure for emergency response. Interfaces – The Eaniel H. Wagner Prize for Excellence in Operations Research Practice; 39(5), 476 – 490.
8. Lee, E.K. et al., (2014) Systems analytics: Modeling and optimizing emergency department workflow. In: Y Hui, EK Lee, eds. Healthcare Data Analytics. Wiley Series in Operations Research and Management Science. John Wiley & Sons, Inc.
9. Marco A.L. Caetano et al., (2008) Optimal Medication in HIV Seropositive Patient Treatment using Fuzzy Cost Function, Proceedings of American Control Conference, IEEE Xplore, Washington, USA.
10. Mason J.E., et al., (2012), Optimizing Statin Treatment Decisions for Diabetes Patients in the Presence of Uncertain Future Adherence. Med Decis Making, 32(1), pp.154 – 156.
11. Persi Pamela, I. et al., (2013), A Fuzzy Optimization Technique for the Prediction of Coronary Heart Disease Using Decision Tree, International Journal of Engineering and Technology, 5(3), pp. 2506 – 2514.
12. Phani Bushan Rao & Ravi Shankar, N. (2011), Ranking Fuzzy Numbers with a Distance Method using Circumcenter of centroids and an Index Modality, Advances in Fuzzy Systems, pp. 1 -7.
13. Prakash, S., (1981), “A transportation problem with objective to minimize cost and duration of transportation”, Opsearch, 18, pp. 235-238.

14. Ravi Shankar. N., et al., (2012), Fuzzy Risk Analysis based on A New Approach of Ranking Fuzzy Numbers using Orthocenter of Centroids, International Journal of Computer Applications, Vol. 42, No. 3, 24 – 36.
15. Suresh Babu. S., et al., (2012), Ranking Generalized Fuzzy Numbers using centroid of centroids, International Journal of Fuzzy Logic Systems (IJFLS), Vol2, No.3, 17 – 32.
16. Zhao Jingwei, et al., (2010), Fuzzy Multi-objective Location – Routing – Inventory Problem in Recycling Infectious Medical Waste, Proceedings of International Conference on E – Business and E – Government, pp. 4069 – 4073.