

Table S1. Comparison of mean THg concentrations (in mg/kg ww) in fish species from Mediterranean locations.

| Fish species | Location (Country) | THg | Reference |
|----------------|-----------------------------|-------|--|
| Sardine | Balearic Islands | 0.050 | This study |
| | València | 0.040 | Yusà <i>et al.</i> , 2008 |
| | Catalonia | 0.080 | Cano-Sancho <i>et al.</i> , 2015; Perelló <i>et al.</i> , 2014 |
| | Sicily (Italy) | 0.080 | Copat <i>et al.</i> , 2012 |
| | | 0.31 | |
| | Augusta Bay (Italy) | 0.082 | Bonsignore <i>et al.</i> , 2013 |
| | Italy | 0.092 | Brambilla <i>et al.</i> , 2013 |
| Anchovy | Balearic Islands | 0.057 | This study |
| | Sicily (Italy) | 0.030 | Copat <i>et al.</i> , 2012 |
| | | 0.13 | |
| | Augusta Bay (Italy) | 0.052 | Bonsignore <i>et al.</i> , 2013 |
| | Italy | 0.11 | Brambilla <i>et al.</i> , 2013 |
| | València | 0.030 | Yusà <i>et al.</i> , 2008 |
| | Catalonia | 0.087 | Perelló <i>et al.</i> , 2014 |
| Hake | Balearic Islands | 0.30 | This study |
| | Catalonia | 0.13 | Perelló <i>et al.</i> , 2014 |
| | València | 0.15 | Yusà <i>et al.</i> , 2008 |
| | Italy | 0.16 | Brambilla <i>et al.</i> , 2013 |
| | Roses (Catalonia) | 0.51 | Torres <i>et al.</i> , 2015 |
| | | 0.62 | |
| | Italy | 0.59 | Perugini <i>et al.</i> , 2014 |
| Sole | Balearic Islands | 0.45 | This study |
| | Catalonia | 0.027 | Martorell <i>et al.</i> , 2011 |
| | | 0.057 | Perelló <i>et al.</i> , 2014 |
| | Italy | 0.14 | Brambilla <i>et al.</i> , 2013 |
| | Sfax (Tunisia) | 1.8* | Zohra and Habib 2016 |
| Angler | Balearic Islands | 0.74 | This study |
| | Italy | 0.13 | Brambilla <i>et al.</i> , 2013 |
| Horse mackerel | Balearic Islands | 0.18 | This study |
| | Italy | 0.50 | Brambilla <i>et al.</i> , 2013 |
| | Augusta Bay (Italy) | 0.13 | Bonsignore <i>et al.</i> , 2013 |
| | Catalonia | 0.053 | Martorell <i>et al.</i> , 2011 |
| | | 0.099 | Perelló <i>et al.</i> , 2014 |
| | Valencia | 0.030 | Yusà <i>et al.</i> , 2008 |
| Red mullet | Balearic Islands | 0.18 | This study |
| | Portmán | 0.17 | This study |
| | Italy | 0.43 | Storelli and Barone, 2013 |
| | | 0.48 | Perugini <i>et al.</i> , 2014 |
| | | 0.23 | Brambilla <i>et al.</i> , 2013 |
| | Catalonia | 0.33 | Perelló <i>et al.</i> , 2014 |
| | Portmán Bay (Spain) | 0.12 | Martínez-Gómez <i>et al.</i> , 2012 |
| | Santa Pola | 0.078 | |
| | València | 0.093 | |
| | Delta de l'Ebre (Catalonia) | 0.074 | |

*THg concentration in dry weight instead of wet weight.

Table S2. Daily fish consumptions, THg weekly intake through fish consumption, estimated weekly intake (EWI) and Provisional Tolerable Weekly Intake percentages (PTWI %), referred to the infant and general Spanish population by AESA (2006) and Madrid Community ENUCAM (2015).

| | | Daily fish consumption g/day | Median THg in fish mg/kg ww | THg weekly intakes µg/week | EWI µg/kg bw | PTWI ^a (%) | PTWI ^b (%) |
|--------------|----------------------|------------------------------|-----------------------------|----------------------------|--------------|-----------------------|-----------------------|
| AESA, 2011 | children | 46.4 | | 94.2 | 2.7 | 68 | 210 |
| | adult | 71.1 | | 144.3 | 2.1 | 53 | 162 |
| ENUCAM, 2015 | adult* | 61.9 | | 125.7 | 1.8 | 46 | 141 |
| | men* | 58.3 | 0.29 | 118.4 | 1.5 | 37 | 113 |
| | women* | 64.9 | | 131.8 | 2.0 | 51 | 157 |
| | Women (18-44 years)* | 57.2 | | 116.1 | 1.8 | 45 | 138 |

Children: 7-12 years with a mean body weight of 34.48 kg; Adults: >17 years with a mean body weight of 68.48 kg (AESA, 2006). *Adults, men: >18 years with a mean body weight of 80.75 kg; *Adults, women: >18 years with a mean body weight of 64.72 kg (ENUCAM, 2015). ^aBased on THg PTWI of 4 µg/kg bw (EFSA, 2012). ^bBased on MeHg PTWI of 1.3 µg/kg bw (EFSA, 2012). *Based on ENUCAM (2015).

References.

- AESA, 2011. Agencia Española de Seguridad Alimentaria (AESA). Modelo de dieta española para la determinación de la exposición del consumidor a sustancias químicas. Madrid (Spain): Ministerio de Sanidad y Consumo. URL: http://www.aecosan.msssi.gob.es/AECOSAN/docs/documentos/para_consumidor/modelo_dieta_espanola.pdf
- Bonsignore, M., Salvagio Manta, D., Oliveri, E., Sprovieri, M., Basilone, G., Bonanno, A., Falco, F., Traina, A., Mazzola, S., 2013. Mercury in fishes from Augusta Bay (southern Italy): Risk assessment and health implication. *Food Chem Toxicol* 56, 184 – 194.
- Brambilla, G., Abete, M.C., Binato, G., Chiaravalle, E., Cossu, M., Dellatte, E., Miniero, R., Orletti, R., Piras, P., Roncarati, A., Ubaldi, A., Chessa, G. 2013. Mercury occurrence in Italia seafood from the Mediterranean Sea and possible intake scenarios of the Italian coastal population. *Reg Toxicol Pharmacol* 65, 269 – 277.
- Cano-Sancho, G., Perello, G., Maulvault, A.L., Marques, A., Nadal, M., Domingo, J.L., 2015. Oral bioaccessibility of arsenic, mercury and methylmercury in marine species commercialized in Catalonia (Spain) and health risks for the consumers. *Food Chem Toxicol* 86, 34-40.
- Copat, C., Bella, F., Castaing, M., Fallico, R., Sciacca, S., Ferrante, M., 2012. Heavy Metals Concentrations in Fish from Sicily (Mediterranean Sea) and Evaluation of Possible Health Risks to Consumers. *Bull Environ Contam Toxicol* 88, 78 – 83.
- EFSA, 2012. Scientific opinion on the risk for public health related to the presence of mercury and methylmercury in food. *EFSA J* 10, 2985 – 3226.
- ENUCAM, 2015. Encuesta de Nutrición de la Comunidad de Madrid. <http://www.madrid.org/bvirtual/BVCM017699.pdf>
- Martínez-Gómez, C., Fernández, B., Benedicto, J., Valdés, J., Campillo, J.A., León, V.M., Vethaak, A.D., 2012. Health status of red mullets from polluted areas of the Spanish Mediterranean coast, with special reference to Portmán (SE Spain). *Mar Environ Res* 77, 50 – 59.
- Martorell, I., Perelló, G., Martí-Cid, R., Llobet, J.M., Castell, V., Domingo, J.L., 2011. Human exposure to arsenic, cadmium, mercury, and lead from foods in Catalonia, Spain: temporal trend. *Biol Trace Elem Res* 142, 309–322.
- Perello, G., Llobet, J.M., Gómez-Catalán, J., Castell, V., Centrich, F., Nadal, M., Domingo, J.L., 2014. Human Health Risks Derived from Dietary Exposure to Toxic Metals in Catalonia, Spain: Temporal Trends. *Biol Trace Elem Res* 162, 26 – 37.
- Perugini, M., Visciano, P., Manera, M., Zaccaroni, A., Olivieri, V., Morena, M., 2014. Heavy metals (As, Cd, Hg, Pb, Cu, Zn, Se) concentrations in muscle and bone of four commercial fish caught in the central Adriatic Sea, Italy. *Environ Monit Assess* 186, 2205 – 2213.
- Storelli, M.M., Barone, G., 2013. Toxic Metals (Hg, Pb, and Cd) in Commercially Important Demersal Fish from Mediterranean Sea: Contamination Levels and Dietary Exposure Assessment. *J Food Sci* 78, 362 – 366.
- Torres, J., Eira, C., Miquel, J., Ferrer-Maza, D., Delgado, E., Casadevall, M., 2015. Effect of Intestinal Tapeworm *Clestobothrium crassiceps* on Concentrations of Toxic Elements and Selenium in European Hake *Merluccius merluccius* from the Gulf of Lion (Northwestern Mediterranean Sea). *J Agr Food Chem* 63, 9349 – 9356.
- Yusà, V., Suelves, T., Ruiz-Atienza, L., Cervera, M.L., Benedicto, V., Pastor, A., 2008. Monitoring programme on cadmium, lead and mercury in fish and seafood from Valencia, Spain: levels and estimated weekly intake. *Food Add Contam Part B Surveil* 1, 22 – 31.
- Zohra, B.S., Habib A., 2016. Assessment of heavy metal contamination levels and toxicity in sediments and fishes from the Mediterranean Sea (southern coast of Sfax, Tunisia). *Environ Sci Poll Res* 23, 13954-13963.