

Supplementary table 12. Behavioral tests data about: latency to fall (seconds) from the accelerating rotarod (ROT), total distance moved in open field arena (OF), sweet-taste solution preference (SPT) at three months post-LPS injection

	Animal	ROT Time (s)	OF Total distance (cm)	SPT sucrose intake
	1	142	581,77	220,68
	2	220	580,35	215,74
	3	172	437,28	202,57
	4	109	2626,81	193,57
SAL WT	5	101	5594,92	176,74
	6	145	3360,56	202
	7	144	1453,24	190,74
	8	113	2731,05	189,64
	9	63	1856,00	171,66
	10	181	2742,31	171,17
	1	111	972,785	214,67
	2	160	287,185	196,62
	3	204	861,943	226,3
LPS WT	4	121	1961,1	184,43
	5	202	4262,29	199,69
	6	131	3696,09	179,01
	7	39	1544,66	183,15
	8	214	2246,82	206,61
	9	46	2446,15	218,49
	1	141	2331,02	216,93
	2	171	2917,33	199,35
	3	154	2918,99	218,07
	4	139	617,23	176,77
SAL <i>Snca</i>+/+	5	158	259,65	153,55
	6	82	524,83	137,31
	7	80	704,48	99,2
	8	89	836,30	180,88
	9	111	3320,93	220,88
	10	118	5683,97	199,22
	1	123	1517,76	230,08
	2	214	2722,86	183,9
	3	108	3168,13	227,16
	4	136	665,487	186,25
LPS <i>Snca</i>+/+	5	139	3476,04	214,46
	6	134	5110,43	210,41
	7	151	3628,69	201,21
	8	80	4041,56	209,46
	9	144	3668,58	217,42