

On taxonomy and distribution of the genus *Carpelimus* Leach, 1819 (Coleoptera: Staphylinidae: Oxytelinae) of Nepal

Mikhail Yu. Gildenkov

Gildenkov M. Yu. 2014. On taxonomy and distribution of the genus *Carpelimus* Leach, 1819 (Coleoptera: Staphylinidae: Oxytelinae) of Nepal. *Baltic J. Coleopterol.*, 14(1): 21 - 31.

New distributional records from Nepal for thirteen species of the genus *Carpelimus* Leach, 1819 are presented. Six species are new for the fauna of Nepal: *Carpelimus (Bucephalinus) laticeps* (Cameron, 1930); *Carpelimus* (s.str.) *congruus* (Cameron, 1930); *Carpelimus* (s.str.) *indicus* (Kraatz, 1859); *Carpelimus (Trogophloeus) bengalensis* (Cameron, 1930); *Carpelimus (Trogophloeus) corticinus* (Gravenhorst, 1806); *Carpelimus (Trogophloeus) palitans* (Cameron, 1930). Two new combinations are proposed: *Carpelimus (Bucephalinus) kathmanduensis* Herman, 2001, comb. n. and *Carpelimus (Bucephalinus) nepalicus* (Coiffait, 1982), comb. n.

Key words: Coleoptera, Oxytelinae, *Carpelimus*, Nepal, new combinations, new records.

Mikhail Yu. Gildenkov. Smolensk State University, Department of Ecology and Chemistry, Przhhevsky str., 4, Smolensk 214000, Russia. E-mail: mgildenkov@mail.ru

INTRODUCTION

The genus *Carpelimus* Leach, 1819 is one of the largest genera within the subfamily Oxytelinae Fleming, 1821; it includes about 500 species in the fauna of the world and is widely distributed in all zoogeographical areas. The majority of species of the genus inhabit damp ground on the supralittoral, where they can be very abundant and play an important role in the functioning of ecological systems.

According to the catalogue of Smetana (2004), twelve species are known in Nepal (the names of species and subgenera are listed according to

this catalogue): *Carpelimus (Boopinus) insularis* (Kraatz, 1858); *C. (Boopinus) nigrita* (Wollaston, 1857); *C. (Boopinus) subpolitus* (Coiffait, 1982); *C. (Paraboopinus) travei* (Coiffait, 1982); *C. (s. str.) kathmanduensis* Herman, 2001; *C. (Taenosoma) apterus* (Coiffait, 1982); *C. (Taenosoma) bertrandi* (Coiffait, 1982); *C. (Taenosoma) brachypterus* (Coiffait, 1982); *C. (Taenosoma) nepalicus* (Coiffait, 1982); *C. (Troginus) despectus* (Baudi di Selve, 1870); *C. (Troginus) exiguus* (Erichson, 1839); *C. (Trogophloeus) vagans* (Cameron, 1930). Seven species from this list are described and known only from the Nepal. The level of endemism is compound 58%.

With the account of the synonymy of the generic groups in *Carpelimus* (Herman, 1970: 389, 390; Gildenkov, 2002: 1462, 1463; Gildenkov, 2002a: 1275; Gildenkov, 2013a: 243) it is necessary to recognize, that subgenera *Boopinus*, *Paraboopinus* and *Taenosoma* are not valid now. Some combinations of taxonomical names were changed during studies. Several species were synonymized (Gildenkov, 2014: 300, 304, 308): *C.* (s. str.) *peregrinus* (Cameron, 1919) = *C.* (s. str.) *subpolitus* (Coiffait, 1982), syn.; *C.* (*Troginus*) *atomus* (Saulcy, 1864) = *C.* (*Troginus*) *travei* (Coiffait, 1982), syn.; *C.* (*Trogophloeus*) *brachypterus* (Coiffait, 1982) = *C.* (*Trogophloeus*) *apterus* (Coiffait, 1982), syn. At the present time *C. nigrita* is differently understood (Schülke, 2004: 398; Gildenkov, 2010: 12; 2014: 299) as both the nominative subspecies *C.* (s. str.) *nigrita nigrita* (Wollaston, 1857), known only on the holotype from the archipelago Madeira (island Porto Santo), and also the widespread subspecies *Carpelimus* (s. str.) *nigrita anthracinus* (Mulsant and Rey, 1861). To assume the discovery of *C. nigrita nigrita* in Nepal (taking into account its incorrect understanding) will be strange as there are no data from Nepal for *C. nigrita anthracinus*.

Thus, ten species of the genus *Carpelimus* are known from the territory of Nepal, of which four species were known only from this country (40 %).

MATERIAL AND METHODS

This paper is based on specimens which are deposited in the following collections: BMNH = British Museum of Natural History (London, Great Britain); cMG = private collection of M. Gildenkov (Smolensk, Russia); cVA = private collection of V. Assing (Hannover, Germany); FMNH = Field Museum of Natural History (Chicago, the USA); HNHM = Hungarian Natural History Museum (Budapest, Hungary); MNHN = Muséum National d'Histoire Naturelle (Paris, France); NHMW = Naturhistorisches Museum Wien (Vienna, Austria); NKME =

Naturkundemuseum Erfurt (Erfurt, Germany); SDEI = Senckenberg Deutsches Entomologisches Institut (Müncheberg, Germany); SMNS = Staatliches Museum für Naturkunde in Stuttgart (Stuttgart, Germany).

In the present study standard methods were used for the taxonomic research of the insects; the preparations were made with the use of the binocular microscope MBS-10. The genital preparations were processed using 10% KOH and fixed in euparal later.

RESULTS

Carpelimus (Bucephalinus) kathmanduensis Herman, 2001 **comb. n.**

(*Trogophloeus (Carpelimus) nepalicus* Coiffait, 1984, nom. praeocc.; *C.* (s.str.) *kathmanduensis*)

Coiffait, 1984: 385

Material. Holotypus, male: "Nepal 8. 83 | Kathmandu gokarna" "*Carpelimus nepalicus* H.Coiffait det. 19" "TYPE (red label)" "Holotypus *Trogophloeus (Carpelimus) nepalicus* Coiffait, 1984 | rev. M.Gildenkov, 2013" (MNHN).

Remarks. This species is known only from the Nepal.

Carpelimus (Bucephalinus) laticeps (Cameron, 1930)

Material. 1 female: "NEPAL: Phewa-Tal; Pokhara. 85°59'E., 28°14'N. 2.500 Ft. 8-10.v.1954. K.H. Hyatt." "In cattle dung." "B.M. Nepal Expdt. B.M. 1954-540." (BMNH).

Remarks. The lectotype was designated from India (Gildenkov, 2013: 57). The species is known from the Oriental region (our data): India, Indonesia (Sumatra), Nepal, Singapore, Sri Lanka, Thailand, Vietnam. It is here reported from Nepal for the first time.

Carpelimus (Bucephalinus) nepalicus (Coiffait, 1982) **comb. n.**
(*C. (Taenosoma) nepalicus*)

Coiffait, 1982: 161

Material. Nepal: Holotypus, male (under the description, aedeagus is not present): “Nepal 21.X.76 | Pokhara 12” “Preparation microscopique” “HOLOTYPE” “*Carpelimus nepalicus* (Coiffait, 1982) | rev. Gildenkov, 2007” “MUSEUM PARIS” (MNHN); 1 female: “Nepal 21.X.79 [error in the label; must be: 21.X.76] | Pokhara 12” “PARATYPE” “*Carpelimus nepalicus* (Coiffait, 1982) | rev. Gildenkov, 2007” “MUSEUM PARIS” (MNHN).

Additional material. Vietnam: 6 females, 2 specimens: “VIETNAM: Cuc Phuong Ninh binh, 3-10.V.1966 Exp. Gy. TOPAL” “Nr. 309 sifted bark of tree” (HNHM; 1 female, 1 specimen – cMG).

Remarks. The species was known from Nepal. It is here reported from Vietnam for the first time. The last record confirms the Oriental distribution of the species.

Carpelimus (s.str.) congruus (Cameron, 1930)

Material. Nepal: 3 males: “NEPAL, Prov. Narayani, Sauraha, Rapti River Ufer, 180 m NN, 27°34'80" N/ 84°29'49" E, LF, 18.IV.2000, leg. A.Weigel” (NKME; 1 male – cMG); 1 male, 2 females: “NEPAL: Chitwan NP Narayani Ganga, 18.4.1995 27°33'N 84°06'E 150m, leg. Malicky” (NHMW).

Remarks. The lectotype is designated from India (Gildenkov, 2013: 55). The species was known from India and Sri Lanka. *C. congruus* is here reported from Nepal for the first time.

Carpelimus (s.str.) indicus (Kraatz, 1859)

(*flavipes* Motschulsky, 1861; *ceylonicus* Bernhauer, 1902; *kreyenbergi* Bernhauer, 1928 – syn. in Gildenkov, 2014: 298)

Material. Nepal: 4 specimens: “NEPAL, P: Narayani, D: Chitwan, Sauraha, Hotel Riversid, 190 m NN – 26.VI-02.VII.2011, LFN 27°34'55" E 84°29'58" leg. M. Hartmann, #54” (NKME); 1 male: “NEPAL, Prov. Narayani, Sauraha, Rapti River, 180 m, 27°34'80" N/ 84°29'49" E, 14-15.VII.2001, leg. Weigel Ufer LF” (NKME); 1 female: “NEPAL, Prov. Koshi distr. ankhawasabha Tumlingtar, Arun-Ufer 27°19'08" N/ 87°10'05" E, 310 m NN, 26.XI.1998, leg. M.Hartman” (NKME); 1 specimen: “NEPAL, Prov. Koshi distr. Sankhuwasabha Tumlingtar, Arun-Ufer 27°19'08" N/ 87°10'05" E, 310 m NN, 26.XI.1998, leg. M.Hartman” (NME); 1 specimen: “NEPAL: 700', Chitwan Nat. Pk. Sauraha, 3-6.vi.1983.” “At light” “M.J.D. Brendell B.M. 1983-222” (BMNH).

Remarks. The lectotype was designated from India (Gildenkov, 2010: 13). The species is widely distributed in the Oriental region (Gildenkov, in lit.): Andaman islands, Cambodia, China (Fuzhou, Guangdong, Guangxi, Guizhou, Hongkong, Hunan, Yunnan, Zhejiang), India, Indonesia (Kalimantan, Sumatra, Java), Japan (Okinawa), Laos, Malaysia (including Kalimantan), Myanmar, Nepal, Philippines, Sri Lanka, Thailand, Taiwan, Vietnam. In the Palaearctic the species known only from Japan (Gildenkov, 2001: 77), China (N-Sichuan) (Gildenkov, 2009: 29), and Pakistan (Gildenkov, 2012: 354). *C. indicus* is here reported for Nepal for the first time.

Carpelimus (s. str.) insularis (Kraatz, 1858)
(spec. propr. *nigrita* auct. nec Wollaston, 1857 – in Schülke, 2004: 398; *calidus* Bernhauer 1943 – syn. in Gildenkov, 2007: 1076; *oculatus* Wollaston 1865 – syn. in Gildenkov, 2007: 1078; *meridioafricanus* Scheerpeltz 1974 – syn. in Gildenkov, 2007: 1078)

Coiffait, 1982: 43 (as *nigrita* Wollaston, 1857)

Remarks. The species was described from Greece; the neotype was designated from southern Spain (Gildenkov, 2007: 1076). It is well known from the Palaearctic: Morocco, Algeria, Tunis, Sudan (on border with Egypt), Spain, France, Italy (including Sicily and Sardinia), Greece, Bul-

garia, Moldova, Lebanon, Turkey, Iran, Iraq, Afghanistan, Russia (southern Dagestan), Azerbaijan, Georgia, Tadjikistan, Turkmenistan, Uzbekistan (Gildenkov, 2001: 94; 2009: 29). It is known from the Tropical Africa (Gildenkov, 2007: 1078; 2007b: 898): Angola, Botswana, Chad, Congo, Ethiopia, Ghana, Kenya, Madagascar, Mali, Namibia, Nigeria, the Republic of South Africa, Reunion, Senegal, Sudan, Tanzania, Uganda, Zambia, Zimbabwe. Recently it was recorded from Pakistan, Portugal and Kirghizia (our data). Any data about the distribution of this species in south-eastern Asia are not present. The record of *C. insularis* based on a single specimen by Coiffait (1982: 43) from Nepal cannot be confirmed herein and it seems to be doubtful, it more probably belonged to the widely distributed in Nepal and rather variable *C. peregrinus*.

Carpelimus (s. str.) peregrinus (Cameron, 1919) (*afghanicus* Coiffait, 1982 – syn. in Gildenkov, 2010: 14; *siamensis siwalikensis* Cameron, 1930 – syn. in Gildenkov, 2014: 300; *tagus* Herman, 1970 – syn. in Gildenkov, 2014: 300; *subpolitus* Coiffait, 1982 – syn. in Gildenkov, 2014: 300)

Coiffait, 1982: 43, 158 (as *Trogophloeus (Boopinus) subpolitus*)

Material. Nepal (as *subpolitus*): Holotypus, male: “Nepal 1977, Siklis, 2150 – B.408” “HOLOTYPE” “*C.subpolitus* (Coiffait, 1982) | rev. Gildenkov, 2007” “MUSEUM PARIS” (MNHN); 1 female: “Nepal 1977, Madi Khola, 2385 m – B.913” “PARATYPE” “*C.subpolitus* (Coiffait, 1982) | rev. Gildenkov, 2007” “MUSEUM PARIS” (MNHN).

Nepal (as *peregrinus*): 1 female: “NEPAL, P: Narayani, D: Chitwan, Sauraha, Hotel Riversid, 190 m NN – 26.VI-02.VII.2011, LFN 27°34'55", E 84°29'58" leg. M. Hartmann, #54" (NKME); 3 males, 1 female, 14 specimens: “NEPAL c. 2001 Prov. Narayani Sauraha, Rapti River, LF, 180m leg. Kopetz 15.VII.” “27°34'51"N 84°29'30"E" (NKME); 1 male – cMG); 1 male: “NEPAL, Prov. Narayani, Sauraha, Rapti River, 180 m, 27°34' 80"

N/ 84°29'49" E, 14-15.VII.2001, leg. Weigel Ufer LF” (cMG); 1 male, 1 female: “NEPAL, Prov. Narayani, Sauraha, Rapti River, 27°34'51" N/ 84°29'30" E, 14-15.VII.2001, 180 m riverside, leg. Kopetz” (NKME); 1 male, 3 females: “NEPAL, Kathmandu, N Bagmati River, 1300 m NN, 06.VI.1995, leg. Hartman” (NKME); 1 female: “NEPAL, Kathmandu Thamel, Hotel Norbhu Linkha, LF, 23.IX.1996, leg. Hartman” (NKME); 2 females: “NEPAL, Prov. Bheri, 15 km W Kusum, 28°02'36" N/ 81°34'35" E, 11-12.07.2001, LF, 170 m, townside, leg. A.Kopetz” (NKME); 1 female: “NEPAL, Prov. Narayani, Sauraha, Rapti River, 180 m LF, 27°34'80" N/ 84°29'49" E, 18.IV.2000, Part-Ufer, leg. Weigel” (NKME); 1 male: “NEPAL, valley of Trisuli Ganga, ca. 20 km NE Trisuli” “85°12'E, 28°03'N 900 m, 23.V.1995, Gy Fábian & L. Ronkay” (HNHM); 1 male: “NEPAL, valley of Trisuli Ganga, ca. 20 km NE of Trisuli, 900m” “28°3'N, 85°12'E, 23.V.1995, leg. Gy Fábian & L. Ronkay” (cMG); 3 specimens: “NEPAL, 700 feet Chitwan National Pk. Sauraha, 4.vi.1983 At MV Light, MJD Brendell BMNH {E} 1983-222” (BMNH); 1 male, 4 specimens: “NEPAL: 4500', Kathmandu, British Embassy, 20.v. – 23.vi.1983.” “M.J.D. Brendell B.M. 1983-222” “At light” (cMG); 1 specimen: “NEPAL: 4500', Kathmandu Dist. Gokarna, 1.viii.1983.” “Mixed primary forest” “Lt. Col. M.G. Allen Brit. Mus. 1983-254” (BMNH); 1 male, 1 specimen: “NEPAL: 700', Chitwan Nat. Pk. Sauraha, 3-6.vi.1983.” “At MV light” “M.J.D. Brendell B.M. 1983-222” (BMNH); 1 male, 2 females, 15 specimens: “NEPAL: 4500', Kathmandu, British Embassy, 20.v. – 23.vi.1983.” “M.J.D. Brendell B.M. 1983-222” “At MV light” (BMNH; 2 specimens – cMG); 2 females, 11 specimens: “NEPAL: 4500', Kathmandu, British Embassy, 20.v. – 23.vi.1983.” “M.J.D. Brendell B.M. 1983-222” “At light” (BMNH); 1 male, 4 females: “600 NEPAL: Kathmandu Baneshwar 1350m, 20.-21.V.2000 leg. W. SCHAWALLER” (SMNS); 1 female: “314 Ilan Distr., Mai Khola Valley, 470m, on river bank, bushes, betw. pebbles, 7 April 1988, J. MARTENS & W.SCHAWALLER leg.” (SMNS); 1 male, 2 females: “NEPAL: 27°52'N 84°38'E 17.4.1995, Trisuli Ganga nr. Kurin Ghat, 300m leg. Malicky” (NHMW).

Remarks. The lectotype is designated from Sri Lanka (Gildenkov, 2010: 14). The species is known from the Oriental region: Andaman islands, Cambodia, China (Shaowu), India, Indonesia (Kalimantan, Sumatra, Java), Malaysia (including Kalimantan), Myanmar, Nepal, Singapore, Taiwan, Thailand, Vietnam (our data). Recently it is recorded from Seychelles islands (our data). As *C. siamensis* it is known from the Palaearctic from Afghanistan (Gildenkov, 2009: 34) and Pakistan (Gildenkov, 2012: 355).

Carpelimus (Troginus) atomus (Saulcy, 1864) (*discolor* Baudi di Selve, 1870; *variegatus* Cameron 1944 – syn. in Gildenkov, 2007a: 1318; *ruandanus* Cameron 1956 – syn. in Gildenkov, 2007a: 1318; *maroccanus* Gildenkov 2004 – syn. in Gildenkov, 2007a: 1318; *travei* Coiffait, 1982 – syn. in Gildenkov, 2014: 304; *formosanus* Cameron, 1940 – syn. in Gildenkov, 2014: 304; *atomarius* Coiffait, 1983 – syn. in Gildenkov, 2014: 304)

Coiffait, 1982: 43, 158 (as *Trogophloeus (Paraboopinus) travei*)

Material. Nepal (as *travei*): Holotypus, male (in the description it is erroneous – female) “Nepal 26.X.76 env. Pokhara 2’200 m 74” “HOLOTYPE” “*C.travei* (Coiffait, 1982) | rev. Gildenkov, 2007” “MUSEUM PARIS” (MNHN).

Nepal (as *atomus*): 1 male, 3 specimens: “NEPAL, P: Narayani Chitwan, Gunganagar V.2005, 200m 27°39’N, 84°19’E leg. D. Ahrens” (NKME); 1 male, 7 specimens: “NEPAL c. 2001 Prov. Narayani Sauraha, Rapti River, LF, 180m leg. Kopetz 15.VII.” “27°34’51”N 84°29’30”E” (NKME); 4 specimens: “NEPAL, Prov. Narayani Sauraha, Rapti River Ufer, 180m NN, 27°34’80”N, 84°29’49”E LF, 18.IV.2000 leg. A. Weigel” (NKME); 2 males, 7 specimens: “NEPAL, Narayani Prov. Sauraha, Chitwan Nat. P., 180 m LF, 27°34’80” N/ 84°29’49” E, 18.IV.2000, Parti-Ufer, leg. Weigel” (NKME); 1 specimen: “NEPAL, Prov. Bheri, Distr. Nepalganj, 40 km N Nepalganj, Babai-River 500 m NN, 03.VI.1995 Ufer, leg. M.Hartman” (NKME); 1 specimen: “NEPAL: 4500’, Kathmandu Dist.

Gokarna, 1.viii.1983.” “Mixed primary forest” “Lt. Col. M.G. Allen Brit. Mus. 1983-254” (BMNH); 11 specimens: “NEPAL: Chitwan NP Narayani Ganga, 18.4.1995 27°33’N 84°06’E 150m, leg. Malicky” (NHMW); 1 specimen: “NEPAL: Kabhre dist. Bebar, 1.12.1999 Bebar Khola, 1514m 27°35’00”N 85°29’50”E leg. S. Khanal (6)” (NHMW).

Remarks. The species is widely distributed in the Oriental region: China (Guangdong, Guizhou, Hongkong, Hubei, Hunan, Yunnan, Zhejiang), India, Indonesia (Kalimantan, Sumatra, Java), Japan (Okinawa), Cambodia, Malaysia (including Kalimantan), Nepal, Philippines, Sri Lanka, Taiwan, Thailand, Vietnam (our data). *C. atomus* is known from the tropical Africa: Angola, Benin, Botswana, Burundi, Cameroon, the Central-African Republic, Congo, Cote d’Ivoire, Ethiopia, Gambia, Ghana, Madagascar, Mauritius, Namibia, Nigeria, the Republic of South Africa, Ruanda, Senegal, Seychelles islands, Sudan, Tanzania, Uganda, Yemen, Zambia, Zimbabwe (Gildenkov, 2007a: 1318; 2007c: 910). Besides that, the species is many times was recorded from the Palaearctic: Cyprus, Syria (Gusarov, 1997: 280), Egypt, Libya, Saudi Arabia (Gusarov, 1997: 280; Gildenkov, 2001: 277; 2003: 380; 2003a: 97; 2009a: 252), Iraq (Gildenkov, 2001: 277; 2003: 380; 2003a: 97), China (Hebei), Japan, South Korea (Gildenkov (as *formosanus*), 2001: 282, 2003: 381; 2003a: 97), Morocco (Gildenkov (as *maroccanus*), 2001: 280, 2003: 381; 2003a: 97), Turkey (Gildenkov, 2007a: 1318; 2007c: 910), Afghanistan (Gildenkov, 2009a: 252) and Pakistan (Gildenkov, 2012: 356). Recently it was recorded from Canary islands, China (N-Sichuan, Peking, Shaanxi), Malta, Madiera, Spain, Tunis (our data). Such distribution confirms the species more likely to belong to the fauna of tropical Africa and Oriental area, than to the fauna of Palaearctic.

Carpelimus (Troginus) despectus (Baudi di Selve, 1870) (*despectus* Mulsant & Rey, 1870; *despectus* var. *leederi* Bernhauer, 1940)

Coiffait, 1982: 44

Remarks. The species is known from the Palaearctic: Austria, Germany, Italy, Poland, Russia (the West and the Center of the European part (Moscow Area, Smolensk Area, Yaroslavl Area), Western Siberia (Tyumen Area)), Switzerland (Gildenkov, 2001: 271; 2009a: 253), Russia (Eastern Siberia (Irkutsk Area)) (Gildenkov & Shavrin, 2012: 53), Byelorussia (Gildenkov & Derunkov, 2012: 62). Recently it was recorded for the Ukraine (our data). It is a species difficult to identify. To use faunistic data of other authors about *C. despectus* outside Europe must be followed with great care. The record of *C. despectus* from Nepal cannot be considered confirmed and it seems to me doubtful because a single specimen of *C. despectus*, specified by Coiffait more probably belonged to the widely distributed in Nepal and rather morphologically variable *C. atomus*.

Carpelimus (Troginus) exiguus (Erichson, 1839) (*glabricollis* Motschulsky, 1860; *luteicornis* Mulsant & Rey, 1878; *aberrans* Rosenhauer, 1856)

Coiffait, 1982: 44

Remarks. The lectotype was designated from Europe (Gildenkov, 1998: 127). The species is widely distributed in Palaearctic: Austria, Bulgaria, Georgia, Germany, Italy, Kazakhstan, Moldova, Romania, Russia (all territory, including the West and the Center of the European part, the Volga region, Northern Caucasus, Eastern Siberia (Krasnoyarsk Territory, Irkutsk Area), the Russian Far East (Sakhalin, Maritime Province)), Turkey, Ukraine (Gildenkov, 2001: 265; 2009a: 253; Gildenkov & Shavrin, 2012: 54), Byelorussia (Gildenkov & Derunkov, 2012: 62). Recently it was recorded from Spain, Kirghizia and Tadjikistan (our data). It is difficult for identification species. To use faunistic data of other authors about *C. exiguus* outside Europe must be followed with great care. The record of species based on the single specimen for Nepal by Coiffait (1982) is not confirmed here and seems to be doubtful, it is probably *C. atomus* which is very variable morphologically. Previously, the

distribution of *C. exiguus* was understood as worldwide, now this does not have any confirmations, it is obvious that *C. exiguus* did not differ with *C. atomus*.

Carpelimus (Trogophloeus) bengalensis (Cameron, 1930)

Material. Nepal: 1 male: "NEPAL, Narayani Prov. Sauraha, Chitwan Nat. P., 180 m LF, 27°34'80" N/84°29'49" E, 18.IV.2000, Parti-Ufer, leg. Weigel" (cMG).

Remarks. The lectotype was designated from India (Gildenkov, 2013: 54). *C. bengalensis* is here reported for Nepal for the first time.

Carpelimus (Trogophloeus) bertrandi (Coiffait, 1982)

Coiffait, 1982: 44, 159

Material. Nepal: Holotypus, male (under the description, aedeagus is not present) "Nepal 1977, Madi Khola, 970 m – B.508" "Preparation microscopique" "HOLOTYPE" "*Carpelimus bertrandi* (Coiffait, 1982) | rev. Gildenkov, 2007" "MUSEUM PARIS" (MNHN); 1 male: "Nepal, 4-XI-76. Col Argan" "PARATYPE" "*Carpelimus bertrandi* (Coiffait, 1982) | rev. Gildenkov, 2007" "MUSEUM PARIS" (MNHN).

Remarks. It is known only from Nepal.

Carpelimus (Trogophloeus) brachypterus (Coiffait, 1982)

(*apterus* Coiffait, 1982 – syn. in Gildenkov, 2014: 308)

Coiffait, 1982: 44, 160; 1982: 44, 160 (as *apterus*)

Material. Nepal (as *brachypterus*): Holotypus, female: "Nepal 19.X.77 | Boudanikantar, 1550 m LD-13" "HOLOTYPE" "*C. brachypterus* (Coiffait, 1982) | rev. Gildenkov, 2007" "MUSEUM PARIS" (MNHN); 1 female: "Nepal IX-X.77 |

Boudanikantar, 1550 m LD-13" "PARATYPE" "*C. brachypterus* (Coiffait, 1982) | rev. Gildenkov, 2007" "MUSEUM PARIS" (MNHN).

Nepal (as *apterus*): Holotypus, female: "Nepal 26.X.76 env. Pokhara 2'200 m 74" "HOLOTYPE" "*Carpelimus apterus* (Coiffait, 1982) syn.n. = *C. brachypterus* (Coiffait, 1982) det. Gildenkov, 2007" "MUSEUM PARIS" (MNHN).

Remarks. It is known only from Nepal.

Carpelimus (Trogophloeus) corticinus (Gravenhorst, 1806)

(*atratus* Stephens, 1834; *minimus* Runde, 1835; *dispersepunctatus* Scheerpeltz, 1947; *nanus* Wollaston, 1854)

Material. Nepal: 1 female: "NEPAL, Kathmandu, N Bagmati River, 1300 m NN, 27.IV.1995, leg. M.Hartman" (NKME).

Remarks. The lectotype was designated from Europe (Gildenkov, 2004: 542; 2004a: 612). The species is widely distributed in Palaearctic region: Albania, Austria, Azerbaijan, Bosnia-Herzegovina, Bulgaria, Croatia, Czech Republic, Estonia, France, Georgia, Germany, Great Britain, Greece, Hungary, Iraq, Italy, Japan, Kazakhstan, the Canary islands, Crete, Latvia, Lithuania, Madiera, Morocco, Moldova, Mongolia, Northern Korea, Poland, Russia (all territory, including the European part, Caucasus, Eastern Siberia and the Far East), Serbia, Spain, Sweden, Tadjikistan, Tunis, Turkmenistan, Turkey, Ukraine, Uzbekistan (Gildenkov, 2001: 176; Gildenkov & Shavrin, 2012: 54), Armenia (Gildenkov, 2009a: 240), Byelorussia (Gildenkov & Derunkov, 2012: 61). Recently it was recorded from Afghanistan, China (Heilungkiang, Peking, Shaanxi), Denmark, Egypt, Iran, Kirghizia, Lebanon (our data). It is known from the Oriental area: China (Yunnan), Nepal and Northern India. *C. corticinus* is here reported for Nepal for the first time.

Carpelimus (Trogophloeus) palitans (Cameron, 1930)

(*gratus* Cameron, 1930 – syn. in Gildenkov, 2014: 309)

Material. Nepal: 1 female: "NEPAL – Chitwan Nat. Park: Sauraha, 100 m 28.IX.92, L.Schmidt" (cVA); 1 male: "NEPAL, P: Narayani, D: Chitwan, Sauraha, Hotel Riversid, 190 m NN – 26.VI-02.VII.2011, LFN 27°34'55", E 84°29'58" leg. M. Hartmann, #54" (NKME); 2 males, 1 female, 10 specimens: "NEPAL c. 2001 Prov. Narayani Sauraha, Rapti River, LF, 180m leg. Kopetz 15.VII." "27°34'51"N 84°29'30"E" (NKME; 1 male – cMG); 1 male, 2 females, 3 specimens: "NEPAL, Prov. Bheri Nepalgunj Hotel, Batika, 28°02'59" N/ 81°36'56" E, 230 m NN, 18.VI.1999, LF, leg. M.Hartman" (NKME; 1male – cMG); 2 males, 3 females: "NEPAL, Prov. Bheri, 15 km W Kusum, 28°02'16" N/ 81°57'16" E, 13.VII.2001, leg. A.Kopetz, deciduous wood KL/HF" (NKME); 1 female: "NEPAL, Prov. Bheri, 15 km W Kusum, 28°02'36" N/ 81°34'35" E, 11-12.07.2001, LF, 170 m, townside, leg. A.Kopetz" (NKME); 1 male, 1 female: "NEPAL, Prov. Narayani, Sauraha, Rapti River, 180 m LF, 27°34'80" N/ 84°29'49" E, 18.IV.2000, Parti-Ufer, leg. Weigel" (NKME); 2 specimens: "NEPAL, 700 feet Chitwan National Pk. Sauraha, 4.vi.1983 At MV Light, MJD Brendell BMNH {E} 1983-222" (BMNH).

Remarks. The lectotype was designated from India (Gildenkov, 2013: 59). The species is known from the Oriental region: China (Fuzhou), India, Indonesia (Sumatra, Java), Malaysia (including Kalimantan), Nepal, Singapore, Sri Lanka, Thailand (our data). *C. palitans* is here reported for Nepal for the first time.

Carpelimus (Trogophloeus) vagans (Cameron, 1930)

Coiffait, 1982: 43

Remarks. India: 1 male: "Kaligad, Dehra Dun. 10. VII. 21" "Syntypus (red label)" "*T. vagans* Cam" "*Carpelimus vagans* (Cameron) | det. M.Gildenkov, 1996" "*Carpelimus vagans* (Cameron, 1930) | det. M.Gildenkov, 2014" "DEI Müncheberg Col – 03432" (SDEI); 1 male "Kaligad, Dehra Dun. 10. VII. 21" "Cotypus don. Cameron (violet label)" "*T. vagans* Cam" "*Carpelimus vagans* (Cameron, 1930) | det.

M. Gildenkov, 2014” “Chicago NHMus M. Bernhauer Collection” (FMNH).

C. vagans refer to a numerous group of species “*taprobanae*” which are very difficult to determine (Gildenkov, 2013b(2014): 292). The discovery of *C. vagans* in Nepal is quite probable, but cannot be considered to be confirmed at the present time.

DISCUSSION.

Thus, twelve species of the genus *Carpelimus* are known from Nepal. The finding of *Carpelimus* (*Trogophloeus*) *vagans* (Cameron, 1930) in Nepal is very expected.

Six species are new for the fauna of Nepal: *Carpelimus* (*Bucephalinus*) *laticeps* (Cameron, 1930); *Carpelimus* (s.str.) *congruus* (Cameron, 1930); *Carpelimus* (s.str.) *indicus* (Kraatz, 1859); *Carpelimus* (*Trogophloeus*) *bengalensis* (Cameron, 1930); *Carpelimus* (*Trogophloeus*) *corticinus* (Gravenhorst, 1806); *Carpelimus* (*Trogophloeus*) *palitans* (Cameron, 1930). Two species have been known for Nepal before, but under other names that have been transferred by the author to synonyms: *Carpelimus* (s. str.) *peregrinus* (Cameron, 1919) = *Trogophloeus* (*Boopinus*) *subpolitus* Coiffait, 1982, syn.; *Carpelimus* (*Troginus*) *atomus* (Saulcy, 1864) = *Trogophloeus* (*Paraboopinus*) *travei* Coiffait, 1982, syn.

Three species from thirteen are known only from Nepal: *Carpelimus* (*Bucephalinus*) *kathmanduensis* Herman, 2001; *Carpelimus* (*Trogophloeus*) *bertrandi* (Coiffait, 1982); *Carpelimus* (*Trogophloeus*) *brachypterus* (Coiffait, 1982). One species (*C. corticinus*) has a palaeartic distribution, other species (*C. atomus*) is widely distributed in Eastern and Afrotropical regions, and penetrates from the south to the Palaeartic. Other eight species (six species from them are recorded here for Nepal for the first time) belong to the Oriental fauna. It confirms our opinion that fauna of the species of the genus *Carpelimus* of Nepal should be considered within the Oriental region.

ACKNOWLEDGEMENTS

The author wishes to thank all colleagues for making material available for study: Mr. Martin Brendell, Dr. Roger G. Booth (London), Dr. Volker Assing (Hanover); Dr. Alfred F. Newton, Dr. Margaret K. Thayer, Mr. James H. Boone (Chicago); Dr. György Makranczy (Budapest); Marc Tronquet (Molitg-les-Bains); Dr. Azadeh Taghavian (Paris); Dr. Harald Schillhammer (Vienna); Dr. Matthias Hartmann (Erfurt); Dr. Lothar Zerche (Müncheberg); Dr. Wolfgang Schawaller (Stuttgart); Dr. Alexey V. Shavrin (Daugavpils).

REFERENCES

- Coiffait H. 1982. Contribution a la connaissance des Staphylinides de l’Himalaya (Nepal, Ladakh, Cachemire) (Insecta: Coleoptera: Staphylinidae). *Senckenbergiana Biologica* 62: 21–179.
- Coiffait H. 1984. Contribution a la connaissance des Staphylinides de l’Himalaya (Coleoptera, Staphylinidae). *Annales de la Societe entomologique de France* (n. ser.) 20(4): 373–387.
- Gildenkov M. Yu. 1998. *Carpelimus minimus* (Kraatz, 1859) as a proprius species, a key to the Oriental species of the subgenus *Troginus* (Coleoptera: Staphylinidae: *Carpelimus*). *Russian Entomological Journal* 7 (3–4): 127–128.
- Gildenkov M. Yu. 2001. Fauna *Carpelimus* Palearktiki (Coleoptera, Staphylinidae). *Problemy vida i vidoobrazovaniya. Chast’ 1. Istoriya izucheniya. Morfo-ekologicheskie osobennosti. Sistema roda. Opisanie vidov* [The fauna of *Carpelimus* of the Palaeartic (Coleoptera, Staphylinidae). Problems of species and speciation. Part 1. The history of research. Morpho-ecological features. System of the genus. Description of species]. Smolensk: SSPU. 304 pp. [in Russian].

- Gildenkov M. Yu. 2002. Sistema roda *Carpelimus* (Coleoptera, Staphylinidae) v predelakh Palearktiki. 1. Obshchaya chast' i opisanie novykh taksonov [Classification of the genus *Carpelimus* (Coleoptera, Staphylinidae) within the Palaearctic Region: 1. Introduction and description of new taxa]. *Zoologicheskij zhurnal* 81(12): 1461–1475. [in Russian; Abstract in English].
- Gildenkov M. Yu. 2002a. Classification of the Genus *Carpelimus* (Coleoptera, Staphylinidae) within the Palaearctic Region: 1. Introduction and description of new taxa. *Entomological Review* 82 (9): 1274–1288.
- Gildenkov M. Yu. 2003. Sistema roda *Carpelimus* (Coleoptera, Staphylinidae) v predelakh Palearktiki. 3. Klyuchi dlya opredeleniya podrodov i vidov podrodov *Trogophloeus* è *Troginus* [Classification of the genus *Carpelimus* (Coleoptera, Staphylinidae) within the Palaearctic Region: 3. Keys to species of the subgenera *Trogophloeus* and *Troginus*]. *Zoologicheskij zhurnal* 82(3): 366–381. [in Russian; Abstract in English].
- Gildenkov M. Yu. 2003a. Classification of the genus *Carpelimus* (Coleoptera, Staphylinidae) within the Palaearctic Region: 3. Keys to species of the subgenera *Trogophloeus* and *Troginus*. *Entomological Review* 83 (1): 82-97.
- Gildenkov M. Yu. 2004. Novye i maloizvestnye palearkticheskie vidy roda *Carpelimus* (Coleoptera, Staphylinidae) [New and little-known Palaearctic species of the Staphylinid genus *Carpelimus* Leach (Coleoptera, Staphylinidae)]. *Entomologicheskoe obozrenie* 83 (3): 538–552. [in Russian; Abstract in English].
- Gildenkov M. Yu. 2004a. New and little-known Palaearctic species of the Staphylinid genus *Carpelimus* Leach (Coleoptera, Staphylinidae). *Entomological Review* 84(5): 609–620.
- Gildenkov M. Yu. 2007. Obzor fauny podroda *Carpelimus* (s.str.) (Coleoptera, Staphylinidae) Tropicheskoj Afriki [A review of the subgenus *Carpelimus* s. str. (Coleoptera, Staphylinidae) from Tropical Africa]. *Zoologicheskij zhurnal* 86 (9): 1073–1085. [in Russian; Abstract in English].
- Gildenkov M. Yu. 2007a. Obzor fauny podroda *Troginus* roda *Carpelimus* (Coleoptera, Staphylinidae) Tropicheskoj Afriki [A review of the fauna of the subgenus *Troginus* (*Carpelimus*, Coleoptera, Staphylinidae) from Tropical Africa]. *Zoologicheskij zhurnal* 86(11): 1315–1327. [in Russian; Abstract in English].
- Gildenkov M. Yu. 2007b. A Review of the Subgenus *Carpelimus* s. str. (Coleoptera, Staphylinidae) from Tropical Africa. *Entomological Review* 87 (7): 893–907.
- Gildenkov M. Yu. 2007c. A Review of the Fauna of the Subgenus *Troginus* (*Carpelimus*, Coleoptera, Staphylinidae) from Tropical Africa. *Entomological Review* 87 (7): 907–919.
- Gildenkov M. Yu. 2009. Novye dannye o rasprostranenii v Palearktike vidov roda *Thinodromus* i vidov roda *Carpelimus* iz podrodov *Carpelimus* s.str., *Paratrogophloeus*, *Bucephalinus* (Coleoptera, Staphylinidae) [The new data on the distribution in Palaearctic Region of species of the genus *Thinodromus* and species of the genus *Carpelimus* from subgenera *Carpelimus* s.str., *Paratrogophloeus*, *Bucephalinus* (Coleoptera, Staphylinidae)]. *Izvestiya Smolenskogo Gosudarstvennogo Universiteta* 2 (6): 25"42. [in Russian; Abstract in English].

- Gildenkov M. Yu. 2009a. Novye dannye o rasprostraneni v Palearktike vidov roda *Carpelimus* iz podrodov *Myopinus*, *Trogophloeus*, *Troginus* (Coleoptera, Staphylinidae) [The new data on the distribution in the Palaearctic Region of species of the genus *Carpelimus* from subgenera *Myopinus*, *Trogophloeus*, *Troginus*]. *Izvestiya Smolenskogo Gosudarstvennogo Universiteta* 3 (7): 237–255. [in Russian; Abstract in English].
- Gildenkov M. Yu. 2010. Novaya sinonimiya i nomenklaturnye tipy dlya palearkticheskikh vidov iz rodov *Thinodromus* i *Carpelimus* (Coleoptera, Staphylinidae, Oxytelinae) [New synonymy and nomenclative types for the Palaearctic species from the genera *Thinodromus* and *Carpelimus* (Coleoptera, Staphylinidae, Oxytelinae)]. *Izvestiya Smolenskogo Gosudarstvennogo Universiteta* 4(12): 7–29. [in Russian; Abstract in English].
- Gildenkov M. Yu. 2012. Novye dannye po taksonomii i rasprostraneniyu roda *Carpelimus* Leach, 1819 v severozapadnom Pakistane (Coleoptera, Staphylinidae, Oxytelinae) [The new data on taxonomy and distribution of the genus *Carpelimus* Leach, 1819 in northwest Pakistan (Coleoptera, Staphylinidae, Oxytelinae)]. *Izvestiya Smolenskogo Gosudarstvennogo Universiteta* 2 (18): 351–357. [in Russian; Abstract in English].
- Gildenkov M. Yu. 2013. Designations of the lectotypes for the oriental species of the genus *Carpelimus* Leach, 1819 (Coleoptera: Staphylinidae). *Zootaxa* 3717 (1): 53–64.
- Gildenkov M. Yu. 2013a. Obzor sostava podroda *Trogophloeus* Mannerheim, 1930 iz roda *Carpelimus* Leach, 1819 dlya Tropichestkoj Afriki (Coleoptera, Staphylinidae, Oxytelinae) [The review of the structure of the subgenus *Trogophloeus* Mannerheim, 1930 of the genus *Carpelimus* Leach, 1819 for Tropical Africa (Coleoptera, Staphylinidae, Oxytelinae)]. *Izvestiya Smolenskogo Gosudarstvennogo Universiteta* 1 (21): 242–251. [in Russian; Abstract in English].
- Gildenkov M. Yu. 2013b. Novye vidy roda *Carpelimus* Leach, 1819 iz grupp «*silvestris*» i «*taprobanae*» (Coleoptera, Staphylinidae, Oxytelinae) [New species of the genus *Carpelimus* Leach, 1819 from the groups «*silvestris*» and «*taprobanae*» (Coleoptera, Staphylinidae, Oxytelinae)]. *Izvestiya Smolenskogo Gosudarstvennogo Universiteta* 4 (24): 292–305. [in Russian; Abstract in English].
- Gildenkov M. Yu. 2014. Novaya sinonimiya i nomenklaturnye tipy dlya vidov roda *Carpelimus* Leach, 1819 (Coleoptera, Staphylinidae, Oxytelinae) [New synonymy and nomenclature types for species from the genus *Carpelimus* Leach, 1819 (Coleoptera, Staphylinidae, Oxytelinae)]. *Izvestiya Smolenskogo Gosudarstvennogo Universiteta* 1 (25): 296–314. [in Russian; Abstract in English].
- Gildenkov M. Yu., Derunkov A.V. 2012. Novye dannye o rasprostraneni predstavitelej roda *Carpelimus* Leach, 1819 (Coleoptera: Staphylinidae: Oxytelinae) v Belarusi [The new data on the distribution of the genus *Carpelimus* Leach, 1819 (Coleoptera: Staphylinidae: Oxytelinae) in Belarus]. *Ecologiya, evolutsiya i sistematika zhivotnykh: materialy Mezhdunarodnoj nauchno-prakticheskoy konferentsii (Rossiya, Ryazan, 13–16 noyabrya 2012)*. Ryazan: NP “Golos Gubernii”: 61–62. [in Russian].
- Gildenkov M. Yu., Shavrin A.V. 2012. New records of *Ochtheophilus* Mulsant & Rey, 1856, *Carpelimus* Leach, 1819 and *Thinodromus* Kraatz, 1857 (Coleoptera: Staphylinidae):

Oxytelinae) from Cis- and Transbaikalia.
Acta Biologica Universitatis
Daugavpiliensis 12 (3): 51–57.

Received: 16.05.2014.
Accepted: 02.06.2014.

Gusarov V. 1997. Staphylinids (Coleoptera: Staphylinidae) from Saudi Arabia and Oman. *Fauna of Saudi Arabia* 16: 277–290.

Herman L. H. 1970. Phylogeny and reclassification of the Genera of the rove-beetle Subfamily Oxytelinae of the World (Coleoptera, Staphylinidae). *Bulletin of the American Museum of Natural History* 142 (5): 343–454.

Smetana A. 2004. Tribe Thinobiini J. Sahlberg, 1876. In: Löbl I. & Smetana A. (Eds.): *Catalogue of Palaearctic Coleoptera. II. Hydrophiloidea – Histeroidea – Staphylinoidea*. Apollo Books, Stenstrup: 519–535.

Schülke M. 2004. Ueber einige von T.V. Wollaston von Inseln des Madeira – Archipels beschriebene Staphyliniden–Arten (Coleoptera: Staphylinidae: Omaliinae, Oxytelinae et Staphylininae). *Linzer biologische Beiträge* 36 (1): 393–415.

