

Caffeine Content Investigation in the Turkish Black Teas

E. Moroydor Derun, A. S. Kipcak, O. Dere Ozdemir, F. Demir, M. Karakoc, S. Piskin

Abstract—Tea is a widely consumed beverage that contains many components. Caffeine belongs to this group of components called alkaloids contain nitrogen. In this study caffeine contents of three types of Turkish teas are determined by using extraction method. After condensation process, residue of caffeine and oil are obtained with evaporation. The oil which is in the residue is removed by hot water. Extraction process performed by using chloroform and the crude caffeine is obtained. From the results of experiments, caffeine contents are found in black tea, green tea and earl grey tea as $3.57\pm 0.43\%$, $3.11\pm 0.02\%$, $4.29\pm 0.27\%$, respectively. Caffeine contents which are found in 1, 5 and 10 cups of tea are calculated. Furthermore, the daily intake of caffeine from black teas that affects human health is investigated.

Keywords—Caffeine, extraction, tea, health.

I. INTRODUCTION

TEA is one of the most widely popular nonalcoholic beverage that comes after water, which are produced about 2.5 million tons of tea in the world every year. Tea is made from the processed leaves of a plant, *Camellia sinensis* (Fig. 1 (a)). Tea is the agricultural product of the leaves, leaf buds, and internodes of the *Camellia sinensis* plant, prepared and cured by various methods [1].

Green tea (Fig. 1 (b)) and black tea (Fig. 1 (c)) are the two most popular types of Turkish Teas. The leaves of green tea are dried and roasted but not fermented, whereas black tea leaves are additionally fermented. In the fermentation, the enzymatic oxidation of tea polyphenols takes place leading to the formation of chemical compounds, which are responsible for the characteristic aroma, and color of teas [2].

To produce green tea, the leaves are chopped, rolled and quickly steamed or heated which inactivates polyphenol oxidase. For black tea the leaves are kept warm for 6 hours and the leaf polyphenols, especially the catechins, are oxidized and condensed. Tea drink is prepared by adding the dried leaf in hot or boiling water. It has a cooling slightly bitter astringent flavor which many people enjoy [3].

E. Moroydor Derun, is with the Yildiz Technical University, Department of Chemical Engineering, Davutpasa Campus, 34210 Esenler, Istanbul, Turkey (phone: 0090-212-3834776; fax: 0090-212-3834725; e-mail: moroydor@yildiz.edu.tr, moroydor@gmail.com).

A. S. Kipcak, O. Dere Ozdemir, F. Demir, M. Karakoc, and S. Piskin are with the Yildiz Technical University, Department of Chemical Engineering, Davutpasa Campus, 34210 Esenler, Istanbul, Turkey (e-mail: skipcak@yildiz.edu.tr, odere@yildiz.edu.tr, demirfundal@hotmail.com, karakocmerve@gmail.com, piskin@yildiz.edu.tr.).



Fig. 1 (a) *Camellia sinensis* (b) black tea (c) green tea [4]

Earl Grey tea is kind of specific tea. Earl Grey tea refers more to the bergamot oil that is used to flavor the tea than the tea base itself. Bergamot is used in Earl Grey tea, and is also often used in aroma therapy, and is considered an essential oil. Bergamot is a citrus fruit, the size of an orange but with an appearance more like a lemon; it is often called the bergamot orange. Bergamot essential oil stimulates the work of the brain, improves concentration and increases immunity [5].

Caffeine (Fig. 2) is a naturally occurring alkaloid which is found in the leaves, seeds or fruits of over 63 plants species worldwide [6]. It is often found in natural products such as tea, coffee and cocoa beans, cola nuts and many others [7]. Caffeine belongs to a large group of natural products that contain nitrogen and have basic properties. This group of natural products is called alkaloids [8].

The amount of caffeine in products varies depending upon the serving size, the type of product, and the preparation method [9].




Fig. 2 Structure of caffeine molecule [10]

Caffeine acts as a stimulant, where as it stimulates the heart, respiration and the central nervous system. It is also a diuretic molecule. Ingesting caffeine generally produces a sensation of heightened alertness. Caffeine is added to soft drinks as a flavoring agent; it imparts a bitterness that modifies the flavors of other components, both sour and sweet [11]. The caffeine content of teas varies from 1-5%.

The purpose of this study is to determine the contents of caffeine in the black, earl grey and green teas.

II. EXPERIMENTAL PROCEDURE

A. Preparation of the Tea Samples

Black tea, earl grey tea and green teas are purchased from the local market in Istanbul, Turkey. Teas are used in the analysis without pre-treatment

B. Caffeine Analysis

Extraction method is used for the analysis of caffeine. In this method, ammonia solution is poured to tea in the flask. After a few minutes, chloroform solution is added to the flask.

Then a condenser is placed on the flask and allowed to stand on the water bath for 30 minutes which is shown in Fig. 3.



Fig. 3 Condensation process

After the condensation process, the mixture is filtered into another flask. First flask is rinsed two times with chloroform solution in order to take all of the teas. The filtrate is evaporated on a water bath. After the evaporation, residue of caffeine and oil is obtained which is shown in Fig. 4.




Fig. 4 Residue of caffeine and oil

To separate oil from the mixture hot pure water is added to the residue. The mixture is heated in a water bath while stirring it gently from time to time with a glass rod and cooled to a room temperature. Potassium permanganate solution is added into the flask and left at room temperature for a few minutes. Then hydrogen peroxide solution is added to remove the color of the solution. If the mixture was still colored the addition of

hydrogen peroxide solution is continued. The mixture is heated in a water bath for 15 minutes and allowed to cool to a room temperature. After cooling, mixture is filtered using a filter paper. The flask is washed three times with distilled water, filtered and the filtrate is collected. Fig. 5 shows the filtration process.




Fig. 5 The filtration process

Filtrate is poured into a separation funnel and rinsed by adding chloroform solution to separate two phases (Fig. 6). The bottom phase is taken into a beaker.




Fig. 6 Two phase in the separator funnel

The collected chloroform solutions are evaporated in water bath and pure caffeine is obtained. In Fig. 7 the crude caffeine is shown.



Fig. 7 Crude caffeine

III. RESULTS AND DISCUSSION

Analysis results are shown in Table I. From the results obtained the maximum caffeine content is seen in the black tea, and followed by green tea and earl grey tea. The average caffeine content in green tea is analyzed as 3.11%. Also the average caffeine content in earl grey and black tea is analyzed

as 3.57 %, 4.29 %, respectively. As can be seen from the table, caffeine content in the earl grey tea is the highest and it is followed by black tea and green tea.

Tea Type	Caffeine Content (%)
Black Tea	3,57±0,43
Green Tea	3,11±0,02
Earl Grey Tea	4,29±0,27




Fig. 8 Caffeine content of teas

IV. CONCLUSIONS

Human daily maximum caffeine intake is given in Table II.

Recommended Maximum Daily Caffeine Intake (mg)	
4-6 Years	45
7-9	60
10-12	85
Women of Childbearing Age	300
Adults	400-450

The green teas and black teas have received a great deal of attention from researchers and the general public due to their possible beneficial health effects. Caffeine is one of the most comprehensively studied ingredients in the food supply. The per capita consumption level of caffeine for all consumers (of all ages) is approximately 120 mg per day, or a mean intake of 1.73 mg/kg body weight/day [11].

Children consume significantly less caffeine than adults. The average daily intake of caffeine by young children ages 1-5 and 6-9 years from all caffeinated beverages was 14 and 22 mg/day, or 0.82 and 0.85 mg/kg body weight/day, respectively [11].

Caffeine is a pharmacologically active substance and, depending on the amount consumed, can be a mild stimulant to the central nervous system. Consumption of caffeine prior to exercise has been shown to improve endurance during physical exercise [13].

From the results obtained in this study it can be said that both three types of teas caffeine content in 1 cup lower than daily maximum caffeine intake for adults. For black tea and green tea, amount of caffeine in 5 cups are lower than daily intake but caffeine content of earl grey tea is equal to maximum value. The value is shown in Tables III and IV.

TABLE III
DAILY CAFFEINE INTAKE

Daily Caffeine Intake (mg)			
	Black Tea	Green Tea	Earl Grey Tea
1 Cup	71,4	62,2	85,8
5 Cup	357	311	429
10 Cup	714	622	858

TABLE IV
DAILY CAFFEINE INTAKE FOR ADULTS

Daily Caffeine Intake for Adults (%)			
	Black Tea	Green Tea	Earl Grey Tea
1 Cup	16,26 – 17,85	14,04-15,55	19,06-21,45
5 Cup	81,33 – 89,25	70,2-77,75	95,3-107,25
10 Cup	162,6 – 178,5	140,4-155,5	190,6-214,5

For a tea addict person who drinks ten cups of black tea, green tea or earl grey tea takes more caffeine than daily maximum intake.

As seen that both three types of teas can reach the maximum daily caffeine intake.

REFERENCES

- [1] D. Desideri, M.A.Meli, C. Roselli, L. Feduzi, 2011, Polarized X Ray Fluorescence Spectrometer (EDPXRf) for the Determination of Essential and Non Essential Elements in Tea
- [2] Rachel R. McCusker, Bruce A. Goldberger, and Edward J. Cone, Caffeine Content of Specialty Coffees, Journal of Analytical Toxicology, Vol. 27, October 2003
- [3] M. A. Herrador, A. G. Gonzalez, 2000, Pattern Recognition Procedures for Differentiation of Green, Black and Oolong Teas According to Their Metal Content from Inductively Coupled Plasma Atomic Emission Spectrometry
- [4] <http://www.naturestealeaf.com/review/product/list/id/449/> retrieved on 20/03/2013
- [5] <http://coffeetea.about.com/od/typesoftea/a/Earl-Grey-Tea.htm>
- [6] Bibby Scientific The quantitative determination of caffeine in beverages and soft drinks using UV wavelength spectroscopy
- [7] Institute of Chemical Technology Prague, Direct analysis of caffeine in soft drinks and coffee and tea infusions.
- [8] NS207 Lab3- Isolation of Caffeine From Tea
- [9] K.L. Mandiwana, N. Panichev, S. Panicheva, 2011, Determination of Chromium(VI) in Black, Green and Herbal Teas
- [10] <http://tr.wikipedia.org/wiki/Kafein> retrieved on 15/03/2013
- [11] International Food Information Council Foundation, Caffeine & Health: Clarifying The Controversies
- [12] <http://www.milk.mb.ca/nutritionOrder/pdf> retrieved on 19/03/2013
- [13] K.L. Mandiwana, N. Panichev, S. Panicheva, 2011, Determination of Chromium(VI) in Black, Green and Herbal Teas



Emek Moroydor Derun was born in Istanbul in 1976. Moroydor Derun was graduated from B.Sc. in 1998, M.Sc. in 2000 and Ph. D. in 2005 from Chemical Engineering Department at Yildiz Technical University, Istanbul. Her research interest is in the area of waste management, lightweight concrete, semi conductive materials and boron technology. She has many articles and studies in international and national conference proceedings and articles.



Azmi Seyhun Kipcak was graduated from Department of Chemical Engineering in Ege University in 2002. After completing the university studies he graduated from Bilgi University from the department of Master of Business Administration in 2004. He worked in Kultur University from 2003 to 2007 as a research assistant then he transferred to Yildiz Technical University at 2008, where he started his M.Sc. studies about Chemical Engineering in 2006. He completed his M.Sc. and Ph.D. studies at Yildiz Technical University in 2009 and 2013, respectively. He studied on neutron shielding with boron minerals and the characterization of boron minerals by using XRD, XRF, FT-IR, Raman, DTA/TG, DSC and ICP-OES at the M.Sc. studies and studied on the synthesis of magnesium borates from different raw materials and wastes at the Ph.D. Also he is improving the neutron shielding studies with the synthesized materials and working on the element analysis of Turkish Teas and Coffees. Another research field about the studies he is working is the zinc borate synthesis.



Ozgul Dere Ozdemir was born in Canakkale in 1982. Ozdemir graduated B.Sc. and M.Sc. in Chemical Eng. Department at Yildiz Technical University, Istanbul. She has been research assistant since 2006.

Her research interest is in the area of waste management, material characterization, chemical technologies and food technologies especially in teas

and coffees.



Funda Demir was born in İstanbul in 1990. She is continuing her B.Sc studies in Chemical Eng. Department at Yildiz Technical University, Istanbul.



Merve Karakoç was born in İstanbul in 1991. She is continuing her B.Sc. studies in Chemical Eng. Department at Yildiz Technical University, Istanbul.



Sabriye Piskin graduated from Istanbul Technical University on Chemical Engineering with M.Sc. degree in 1974. She completed a Ph.D. degree at the same department in 1983. Her research interests include boron minerals and compounds, hydrogen storage technologies, fuel cell applications, materials characterization, coal, waste management, corrosion, implants and synthetic materials production.