

A Review On 1,2,4-Triazoles

Kishor Raval*, Kaushik Patel, Shivani Patel, Ronak Patel, Snehal Patel

APMC College of Pharmaceutical Education and Research, Himmatnagar-383001, Gujarat, India.

Received on: 22-08-2012; Revised on: 23-08-2012; Accepted on: 29-08-2012


ABSTRACT

Triazole is a five membered heterocyclic system consisting of two carbon atoms and three nitrogen atoms shows wide range of biological activities. Triazoles can be synthesized using Einhorn-Brunner reaction or the Pellizzari reaction from acyl hydrazides and various different method. Triazole derivatives are showing very promising and excellent therapeutic effectiveness. The major activities exhibited by these derivatives include insecticidal, antifungal, antiviral, antibacterial, sedative, hypnotic, anticonvulsant and anti-inflammatory action. In recent year heterocyclic compounds analogues and derivatives have attracted strong interest due to their useful biological and pharmacological properties. The small and simple triazole nucleus is present in compounds involved in research aimed at evaluating new products that possess biological activities, such as, anti-microbial, anti-tumor, anthelmintic, anti-leishmanial, anti-convulsant and anti-inflammatory.

Key words: Triazole, 1H-1, 2,4-Triazole, Heterocycle, Total synthesis, Biological activities.

INTRODUCTION

Triazoles are heterocyclic compounds featuring five member ring of two carbon atoms and three nitrogen atoms as part of the aromatic five-member ring. Triazole refers either one of a pair of isomeric chemical compounds with molecular formula C₂H₃N₃.



During the last few decades, a considerable attention has been devoted to synthesis of 1,2,4 triazole derivatives due to their wide spectrum of biological activities such as anticonvulsant [1], antidepressant [2], antibacterial [3], antifungal [4], anti-inflammatory [5], analgesic [6], anticancer [7] and anti viral [8] activities. The anticonvulsant drug design is based on the presumption that for the activity in maximal electroshock seizure (MES) evaluation, at least one phenyl or similar aromatic group in close proximity to a two electron donor atom is required. For activity in pentylenetetrazole (PTZ) evaluation, an alkyl or amine group close to a two electron donor atom is needed [9]. In view of the above findings, it was thought worthwhile to synthesize a series of ten compounds containing the features of 1,2,4-triazole moiety with different substituents and studied their anticonvulsant activity by electroshock and chemo shock methods. The chemistry of 1,2,4-triazoles described briefly in the literature [9, 10]. The first 1,2,4-triazole derivatives have been synthesised by Bladin in 1885. 1,2,4-triazole derivatives are also useful as analytical reagents [11], photographic chemicals [12] and in polymer synthesis [13].

Synthetic Aspect:

1. Base-catalyzed, direct synthesis of 3,5-disubstituted 1,2,4-triazoles:


A convenient and efficient one step, base-catalyzed synthesis of 3,5-disubstituted 1,2,4-triazoles by the condensation of a nitrile and a hydrazide is presented. A diverse range of functionality and heterocycles are tolerated under the reaction conditions developed, and the reactivity of the nitrile partner is relatively insensitive to electronic effects [14].

*Corresponding author:


Kishor Raval

APMC College of Pharmaceutical Education and Research,
Himmatnagar-383001, Gujarat, India.


*E-Mail: rx.kish148@gmail.com




2. Reid and Heindel^[15] have reported the reaction of aryl acid hydrazide with CS₂/KOH and Hydrazine hydrate furnished triazoles:



Bahittin Kahveci^[16] have reported that the iminoester hydrochlorides are converted into ester ethoxycarbonyl hydrazones, which turn into a new series of substituted 4-amino-4,5-dihydro-1H-1,2,4-triazole-5-ones.




4. Khosrow Zamani and co-worker^[17] have prepared triazole derivatives from respective substituted pyridine carboxylic acid hydrazides and 4-methylphenyl isothiocyanate.




5. Li-Li Liu and Guang Yang^[18] were described a facile method to prepare 1,2,4-triazole by the reaction of isonicotinonitrile with excess amount of hydrazine.

6. Shuki Araki and co-worker^[19] have synthesized 1,2,4-triazoles from 2,3-diphenyltetrazolium salt. using triethylamine as a base.




7. The synthesis of dendrons bearing 1,2,4-triazoles moiety is described by Enrique Diez-Barra et al. [20] The synthesis of these compounds was carried out using different approaches.



8. Kee-Jung lee et al. [21] have reported 1,2,4-triazole from the electrocyclic reaction of conjugated heterocumulenes.

8. Chande et al. [22] have reported 4-anilino-5-mercapto-5-triazoles from α -acyl dithiocarbamate and phenyl hydrazine at 140-160°C.

9. Shin-ichi Nagai [23] have synthesised triazole by the reaction of thiosemicarbazide with formic acid in the presence of acetic anhydride (AC₂O).



10. Yan Shiquang [24] have prepared triazole by the treatment of ferrocenecarboxylic acid hydrazide with aryl isothiocyanate and cyclization of the product gave 3,4-disubstituted 4H-1,2,4-triazole-5-thiol.


11. B. J. Rai and Co-workers [25] have synthesised 3-amino-5-aryl-2-phenyl-2H-1,2,4-triazole by reaction of aroyl-cyanide with phenyl hydrazine.

Therapeutic Importance:


1,2,4-triazole derivatives have been associated with diverse biological activities. Therapeutic activity of 1,2,4-triazoles are listed as under.

1. Antiinflammatory [26]
2. Diuretic [27]
3. Antiviral [28]
4. Antihypertensive [29]
5. Anthelmintics [30]
6. Bactericidal [31]
7. Anticonvulsant [32]
8. Herbicidal [33]
9. Insecticidal & Acaricidal [34]
10. Fungicidal [35]
11. Antimicrobial [36]
12. Anticancer and anti-HIV [37]
13. Plant growth regulator [38]
14. Antileishmania [39]
15. Antitumor [40]
16. Antidepressant and Anxiolytic [41]

A new series of clubbed thiazolidinone-triazole derivatives(I) were synthesized by Mahendra R. Shiradkar et al. [42] The compounds were evaluated for their anticonvulsant activity in two animal models of seizures.





Rossi Carla [43] was investigated 1,2,4-triazoles having antigestative immunosuppressant and antitumor activity. Michael et al. [44] have reported triazoles as potential antibacterial agents. H. Mikali et al. [45] have documented triazoles as antimicrobial agents. Substituted 1,2,4-triazole have been reported for their pharmacological activity by Bahittin Kahveci et al. [46] and antifungal activity [47]. Liu. Chanjian et al. [48] have investigated triazoles (II) as IMPDH inhibitors.





(II)

A series of 4-amino-5-(substituted phenyl) [1,2,4]triazoles(III) and (IV) were synthesized by Loredana Salerno et al. [49] with the aim of obtaining new selective 5-HT_{1A} ligands. All New compounds were tested in radioligand binding experiments, from many of them showed a preferential affinity for the 5-HT_{1A} receptor.




(III) & (IV)


Mari Makoto et al. [50] have prepared water soluble triazoles as fungicides. Laddawahetty et al [51], have synthesised triazole as selective human GABA receptor for the treatment of anxiety and enhancing cognition. Eight novel compounds like (V) were synthesized and their insecticidal activities were tested by Bing Chai and co-workers [52]. The newly designed substituted triazole derivatives (VI) were synthesized by Maria Grazia Mamolo et al. [53] and tested for their *in vitro* antifungal and a n t i mycobacterial activity.



New 1,2,4-triazole compounds (VII), containing a *D,L*-methionine moiety were synthesized by Otilia Pintilie et al.^[54]. All compounds exhibited promising antimicrobial activities. A series of 4-amino-3-(2-furyl)-5-mercapto-1,2,4-triazole (VIII) were prepared by Jingde Wu and co worker^[55]. All the synthesized substituted triazole derivatives were reported as anti-HIV-1 agents by examined their inhibition activity of HIV-1-induced cytopathogenicity in MT-4 cells and by determined their inhibitory effect on HIV-1 reverse transcriptase.




(VII)




(VIII)

A variety of fluconazole derivatives were synthesized by Ruchita Ohlan and co workers^[56]. The synthesized compounds (IX) were evaluated for their in vitro antifungal activity against *C. albicans* and *A. niger*. Jian-yu Jin and co workers^[57] have discovered new 1,2,4-triazole derivatives (X), which may possess significant biological activities. Plant growth-regulating activity tests of all compounds showed remarkable effects on the growth of radish and wheat.




(IX)




(X)

John W. Hull Jr. et al.^[58] have investigated a new family of functionalized 2,6-dihaloaryl 1,2,4 triazole (XI) & (XII) insecticides featuring. Ibtelhal A. Al-Juwaier et al.^[59] have been synthesized anhydronucleosides of 1,2,4-Triazoles which showed pronounce biological activities.




(XI)




(XII)

Mevlut Sedar and co worker^[60] have synthesized a new series of triazole(XIII) and the all compounds were screened for their antimicrobial and antifungal activities. Thirteen new triazoles (XV) containing 1,3-dioxolane rings were synthesized by Liang-Zhong Xu et al.^[61] and their results of preliminary biological tests show that all of these compounds possess some fungicidal and plant growth regulant activities.



(XIII)



(XIV)

Giorgia pastorin et al.^[62] have documented 1,2,4-triazoles as adenosine receptor antagonist and also as human A3 and A2B adenosine

receptor. B.Shivarama Holla et al.^[63] have screened 1,2,4 triazoles for anticancer property. Uesaka et al.^[64] documented triazoles as adrenergic $\alpha 2C$ receptor antagonists. Biologically active triazole were reported by P. V. Chepal et al.^[65] Aoki Satosh et al.^[66] have investigated triazoles as potent cyclooxygenase inhibitor.

CONCLUSION

Triazole is a unique template that is associated with several biological activities. This article highlighted research work of many researchers reported in literature for different pharmacological activities on triazole compounds synthesized. This review has presented comprehensive details of triazole analogues, potent compounds reported for particular pharmacological activity and the method or technique involved in evaluation process. More investigations must be carried out to evaluate more activities of triazole for many diseases whose treatment are difficult in the medical sciences.

REFERENCES:

- Kucukguzel I, Guniz-Kucukguzel S, Rollas S, Otuk-Sanis G, Ozdemir O, Bayrak I, Altug T & Stables J P. *Farmaco.*, **2004**; 59(11); 893.
- Kane J M, Dubley M W, Sorensen S M & Miller F P. *J. Med. Chem.*, **1988**; 31; 1253.
- Karabasanagouda T, Adhikari A V & Shetty N S. *Eur. J. Med. Chem.*, **2007**; 42; 521.
- Sztanke K, Tuzimski T, Rzymowska J, Pasternak K & Kandefers-Szerszen M. *Eur. J. Med. Chem.*, **2008**; 43(2); 404.
- Mullican M D, Wilson M W, Cornor D T, Kostlan C R, Schrier D J & Dyer R D. *J. Med. Chem.*, **1993**; 36; 1090.
- Tozkoparan B, Kupeli E, Yesilada E, Isik S, Ozalp M & Ertan M, *Arzneimittelforschung.*, **2005**; 55(9); 533.
- Demirba N, Uurlulu R & Demirba A. *Bioorg. Med. Chem.*, **2002**; 10; 3717.
- E-Essawy A, E-Sayed W A, E-Kafrawy S A, Morshedy A S & Abdel-Rahman A H. *Z Naturforsch [C]*. **2008**; 63(9-10); 667.
- E. Hoggarth; In 'Chemistry of Carbon Comp.' edited by E. H. Rodd., IV A. P. 452-63.
- V. Meyer, and P. Jacopson; 'Hand book of Organic Chemistry', 2, Pt. 3, Soc., **1920**; 1-2: 590.
- M. Busch; *Ber.*, **1905**; 38; 856.
- T. Nakaic, S. Meedu, and T. Kurahashi. *Jpn. Pat.*, **1973**; 73; 89932; *Chem. Abstr.*, 81, 65182 (1974).
- R. J. Colter, and M. Matzner; 'Ring terming Polymerisation, Part-B-1, 'Heterocyclic ring', Academic, New York, **1972**.
- Kap-Sun Yeung, Michelle E. Farkas, *Tetrahedron Letters*, **2005**; 46; 3429-3432.
- Jack. R. Reid and Ned D. Heindel. *J. Heterocyclic Chem.*, **1976**; 13; 925.
- Bahittin Kahveci. *Molecules*, **2005**; 10; 376.
- Khosrow Zamani, Khalil Faghihi, M. Reza Sangi and Javad Zolgharnein. *Turk. J. Chem.*, **2003**; 27; 119.
- Li Liu and Guang Yang. *Molbank*, **2008**; 56; 213.
- Shuki Araki, Satoshi Hirose, Yoshikazu Konishi, Masatoshi Nogura et al., *Beilstein Journal of Organic Chemistry*, **2009**; 5; 8.
- Enrique Diez-Barra, Joaquin C. Garcia-Martinez, Javier Guerra, et al., *ARKIVOC*, **2002**; V; 17.
- Kee-Jung Lee and Dong-Wook Kim. *J. Heterocyclic Chem.*, **1997**; 34; 1301.
- Madhuka S. Chande, Jayesh D. Bhandari and Vishwas R. Joshi. *Indian J. Chem.*, **1993**; 32B; 1218.
- Shin-ichi Nagai, Taiser Wada, and Mastoshi Takamura. *J. Heterocyclic Chem.*, **1998**; 35; 293. Kee-Jung Lee and Dong-Wook Kim. *J. Heterocyclic Chem.*, **1997**; 34; 1301.
- Yan Shiquang, Chen ning and Wa Hui. *He Cheng Huaxue.*, **1996**; 4(3); 220. *Chem. Abstr.*, **1997**; 126; 171680w.
- B. J. Rai, and R. Lakhan; *Tetrahedron Asymmetry*, **1996**; 7(6); 16.
- Bozo Eva, Szilagyi Geza, Janky Judit, et al., *Arch. Pharm. (Weinheim. Ger.)* **1989**; 322; 58; *Chem. Abstr.*, **1990**; 112; 55731a.
- L. G. Chekovskaya, E. G. Khush, G. K. Rogylchenko, et al., *Farm. Zh (kiev)* **1989**; 5; 67; *Chem. Abstr.*, **1990**; 112; 198233a.
- S. Archana Jyoti, S. Sanjay, V. K. Saxena et al., *J. Indian Chem. Soc.*, **1991**; 68; 103.
- D. B. Reigz; *Eur. Pat. EP 508445*, (Cl. A61k 31/44); *Chem. Abstr.*, **1993**; 118; 59716w.
- S. M. Kudari, S. M. Beede and Munera Wahaja. *Asian. J. Chem.*, **1997**; 9; 20; *Chem. Abstr.*, **1997**; 126.
- F. Pourmorad and A. Shafiee; *J. Sci. Islamic Repub. Iran*, **1998**; 9; 30; *Chem. Abstr.*, **1998**; 129; 175593e.
- N. Karali, G. Capan, N. Ergene, A. Gursoy et al., *Sci. Pharm.*, **1997**; 65; 277; *Chem. Abstr.*, **1998**; 128; 167389g.
- Fusaka Takafumi, Ujikawa Osamu, Kajiwaru Takeshi, and Tanaka Yasushi. *PCT Int. Appl. WO 97, 11, 075*; *Chem. Abstr.*, **1997**; 126; 293354p.

34. Shibata Yasushi, Aoyagi Koichiro, Ichikawa Naomi et al., PCT Int. Appl. Wo. 99 02507; Chem. Abstr., **1999**: 130; 95554y.
35. Itoh Katsumi, Kitazaki Tomoyuki, Matsushita Yoshihiro et al., Eur. Pat Appl. EP 884311; Chem. Abstr., **1999**: 130; 66505c.
36. V. A. N. Murthy Sreenivasa, L. V. G. Nagappa, Nargund et al., Indian J. Heterocycl. Chem., **1998**: 8; 23; Chem. Abstr., **1999**: 130; 66448m.
37. Mark D. Abel, Randy T. Hewgill, Katherine J. Malczylc et al., J. Heterocycl. Chem., **1998**: 35; 193.
38. Chu Chang Hu, Sun Xiao Wen, Zhang Zi Yi, Li Zhi Chun et al., Chin Chem. Lett., **1999**: 10; 361.
39. R. S. Verma, Vinita Bajpai and A. Kapil. Indian J. Heterocycl. Chem., **2000**: 10; 117.
40. Michael J. Ellis, Malcolm F. G. Stevens et al., J. Chem. Soci., **2001**: 23; 3180 (Eng.); Chem. Abstr., **2002**: 137; 185433g.
41. G. Turan-Zitouni, Z. A. Kaplancikli, F. S. Kilic et al., Chem. Abstr., **2003**: 138; 271613t.
42. Mahendra R. Shiradkar, Mangesh Ghodake, Kailash G. Bothara et al., ARKIVOC, **2007**: XIV; 58.
43. Rossi Carla. PCT Int. Appl. WO 98 55463; Chem. Abstr., **1999**: 130; 3836w.
44. Michael J. Genin, Debra A. Allwine, and David J. Anderson. J. Med. Chem., **2000**: 43; 953.
45. Mikail H. Gezinci, Arnold R. Martin and Scott G. Franzblary et al., J. Med. Chem., **2001**: 44; 1560.
46. Bahittin Kahveci, Mevlut Serder and A. Aykutlkizler. Indian J. Heterocycl. Chem., **2001**: 10; 201.
47. Chang-Hu Chu, Xiao-Wen Sun, Zi-Yi, Zi-Yi Zhang et al., Indian J. Chem., **2001**: 40B; 854.
48. Liu Chanjian, Murali T. G. Dhar, Henry H. Gu, S. Gregory et al., U.S Pat. 143, 176, (Cl. 544-59; C07D279/12), (**2002**).
49. Loredana Salerno, Mariangela Siracusa, Francesco Guerrera, Giuseppe Romeo, et al., ARKIVOC, **2004**: V; 312.
50. Mari Makoto, Kagoshima, Yohiko, Uchida, Takaya. PCT Int. Appl. WO 02 66465, (**2002**); Chem. Abstr., **2002**: 137; 201316d.
51. Laddawahetty, Tamara, Merchant, Kevinjohn, Sternfeld, Francine et al., Chem. Abstr., **2002**: 137; 201339p.
52. Bing Chai, Xuhong Qian, Song Cao, Haidong Liu, and Gonghua Song, ARKIVOC, **2003**: ii; 141.
53. Maria Grazia Mamolo, Daniele Zampieri, Valeria Falagiani, Luciano Vio, et al., ARKIVOC, **2004**: V; 231.
54. Otilia Pintilie, Lenuta Profire, , Valeriu Sunel, Marcel Popa and Aurel Pui. Molecules, **2007**: 12; 103.
55. Jingde Wu, Xinyong Liu, Xianchao Cheng, Yuan Cao, Defeng Wang et al., Molecules, **2007**: 12; 2003.
56. Ruchita Ohlan, Sucheta Ohlan, Vikramjeet Judge, and Balasubramanian Narasimhan. ARKIVOC, **2007**: xiv; 172.
57. Jian-yu Jin, Li-xue Zhang, An-jiang Zhang, Xin-Xiang Lei and Jiang-Hai Zhu. Molecules, **2007**: 12; 1596.
58. John W. Hull Jr., Duane R. Romer, David E. Podhorez, Mezzie L. Ash et al., Beilstein Journal of Organic Chemistry, **2007**: 3; 23.
59. Ibtehal A. Al-Juwaiser, Elizabeth John, Maher R. Ibrahim, Yehia A. Ibrahim et al., ARKIVOC, **2009**: ix; 224.
60. Mevlut Serdar, Nurhan Gumrukc, Neslihan Demirbas et al., Turk. J. Chem., **2007**: 31; 315.
61. Liang-Zhong Xu, Shu-Sheng Zhang, Shu-Yan Niu, Yong-Qi Qin et al., Molecules, **2004**: 9; 913.
62. Giorgia Pastorin, Tatiana Da Ros, Giampiero Spalluto, Francesea Deflorian et al., J. Med. Chem., **2003**: 46; 4287.
63. B. Shivarama Holla, B. Veerendra, M. K. Shivananda, and Boja Poojary. Eur. J. Med. Chem., **2003**: 38; 759.
64. Uesaka oriak, Imma Hironori, Kashima Hajime and Kurokawa Masako. PCT Int. Appl. WO 03 28732, **2003**.
65. P. V. Chepal, O. I. Pansonko, V. P. Buriak, V. I Galdash et al., Chem. Abstr., **2003**: 138; 137235t.
66. Aoki Satosh, Nagakawa Toshiya, Konish Nobukiyo et al., PCT Int. Appl WO 03 40110, Chem. Abstr., **2003**: 138; 368895z (2003).

Source of support: Nil, Conflict of interest: None Declared