

Table 1 - Capacity of interception of mild acute FPIES by the previous considered criteria and the new criteria

| Age (months) at diagnosis | Food involved | N episodes - symptoms | ICG diagnostic criteria (ref. 1) | Miceli Sopo diagnostic criteria (ref. 2) | Miceli Sopo suspicion criteria (ref. 2) | Vazquez-Ortiz& Infante criteria (ref. 3) | New diagnostic criteria | New suspicion criteria |
|---------------------------|-------------------------|-----------------------|----------------------------------|--|---|--|-------------------------|------------------------|
| 61 | Fish | 3 - S | X | X | X | X | V | X |
| 11 | Chicken's egg | 3 - R | X | X | X | V | V | X |
| 8 | Cow's milk | 3 - S | X | X | V | X | V | X |
| 11 | Cow's milk | 3 - R | X | X | X | V | V | X |
| 72 | Fish | 3 - R | X | X | X | V | V | X |
| 76 | Rice | 4 - S | X | X | V | X | V | X |
| 11 | Chicken's egg | 3 - R + D | X | X | X | V | V | X |
| 18 | Cow's milk | 3 - R + D | X | X | X | V | V | X |
| 6 | Rice and cow's milk | 2 - R | X | X | X | V | X | V |
| 13 | Cow's milk and soy milk | 2 - R | X | X | X | V | X | V |
| 77 | Chicken's egg | 4 - S | X | X | V | X | V | X |
| 36 | Fish | 2 - R | X | X | X | V | X | V |
| 17 | Chicken's egg | 4 - R | X | X | X | V | V | X |
| 38 | Fish | 3 - R | X | X | X | V | V | X |
| 72 | Chicken's egg | 2 - R | X | X | X | V | X | V |

Legend: S = single vomiting; R = repeated vomiting; D = diarrhea; V = identified; X = identified. No patient has ever presented pallor, lethargy, hypotension.