

Machine learning applied to species occupancy probability and interactions: the missing link in biodiversity assessment and modelling of Antarctic plankton distribution

Marco Grillo^{1,2,*}, Stefano Schiaparelli^{2,5}, Tiziana Durazzano⁷, Letterio Guglielmo⁴, Antonia Granata⁶, and Falk Huettmann³

¹ Department of Physical Sciences, Earth and Environment (DSFTA), University of Siena, Siena, Italy;

² Italian National Antarctic Museum (MNA, Section of Genoa), University of Genoa, Viale Benedetto XV No. 5, 16132 Genoa, Italy;

³ EWHALE lab Inst of Arctic Biology, Biology & Wildlife Dept. University of Alaska Fairbanks;

⁴ Integrative Marine Ecology Department, Stazione Zoologica Anton Dohrn, Napoli, Italy;

⁵ Department of Earth, Environmental and Life Sciences (DISTAV), University of Genoa, Corso Europa 26,
16132 Genoa, Italy;

⁶ Department of Biological, Chemical, Pharmaceutical, and Environmental Sciences (ChiBioFarAm), University of Messina, 98100 Messina, Italy

⁷ Faculty of Biosciences, Fisheries and Economics, Department of Arctic and Marine Biology, UiT The Arctic University of Norway, Tromsø, Norway;
tiziana.durazzano@uit.no

* Correspondence m.grillo@student.unisi.it; Tel.: +39-010-3538329

SUPPLEMENTAL MATERIAL - CORRLOT




Figure 1. Correlations between RIO values obtained from phytoplankton classes and copepod species.