

GEOLOGY AND PALEONTOLOGY OF A NEW UPPER CRETACEOUS (MAASTRICHTIAN) PALEOGENE TRANSITION SITE (K-Pg) DISCOVERED NEAR MULLICA HILL, GLOUCESTER COUNTY, NEW JERSEY

Les Spears¹ and William Kuehne²

¹Amateur paleontologist, Mullica Hill, New Jersey

²The Delaware Valley Paleontological Society, Inc., P.O. Box 686, Plymouth Meeting, Pennsylvania 19462-0686, wkuehne@comcast.net

<https://doi.org/10.5281/zenodo.11238691>

Abstract - The Turtle Creek tributary of Raccoon Creek near Mullica Hill, Gloucester County, New Jersey, exposes the contact between the Upper Cretaceous (Maastrichtian) Navesink Formation and the overlying Paleogene Hornerstown Formation, a nominal K-Pg transition site. The stratigraphy at Turtle Creek is disturbed. In some sections the Kirkwood makes an appearance, but the expected intermediate Vincentown Formation is entirely missing. Although the site is near the historic prolific Mullica Hill Pond site, fossils native to the same formations exposed at Turtle Creek are relatively sparse. *Cucullaea* sp. steinkerns and fragments that appear to be bone were recovered just above the K-Pg contact at the expected horizon of the Main Fossiliferous Layer (MFL) characteristic of K-Pg sites on the Coastal Plain. The Navesink and Hornerstown Formation exposures yield fossils typically associated with these formations including *Cucullaea* and *Pycnodonte* bivalves as steinkerns, an indurated *Exogyra* and other bivalves, *Peridonella dichotoma* and lithified *Thalassinoides* burrows. Silicified wood fragments found in the stream bed gravel correlate with the Kirkwood Formation which outcrops in a downstream section. All fossils at this site are poorly preserved except the silicified wood and the sponge *Peridonella dichotoma*. Stratigraphy reveals an unexpected layer of gravel with embedded cobbles and boulders with Paleozoic fossils above the Hornerstown and Kirkwood Formations attributed to the Bridgeton Formation several kilometers outside of its mapped bedrock range. Some of the clasts exhibit striations and other evidence of glacial derivation. The site is unique for abundant Paleozoic fossils representing Cnidaria anthozoans: tabulata and rugosa, porifera, crinoidea and brachiopoda taxa traced to the tentative Bridgeton Formation hosting layer. Systematics and photographs for all invertebrate native fossils recovered from the site and representative Class taxa of the Paleozoic fossils are included.

INTRODUCTION

Sites in New Jersey exposing the Upper Cretaceous (Maastrichtian)-Paleogene, K-Pg, contact have long attracted much attention from geologists who believe the transition reveals a great mass extinction event ostensibly caused by an asteroid impact in the Yucatan Peninsula at the end of the Cretaceous Period. Faunal and

lithology changes across this transition at such sites support this extinction hypothesis. Gallagher (1993) provided a rather complete overview of New Jersey sites where the transition is exposed. Gallagher, Parris and Spamer (1986) earlier discussed the paleontology, biostratigraphy and depositional environments of this transition and the role of

glauconite across the New Jersey Coastal Plain. Rowan University is conducting a major investigation of the transition and the MFL which are well exposed at their research site at the former Inversand glauconite quarry in Sewell, New Jersey (Rowan, 2021). The aim of this report is to add Turtle Creek in Mullica Hill to the list of known K-Pg sites in New Jersey and to investigate the exposures there for faunal and lithology changes across the transition which could reflect on the postulated K-Pg extinction event.

Site location and description

East sidewall of the inner ravine of Turtle Creek, a small tributary of Raccoon Creek, near Mullica Hill in Gloucester County, New Jersey, located on the Inner Coastal Plain, typical exposure **Figure 1**, Left, site map, Right, topography including the Mullica Hill Pond site. The inner ravine is 5 meters deep, 50 meters wide and the upper surface elevation 21 meters. A highlighted blue line impressed on the topography traces the relevant section of Turtle Creek. Site details are included in **Table 1**.

| | |
|----------------------|--|
| Site | Turtle Creek, tributary of Raccoon Creek |
| Accessibility | Private |
| Municipality | Harrison Township |
| County | Gloucester |
| Altitude, meters | 21 |
| Coordinates, dd | Latitude: 39.783166° N, Longitude: 75.209007° W |
| Site description | Small underfit stream, unconsolidated bedrock exposures along east bank |
| Surficial geology | Qwcp, Weathered Coastal Plain Formations (Stanford 2000). |
| Surficial alluvium | Decomposition residue of Quaternary and Tertiary age on other sedimentary rocks (Fullerton et al., 2016) |
| Bedrock geology | Formations: Navesink, Lower Hornerstown, Lower Member Kirkwood, Bridgeton |
| Source of fossils | Exposures of formations and loose in stream bed gravel |
| Collecting method | Hand pick in-situ or screen gravel in stream bed |
| Residence of fossils | Spears personal collection |
| Age | Upper Cretaceous (Maastrichtian), base, Paleogene to Holocene, upper |

Table 1. Turtle Creek fossil site details.




Figure 1. Left. Typical stretch of Turtle Creek showing boulders, cobbles and gravel in the stream bed from which most of the fossils were recovered. View is upstream with formation exposures on the right (east) bank. Photograph by author. Right, Location section of USGS topographic map of the Turtle Creek site relative to the more familiar classic Mullica Hill Pond site. Turtle Creek is highlighted. The Rowan University Edelman Fossil Park site (former Inversand) is off the map to the right. ArcGIS map by author.

RESULTS

Geology

On the bedrock geology map, **Figure 2**, Left, the Turtle Creek site when viewed from above is situated where the Paleocene Hornerstown Formation adjoins the Miocene Kirkwood Formations without the sequential intermediate formations, Vincentown, Manasquan and Shark River. As the formations generally dip to the southeast and the underground stratigraphy is disturbed, **Figure 3**, the Upper Cretaceous Navesink Formation intrudes only the central part of the site profile so that it was necessary to construct two columns to completely describe the stratigraphy at Turtle Creek. Stratigraphic Section B includes the Navesink-Hornerstown Formations and Section C the Hornerstown-Kirkwood Formations with the Bridgeton Formation overlying both sections. The surface of the formations is overlain with Qwcp, Weathered coastal plain deposits, thickness at site discontinuous and less than 10 ft. (Thickness of Surficial Deposits in New Jersey, online reference) but the description of Qwcp notably

does not report cobbles or boulder sized clasts which are abundant in the stream bed and in the apparent layer of Bridgeton Formation central to the site stratigraphy, **Figure 3**. The Weathered coastal; plain deposits, Qwcp, is described as:

Exposed sand and clay of Coastal Plain bedrock formations. Includes thin, patchy alluvium and colluvium, and pebbles left from erosion of surficial deposits (Stanford, 2000)

It is enigmatic that the apparent Bridgeton Formation appears in these sections as surficial geology shows only its outliers on higher elevations not closer than two kilometers, **Figure 2**, Right. Apparently, the Bridgeton Formation extends outward from these elevated exposures, under the weathered surface plain deposits. The Turtle Creek watershed extends northeast into one of the exposures. Owens and Minard (1979), however, show Bridgeton and Pensauken Formation blanketing the region, Gallagher, Parris and Spamer (1986) review of the

Inversand site (now Rowan University Edelman Fossil Park) also assert that:

The yellowish-orange gravel that unconformably overlies the Hornerstown greensand is the Pensauken Formation, a Pleistocene outwash deposit containing fossiliferous chert pebbles derived from Paleozoic strata in the Appalachians.

Their article does not, however, provide support for this assertion, and, for purposes of the present report, we will assume that it is the Bridgeton Formation which overlies the Hornerstown Formation at Turtle Creek as in **Figure 3**, based on bedrock geology (Gallagher, Parris and Spamer, 1986).




Figure 2. Geology of the Turtle Creek fossil site. Left, bedrock (Sugarman, P., Castelli, M. and Kopcinski, K., 2021), Right, surficial (Fullerton, D., Bush, C. and Pennell, J., 2016), Blue line is the approximate position of Turtle Creek. ArcGIS maps by author.

Stratigraphy profile

The profile of the stratigraphy at Turtle Creek as inferred is sketched in **Figure 3** illustrating the disturbed strata at the site.




Figure 3. Subsurface geology profile sketched as interpreted along the east bank of the Turtle Creek site, downstream to the right. Sketch by author, MS Word.

Stratigraphy

Sketches and descriptions of the two stratigraphic columns constructed in the positions shown in the site geology profile **Figure**

3 are presented below. Sections include lithology symbols for recovered fossils at approximate levels as found in the column. Section B, **Figure 4** and Section C, **Figure 5**.




Figure 4. Stratigraphic Section B Turtle Creek site. See site profile Figure 3 for location. The white band at the Kns-Tht contact ranges from 10 cm to 46 cm in width. The bones were recovered from the thicker section of this band off to the right. Sketch by author, MS Word. Photograph by author.

Turtle Creek site, description of stratigraphy at Section B, **Figure 4**. This section is representative of a part of the profile which is only about three meters wide but contains the only exposure of the Navesink along this part of the ravine. Reading from the stream bed upward in section.

0 Stream bed gravel. Fossil sponges cf. *Peridonella dichotoma*, Indurated mollusks, *Exogyra* and other unidentified bivalves, lithified *Thalassinoides* burrow fillings, oolites, and free form Paleozoic fossils.

0 to 0.46 m Navesink formation: Dark greenish black glauconitic sand with clay content. In the top of this section a few *Cucullaea* sp. steinkerns reside. At the 0.46 m level there is a distinct color change in the glauconitic sand from the Navesink to the Hornerstown. Also, at this contact there

appears to be a thin concretion layer cemented with iron oxides.

0.46 to 1.50 m Lower Hornerstown, the basal part of this section appears to be a very complex. In some parts the glauconitic sand appears to be a lighter color, almost white and partly cemented possibly by the clay content. There is also a mix of glauconite grains that range from bluish-black to dark green. The following fossils came from the lighter glauconitic sand that appears to be cemented by the clay content, a few *Cucullaea* sp. steinkerns and some bone fragments which may trace to the MFL. The contact with underlying Navesink Formation appears gradational marked by a distinct color change, a white band and a quartz slab which may be a cemented part of the band of medium to coarse-grained quartz sand which is known to

occur at the Kns-Tht contact. In the upper part of this section the glauconitic sand, appears to be almost pure glauconite, dusky green in color. At about midway in this section some *Pycnodonte steinkerns*.

1.5 to 2 m This appears to be the Miocene Bridgeton formation gravel resting on the Hornerstown Formation at this site. 0.5 m layer of gravel, with fine clay, a few cobble size rocks of quartzite, schist, bits of gneiss and a least one boulder size sandstone rock. At some areas of the contact with the Hornerstown there is a thin concretion layer with iron oxides.

2 to 3.35 m Surface. Layer of brown sand with scattered pebbles up to six centimeters with topmost layer of topsoil

The Turtle Creek site exposes what appears to be the contact of the Lower Hornerstown Formation with the Navesink Formation in Section B, but near the base of the Lower Hornerstown Formation the expected MFL which is reported to stretch across the state (Rowan 2021, online reference) contains only *Cucullaea* sp. steinkerns and a few bone fragments. The ichnofossil, *Thalassinoides*, has been used to place the MFL at K-Pg exposure sites (Weist et al., 2016). Gallagher, Parris and Spamer, 1986) reported burrows hanging down from the contact but, unfortunately, no lithified burrows or tunnel traces of this taxon have been observed in-situ across this boundary at the Turtle Creek site. Also, the rich faunal content of the Navesink Formation exhibited at the Mullica Hill Pond site (Gallagher, 1993) and at the Rowan Inversand site (Rowan, 2021, online reference) 64 invertebrate taxa recovered from Mullica Hill Pond (Gallagher 1993), six present at Turtle Creek (this report). Although the Vincentown

Formation is the next Formation above in the Paleogene stratigraphic sequence above the Hornerstown, it is missing in the Turtle Creek section probably eroded away and replaced by the Miocene Bridgeton Formation. Neither is the diagnostic brachiopod *Oleneothyris harlani* (Morton) present in the stream bed, another indication that the Vincentown Formation was never present in the Turtle Creek upstream watershed. Nor is there any evidence of this taxon at the nearby Mullica Hill Pond site. In Section C, **Figure 5** below, the Kirkwood Formation overlies the Upper Hornerstown Formation without the Vincentown formation. No silicified wood was found in-situ in the Kirkwood, but some fragments were recovered from the stream bed gravel providing evidence of its presence somewhere along the upstream reach of Turtle Creek.

Note that Cross Section B of the Pitman West bedrock geological map shows the Cohansey Formation above the Kirkwood at higher elevations in the vicinity of Raccoon Creek and does not include the Bridgeton Formation (Sugarman, P., Castelli, M. and Kopcznski, K., 2021). The description of the Cohansey does not include any clasts larger than gravel, maximum size 6 centimeters (Pebble gravel, Udden-Wentworth scale) except ironstone which does not appear at Turtle Creek (Cohansey Formation, online reference). As Cohansey lithology does not differ markedly from the surficial weathered coastal plain deposits, Qwcp, it is conceivable that some Cohansey may be mixed in with the surface deposits. The Cohansey Formation is also a source of Paleozoic fossils (Kuehne, Hajzer and Spears, 2022).




Figure 5. Stratigraphic Section C, showing the layer of Kirkwood Formation, Turtle Creek site. See site profile Figure 3 for location of this section. Sketch by author, MS Word.

Description of stratigraphy at Section C, **Figure 5**.
Reading upward from stream bed

0 – 0.6 m Hornerstown Formation, glauconitic sand, dusky green, almost pure glauconite

0.6 to 0.9 m Kirkwood Formation, mix of light gray sand and clay. Note, In this area of Mullica Hill, the Kirkwood is deeply weathered in the outcrop to shades of orange, dark yellowish orange, grayish orange and light gray.

0.9 to 2.4 m Bridgeton Formation, red gravel with small pebbles throughout with some small cobbles.

0.3 m Layer of topsoil at top

The paucity of fossils in the Navesink at Turtle Creek compared to Rowan Inversand and Mullica

Hill Pond is probably because fossils are more abundant lower in the Navesink Formation, clear from stratigraphic columns at these sites and from Sugarman, Castelli and Kopcznski (2021) description of the Navesink Formation.

White and glauconitic oolites are present on some indurated molluscan slabs recovered from the stream bed at the Turtle Creek site at Section B **Figure 4**, white form pictured in **Figure 6**. Ooids form in shallow seas in warm, wave agitated water. The basal Hornerstown Formation is characterized by transgressive and regressive sequences which may be favorable to oolite formation (Hornerstown Formation, online reference).




Figure 6. Indurated Mollusca slab from stream bed at Section B, Figure 4 with interior oolites. Photographs by author.

Paleontology

The assemblage of fossils found at the Turtle Creek site is presented in **Table 2** below and systematics, descriptions and photographs in the Appendix, complete for native fossils except bones but only representative Class assignments for the exotic Paleozoic fossils. Except for *Peridonella dichotoma* and silicified wood, all fossils are poorly preserved. Three vertebrate fossils which appear to be bone were recovered from the expected location of the MFL in the lower Hornerstown Formation, symbolized on Section B, **Figure 4**. These vertebrates are highly pyritized, encrusted and poorly preserved. Because of their poor quality, photographs of

these vertebrates are not included in the Appendix.

Salisbury and Knapp (1917) speculate that that the chert containing the Paleozoic fossils is in the Miocene Bridgeton Formation and that the fossils there are Devonian, consistent with conclusions at Turtle Creek. The source of Paleozoic fossils on the New Jersey Atlantic Coastal Plain is discussed in Kuehne, Hajzer and Spears (2022). It is remarkable that these Paleozoic fossils, weathered out of probably Devonian diagenic limestone at least 150 kilometers north, have survived with minimal attrition and weathering, another indication of possible ice rafting transport.

| <u>Formation</u> | <u>Taxon</u> |
|--|--|
| Weathered coastal plain deposits, Qwcp. | No in-situ fossils found. |
| Bridgeton | Representative Classes of Paleozoic fossils in stream bed, two Rugosa fossils recover in-situ |
| | Tabulata |
| | Rugosa |
| | Brachiopoda |
| | Bryozoa |
| | Crinoidea |
| Kirkwood | Plantae: silicified wood in stream bed, none in-situ |
| Hornerstown | <i>Cucullaea</i> sp. steinkerns |
| | Bone fragments |
| | <i>Pycnodonte dissimilaris</i> steinkerns |
| Navesink | <i>Cucullaea</i> sp. (steinkern) |
| Stream bed gravel | <i>Peridonella dichotoma</i> Gabb |
| | <i>Thalassinoides</i> lithified burrows |
| Indurated Mollusca slab in stream bed | Exogyra mold cf. <i>Exogyra costata</i> |
| | Bivalves unidentified |

Table 2. Fossil assemblage Turtle Creek site

BOULDERS

Clasts up to cobble and even boulder size litter the bed of Turtle Creek at the site location. Striations, facets and chatter marks mark some of these clasts as glacially derived indicating Pleistocene age but enigmatically appear to reside in the Miocene Bridgeton Formation with many downwashed into the stream bed. Only the surficial weathered coastal plain deposits (Qwcp) are reported to contain materials of Pleistocene age but no glacial derivatives have

been noted in the Turtle Creek exposures of Qwcp, Figure 1. Apparent striations appear on one cobble recovered from the stream bed **Figure 8**. The surface above the bedrock formations is overlain with Qwcp, Weathered coastal plain deposits, thickness at site discontinuous and less than 10 ft. (Thickness of Surficial Deposits in New Jersey, online reference) but notably does not contain cobbles or boulder sized clasts which are plentiful in the stream bed and in the apparent layer of

Bridgeton Formation central to the site stratigraphy profile, **Figure 3**.

An assortment of stream bed clasts is pictured in **Figure 7**. Many lithologies are represented consistent with the description of Newell et al. (2000) including red shale, schist (cf. Manhattan Schist) and gneiss forms representing bedrock found at least 150 kilometers north in the Newark Group and Ridge and Valley provinces of northern New Jersey or farther into New York State. From Sternberg's law of exponential downstream attrition Chatanantavet, et al., (2010), however, it is unlikely that these irregular clasts have transgressed any significant distance (>100 kilometers) by fluvial transport, suggesting

ice rafting along the Delaware River glacial outwash channel. (Kuehne, 2021, online reference). The Bridgeton Formation was deposited by east flowing wash across the coastal plain opposite to the present direction of flow of the Mullica River and its tributaries (Owens and Minard, 1979). Additionally, in fluvial transport, the degree of roundness increases by the logarithm of the distance traveled, so that over a reach of 100 kilometers, only rounds and half-rounds (broken rounds) are expected to survive. None of the large clasts in the bed of Turtle Creek are rounded in this fashion **Figure 7**. Apparent glacial striations on one clast are illustrated in **Figure 8**.



Figure 7. Cobble sized clasts from Turtle Creek stream Bed pictured in Figure 1. Scale bar is 12 inches. Photographs by author.




Figure 8. Large cobble from the Turtle Creek site, 10 x 18 centimeters. Appears to have striations on the right side, adjoining inset. Photograph by author.

DISCUSSION

Although Turtle creek is not a spectacular fossil site when compared to the prolific vertebrate and invertebrate discoveries at Mullica Hill Pond and former Inversand site (Rowan, 2021, online reference), the findings at the Turtle Creek site do add to the body of scientific knowledge of the paleontology of New Jersey particularly about the K-Pg transition and the associated extinction event. The site is unusual, however, in many respects including:

1. Unexpected and unexplained sparseness of native fossils in formations exposed at Turtle Creek correlative with Mullica Hill Pond and Inversand (Rowan, 2021).
2. Only a few bone fragments and *Cucullaea* sp. steinkerns were found at the expected location of the usually fossil rich MFL, an exception to the perceived ubiquitous nature of the MFL across the New Jersey Coastal Plain (Gallagher, 1993).
3. Discovery of Paleozoic fossils in-situ in the tentatively identified Bridgeton Formation
4. Large number of Paleozoic fossils lodged in stream bed gravel of Turtle Creek.

5. Incongruous outlier of the Bridgeton Formation in the Turtle Creek site stratigraphy
6. The presence of apparent glacially derived clasts in the stream bed

DATA REPOSITORY

The complete assemblage of Paleozoic fossils recovered from Turtle Creek is included in the data repository at DOI: 10.13140/RG.2.2.10504.70405 Resolve doi name at <https://www.doi.org/>

REFERENCES

Published references.

- Chatanantavet, P., Lajeunesse, E., Parker, G., Malverti, G. and Meunier, P., 2010, Physically based model of downstream fining in bedrock streams with lateral input. *Water Resour.* 46, W02518. doi:10.1029/2008WR007208.
- Fullerton, D., Bush, C. and Pennell, J., 2016, Map of Surficial Deposits and Materials in the Eastern and Central United States (East of 102° West Longitude). U.S. Geological Survey, Geologic Investigations Series I-2789, U. S. Geological Survey, Reston, Virginia.

Gallagher, W., 1993, The Cretaceous/Tertiary mass extinction event in the northern Atlantic Coastal Plain. *The Mosasaur*, Vol. 5, 75-154.

Gallagher, W., Parris, D. and Spamer, E., 1986, The Paleontology, biostratigraphy and depositional environments of the Cretaceous-Tertiary transition on the New Jersey Coastal Plain. *The Mosasaur*, Vol. 3, 1-35.

Kuehne, W, Hajzer, F. and Spears, L., 2022, Provenance of Silicified Paleozoic Invertebrate Fossils Dispersed in Neogene - Quaternary Age Sediments across the New Jersey Atlantic Coastal Plain. *The Mosasaur*, Vol. 12, 7-28.

Newell, W., Powars, D., Owens, J., Stanford, S. and Stone, B., 2000, Surficial Geologic Map of Central and Southern New Jersey, MAP I-2540-D and Text to accompany MAP I-2540-D. U.S. Geological Survey, Reston, Virginia.

Owens, J. and Minard, J., 1979, Upper Cenozoic Sediments of the Lower Delaware Valley and the Northern Delmarva Peninsula, New Jersey, Pennsylvania, Delaware, and Maryland, Professional Paper 1067-D; U.S. Geological Survey, Reston, VA. doi:10.3133/pp1067D.

Salisbury, R. and Knapp, G., 1917, The Quaternary Formations of Southern New Jersey. in Final Report Series of the State Geologist, Vol. VIII; New Jersey Geological and Water Survey, Trenton, New Jersey.

Stanford, S., 2000, Surficial Geology of the Long Branch Quadrangle Monmouth

County, New Jersey, Open File Map OFM 38. U.S. Geological Survey, Reston, Virginia.

Sugarman, P., Castelli, M. and Kopcinski, K., 2021, Bedrock Geologic Map of The Pitman West Quadrangle Gloucester and Salem Counties, New Jersey. Open File Map OFM 139. New Jersey Geological and Water Survey, Trenton, New Jersey.

Wiest, L., Buynevich, I., Grandstaff, D., Terry, D., Maza, Z. and Lacovara, K., 2016, Ichnological Evidence for Endobenthic Response to the K–Pg Event, New Jersey, U.S.A. *Palaios*, 31(5) 231-241. URL: <http://www.bioone.org/doi/full/10.2110/palo.2015.080>

Online references

Cohansey Formation. Online reference available at: <https://mrdata.usgs.gov/geology/state/sgmc-unit.php?unit=NJTch%3B1> U. S. Geological Survey, Mineral Resources, Reston, Virginia. Accessed 10 January 2023.

Hornerstown Formation Online reference available at: <https://mrdata.usgs.gov/geology/state/sgmc-unit.php?unit=NJTht%3B1> U. S. Geological Survey, Mineral Resources, Reston, Virginia. Accessed 21 January 2023.

Kuehne, W., 2021, The Provenance of Exotic Boulders Embedded in Quaternary Sediments on the Atlantic Coastal Plain within the Relict Pleistocene Periglacial Region of Southwestern New Jersey, ResearchGate. Online reference available at: <https://www.researchgate.net/> DOI: [10.13140/RG.2.2.26664.65281](https://doi.org/10.13140/RG.2.2.26664.65281).

Lower Member of the Kirkwood Formation.
Online reference available at
<https://mrdata.usgs.gov/geology/state/sgmc-unit.php?unit=NJTkI%3B1> U. S.
Geological Survey, Mineral Resources,
Reston, Virginia. Accessed 30 December
2022.

Oolites. Online reference available at:
<https://www.sandatlas.org/oolite/> Sand
Atlas. Accessed 21 January 2023.

Rowan researchers deploy 3D mapping tools
to define fossil layer, 2021, Online reference
available at:
<https://today.rowan.edu/news/2021/11/rowan-researchers-deploy-3d-mapping-tools-to-define-fossil-layer.html> Accessed 17
January 23.

Thickness of Surficial Deposits in New Jersey,
2013. Online reference available at
<http://www.state.nj.us/dep/njgs/geodata/index.htm> New Jersey Department of
Environmental Protection, New Jersey
Geological and Water Survey. Accessed 9
January 2023.

APPENDIX

Paleontology systematics and description

Plate reference key

| | | |
|----|----|----|
| 1 | 2 | 3 |
| 4 | 5 | 6 |
| 7 | 8 | 9 |
| 10 | 11 | 12 |

Plate I, Native fossils from Turtle Creek

1. Porifera. *Peridonella dichotoma* Gabb. Individuals collected from stream bed.
2. Porifera. *Peridonella dichotoma* collected from stream bed.
3. Crustacea: Ichnofossils, *Thalassinoides*. Lithified burrows recovered from stream bed. Outside diameter ranges from 1.3 to 2 cm. minimal branching, Light gray exterior indicates Navesink Formation host, reddish sediment filling suggests Hornerstown Formation.
4. Crustacea: Ichnofossils, *Thalassinoides* lithified burrow end view, reddish sediment filling suggests oxidized glauconite filling (Hornerstown Formation, online reference).
5. Crustacea: ichnofossils, *Thalassinoides* lithified burrow end view green filling, probable pure glauconite from the Hornerstown Formation (Hornerstown, online reference).
6. Mollusca, Bivalvia: *Pycnodonte dissimilaris* poorly preserved pyritized steinkerns.
7. Mollusca, Bivalvia: *Exogyra*. cf. *Exogyra costata*. traced outline on indurated slab of mollusks.
8. Indurated bivalves. Opposite side of No. 7.
9. Mineral, Oolites on No. 8, enlarged view.
10. Mollusca, Bivalvia: *Cucullaea* sp. cf. *vulgaris*, poorly preserved steinkern with some thin patches of shell material. Found in-situ in the basal Hornerstown Formation. From on-line research, *Cucullaea* fossils are often only internal molds.
11. Mollusca, Bivalvia. *Cucullaea* sp. from Hornerstown Formation, poorly preserved steinkern. Opposite side view of No. 10.
12. Plantae: silicified wood fragments from stream bed.

Plate I - Native fossils of Turtle Creek.



Plate 2 Paleozoic fossil Classes with representative specimens. References [n] follow plate.

Plate reference key

| | | |
|----|----|----|
| 1 | 2 | 3 |
| 4 | 5 | 6 |
| 7 | 8 | 9 |
| 10 | 11 | 12 |
| 13 | 14 | 15 |

Brachiopoda

1. MH 101. Brachiopoda, cf. *Leptaena rhomboidalis*, 24 mm wide, buff, Ordovician (Llanvirn) Devonian (Pragian)[2], [1](p. 38, pl. 132, figs. 16-20), found in Devonian of New York and Pennsylvania
2. 2nd view, *Leptaena rhomboidalis*
3. MH 172. Brachiopod, spiriferid, 10 mm, tan, external mold on poriferid fossil.

Porifera

4. MH 110. Porifera, Demospongea, cf. *Hindia* sp., 32 mm wide, white, spherical, with narrow canals that radiate from the center of the sphere, found in New York, New Jersey, Ordovician through Permian, [5].
5. MH 112. Porifera, Stromatoporoid, cf. *Labechia* sp., 45 mm long, buff. Silurian, Earlton, Ontario [6].
6. MH 113. Porifera, Schlerosponges, cf. *Chaetetes* sp., length 50 mm, width 22 mm, buff. colony of long slender tubes that radiate from center, corallites polygonal to rounded, septa absent in calices, widespread in North America, Middle Devonian through Permian [5].

Rugosa

7. MH 107. Rugosa, cf. *Heliophyllum canadense*, 31 mm long, buff, horn

shape, calice 23 mm wide, Devonian, Onondaga limestone of Mendon, Ontario [3].

8. MH 103. Rugosa, cf. *Heliophyllum* sp., 10 mm long, white. small solitary, conical, calice 15 mm wide with septa, widespread in eastern North America, Lower and Middle Devonian [5].
9. MH 108. Rugosa, cf. *Hexagonia* sp., colonial, irregular shape, buff, corallites averaging 2 to 3 mm in diameter, some with faint septa, found in the Devonian Limestone of Genesee County, New York [1].

Tabulata

10. MH 147. Tabulata, favositid, *Favosites turbinatus*, 25 mm, buff.
11. MH 166. Tabulata, cf. *Favosites* sp., corallites 3 mm wide, white, widespread in North America, Upper Ordovician through Middle Devonian [5].
12. MH 148. Tabulata, favositid, cf. *Favosites* sp., 20 mm wide, white. oval, corallites 1 mm wide, polygonal, white, widespread in North America, Upper Ordovician through Middle Devonian [5].

Miscellaneous

13. MH 177. Crinoidea in chert matrix on MH 174.
14. MH 170. Porifera, Archaeocyathid, cf. *Archaeocyatha* sp., partial specimen in chert clast, buff, found in the Cambrian [3].
15. MH 111. Tabulata, favositid, cf. *Favosites* sp., 48 mm wide, buff, mound shape with bottom center stalk, tightly packed corallites, less than 1 mm wide, widespread in North America, Upper Ordovician through Middle Devonian [5].

Plate 2. Paleozoic fossils Turtle Creek site

SYSTEMATICS REFERENCES FOR PALEOZOIC FOSSILS

1. Shimer, H. W. and Shrock, R., 1944, Index Fossils of North America. The M. I. T. Press, Cambridge, Massachusetts.
2. Digital Atlas of Ordovician Life. Available online at: <http://www.ordovicianatlas.org/atlas/brachiopoda/strophomenata/strophomenida/rafinesquinidae/leptaena/>

3. Digital Atlas of Ancient Life. Morphology. Available online at: <https://www.digitalatlasofancientlife.org/learn/cnidaria/anthozoa/rugosa/>
5. Thompson, 1994, National Audubon Society Field Guide to North American Fossils. National Audubon Society.
6. Fossilarium. Available online at: <https://www.fossiles.ca/en/learn/local-fossil-classification/stromatoporoids-rough-sponges/>