

Preoperative Atelectasis

Part 7: Posthoc Analyses

Javier Mancilla Galindo

2024-04-17

Table of contents

Rationale	2
Setup	3
Outcome variable	5
Prevalence of atelectasis	5
Atelectasis - obesity class	6
Prevalence Ratio	14
Table 2 appendage	14
Ordinal Logistic Regression Model	15
Check proportional odds assumption for main variable of interest:	15
Univariate models for covariates:	18
Multivariable model	22
Package References	24

Rationale

We observed that SpO2 starts decreasing at BMIs above 40-45. Thus, by having used the WHO obesity class categories, detail on differences above BMI 40 for the extent of atelectasis percentage may have been lost. The WHO obesity class categories do not reflect the extent of variation in BMI observed in this sample of patients:

- Class 1, **BMI [30,35)**: ~25% participants
- Class 2, **BMI [35,40)**: ~25% participants
- Class 3, **BMI >40**: ~50% of participants, with a median BMI above a 5 units range.

Thus, creating subcategories within the class 3 obesity may allow to assess the impact of BMI increases above 40 on atelectasis percentage with more detail.

Thus, I will extend the categories of BMI with the following categories:

- **BMI [30,35)** kg/m²
- **BMI [35,40)** kg/m²
- **BMI [40,45)** kg/m²
- **BMI [44,50)** kg/m²
- **BMI >50** kg/m²

Setup

Packages used

```
if (!require("pacman", quietly = TRUE)) {  
  install.packages("pacman")  
}  
  
pacman::p_load(  
  tidyverse, # Used for basic data handling and visualization.  
  table1, #Used to add labels to variables.  
  RColorBrewer, #Color palettes for data visualization.  
  gridExtra, #Used to arrange multiple ggplots in a grid.  
  grid, #Used to arrange multiple ggplots in a grid.  
  mgcv, #Used to model non-linear relationships with a general additive model.  
  ggmosaic, #Used to create mosaic plots.  
  car, #Used assess distribution of continuous variables (stacked Q-Q plots).  
  simpleboot, boot, # Used to calculate mean atelectasis coverage and  
    # 95%CI through bootstrapping.  
  broom, #Used to exponentiate coefficients of regression models.  
  sandwich, #Used to calculate robust standard errors for prevalence ratios.  
  EValue, #Used to calculate E-values as sensitivity analysis.  
  flextable, #Used to export tables.  
  rms, #Used to model ordinal outcome (atelectasis percent) and  
    #test proportional odds assumptions.  
  VGAM, #Used to model partial proportional odds model.  
  gt, #Used to present a summary of the results of regression models.  
  report #Used to cite packages used in this session.  
)
```

Session and package dependencies

R version 4.3.3 (2024-02-29 ucrt)
Platform: x86_64-w64-mingw32/x64 (64-bit)
Running under: Windows 11 x64 (build 22631)

Matrix products: default

locale:

```
[1] LC_COLLATE=Spanish_Mexico.utf8 LC_CTYPE=Spanish_Mexico.utf8
[3] LC_MONETARY=Spanish_Mexico.utf8 LC_NUMERIC=C
[5] LC_TIME=Spanish_Mexico.utf8
```

```
time zone: Europe/Berlin
tzcode source: internal
```

```
attached base packages:
```

```
[1] splines      stats4      grid         stats        graphics    grDevices datasets
[8] utils        methods     base
```

```
other attached packages:
```

```
[1] report_0.5.8      gt_0.10.1         VGAM_1.1-10       rms_6.8-0
[5] Hmisc_5.1-2       flextable_0.9.5   EValue_4.1.3      sandwich_3.1-0
[9] broom_1.0.5       boot_1.3-30       simpleboot_1.1-7  car_3.1-2
[13] carData_3.0-5     ggmosaic_0.3.3    mgcv_1.9-1        nlme_3.1-164
[17] gridExtra_2.3     RColorBrewer_1.1-3 table1_1.4.3       lubridate_1.9.3
[21] forcats_1.0.0     stringr_1.5.1     dplyr_1.1.4       purrr_1.0.2
[25] readr_2.1.5       tidyr_1.3.1       tibble_3.2.1      ggplot2_3.5.0
[29] tidyverse_2.0.0   pacman_0.5.1
```

```
Set seed (for reproducibility of bootstrapping) as the current year 2023:
```

```
seed <- 2023
```

Outcome variable

Corroborate that the new BMI breaks category was created successfully:

```
type_obesity
30-35 35-40 40-45 45-50    50
    63    53    57    31    32
```

Percentages:

```
type_obesity
30-35 35-40 40-45 45-50    50
 26.7  22.5  24.2  13.1  13.6
```

Prevalence of atelectasis

```
           Yes    No
frequencies 77.0 159.0
percent    32.6  67.4
```

Prevalence of atelectasis with 95% confidence interval

```
1-sample proportions test without continuity correction

data:  frequencies, null probability 0.5
X-squared = 28.492, df = 1, p-value = 9.411e-08
alternative hypothesis: true p is not equal to 0.5
95 percent confidence interval:
 0.2696526 0.3884549
sample estimates:
           p
0.3262712
```

The prevalence of atelectasis was **32.6 (95%CI: 26.97, 38.85)**.

Atelectasis - obesity class

Mean expected frequency:

	mean_expected_freq
1	23.6


Frequencies:

	atelectasis	
type_obesity	Yes	No
30-35	8	55
35-40	15	38
40-45	7	50
45-50	15	16
50	32	0

Percentage:

	atelectasis	
type_obesity	Yes	No
30-35	12.70	87.30
35-40	28.30	71.70
40-45	12.28	87.72
45-50	48.39	51.61
50	100.00	0.00

Mosaic Plot



Pearson's Chi-squared test

```
data: frequencies
X-squared = 92.149, df = 4, p-value < 2.2e-16
```

Atelectasis location by obesity class

Mean expected frequency:

```
mean_expected_freq
1                7.7
```

Mean expected frequency is greater than 5.0, so chi-squared without continuity correction is adequate.


Frequencies:

```
atelectasis_location
type_obesity Bilateral lung bases Right lung base predominance
30-35          1                7
35-40          5               10
40-45          4                3
45-50          4               11
50            10               22
```

Percentage:

	atelectasis_location	
type_obesity	Bilateral lung bases	Right lung base predominance
30-35	12.50	87.50
35-40	33.33	66.67
40-45	57.14	42.86
45-50	26.67	73.33
50	31.25	68.75

Mosaic Plot



Pearson's Chi-squared test

data: frequencies
X-squared = 3.6755, df = 4, p-value = 0.4517

Prevalence of atelectasis with 95% confidence intervals calculated with sourced script ***Prevalence_atelectasis.R***

atelectasis	type_obesity	n	prev	confint	Bilateral lung bases	Right lung base predominance
Yes	Total	77	32.63	26.77 - 39.06	24 (31.17%)	53 (68.83%)
No	Total	159	67.37		24 (31.17%)	53 (68.83%)
Yes	30-35	8	12.70	6.03 - 24.04	1 (12.5%)	7 (87.5%)

No	30-35	55	87.30		1 (12.5%)	7 (87.5%)
Yes	35-40	15	28.30	17.2 - 42.56	5 (33.33%)	10 (66.67%)
No	35-40	38	71.70		5 (33.33%)	10 (66.67%)
Yes	40-45	7	12.28	5.49 - 24.29	4 (57.14%)	3 (42.86%)
No	40-45	50	87.72		4 (57.14%)	3 (42.86%)
Yes	45-50	15	48.39	30.56 - 66.6	4 (26.67%)	11 (73.33%)
No	45-50	16	51.61		4 (26.67%)	11 (73.33%)
Yes	50	32	100.00	86.66 - 100	10 (31.25%)	22 (68.75%)

Atelectasis Percent

Mean atelectasis percentage

The following would be the mean atelectasis percentage coverage if a normal distribution were assumed, which is what has been done in some prior studies:

```

      mean      sd
1 2.658898 4.687145

```

And by obesity class:

```

# A tibble: 5 x 3
  type_obesity  mean    sd
  <fct>        <dbl> <dbl>
1 30-35        0.913  2.89
2 35-40        1.56   3.15
3 40-45        0.702  2.05
4 45-50        3.63   4.22
5 50          10.5    5.40

```

As is evident from these numbers, assuming normality causes standard deviation to capture negative values, which is impossible in reality for this variable.

Thus, bootstrapping the mean and 95%CI is expected to lead to more appropriate estimates.

I will calculate this for class 3 subgroups:

Subgroup 1

Mean:

[1] 0.7019737

95% CI:

BOOTSTRAP CONFIDENCE INTERVAL CALCULATIONS
Based on 10000 bootstrap replicates

CALL :
boot.ci(boot.out = boot_sub1)

Intervals :
Level Normal Basic
95% (0.1755, 1.2276) (0.1316, 1.1842)

Level Percentile BCa
95% (0.2193, 1.2719) (0.2193, 1.2719)
Calculations and Intervals on Original Scale

Subgroup 2

Mean:

[1] 3.636355

95% CI:

BOOTSTRAP CONFIDENCE INTERVAL CALCULATIONS
Based on 10000 bootstrap replicates

CALL :
boot.ci(boot.out = boot_sub2)

Intervals :
Level Normal Basic
95% (2.160, 5.084) (2.097, 5.000)

Level Percentile BCa
95% (2.258, 5.161) (2.177, 5.081)
Calculations and Intervals on Original Scale

Subgroup 3

Mean:

```
[1] 10.46127
```

95% CI:

BOOTSTRAP CONFIDENCE INTERVAL CALCULATIONS

Based on 10000 bootstrap replicates

CALL :

```
boot.ci(boot.out = boot_sub3)
```

Intervals :

Level	Normal	Basic
95%	(8.65, 12.31)	(8.52, 12.26)

Level	Percentile	BCa
95%	(8.67, 12.42)	(8.83, 12.50)

Calculations and Intervals on Original Scale

The mean atelectasis percentage coverage in class 3 obesity subcategories was: subgroup 1 (0.7%, 95%CI:0.22-1.27), subgroup 2 (3.64%, 95%CI:2.18-5.08), and subgroup 3 (10.46%, 95%CI:8.83-12.5).

Atelectasis percentage by obesity subgroups

Now, I will continue assessing atelectasis percentage if assumed to be categorical ordinal:

Mean expected frequency:

	mean_expected_freq
1	5.244444

Mean expected frequency is very close to 5.0, so I will use chi-squared with continuity correction.

Frequencies:

	type_obesity				
atelectasis_percent	30-35	35-40	40-45	45-50	50
0	55	38	50	16	0
2.5	2	7	2	0	0
5	1	2	1	5	5
7.5	4	4	4	7	14
10	0	1	0	2	3
12.5	0	0	0	0	1
15	0	1	0	1	2
17.5	1	0	0	0	6
27.5	0	0	0	0	1

Pearson's Chi-squared test

data: frequencies
X-squared = 145.94, df = 32, p-value < 2.2e-16

Figure S4

Figure created with sourced script *FigureS4.R*




Figure 1: Figure S4. Atelectasis percentage on chest CT by obesity categories.

Prevalence Ratio

This [paper](#) and accompanying code were used to calculate prevalence ratios.

A modified Poisson regression model with robust errors will be applied to obtain prevalence ratios.

Prevalence ratios were calculated with the accompanying sourced script ***Prevalence_Ratio_subgroups.R***


Table 2 appendage

Category	PR	SE	95%CI	aPR	aSE	a95%CI	Evalue	Evalue_lower
40-45	0.97	0.48	0.37-2.5	0.85	0.46	0.35-2.1	1.63	NA
45-50	3.81	0.38	1.81-8.01	3.52	0.39	1.63-7.61	6.50	2.64
50	7.87	0.33	4.12-15.05	8.00	0.33	4.22-15.17	15.48	7.91

Ordinal Logistic Regression Model

This modelling strategy was performed according to:

- Harrel, Frank. March, 2022. "Assessing the Proportional Odds Assumption and its Impact". Statistical Thinking. March 9, 2022.



Check proportional odds assumption for main variable of interest:

	Model Likelihood Ratio Test	Discrimination Indexes	Rank Discrim. Indexes
Obs 236	LR ² 110.61	R^2 0.416	0.556
Distinct Y 8	d.f. 4	$R^2_{4,236}$ 0.363	
$Y_{0.5}$ 1	$\Pr(>^2)$ <0.0001	$R^2_{4,163.1}$ 0.480	
max log	Score ² 130.41	$ \Pr(Y$	
$L/$ 3×10^{-7}	$\Pr(>^2)$ <0.0001	median) $^{-1/2}$ 0.303	

		S.E.	Wald Z	$\Pr(> Z)$
y 2.5	-1.9013	0.3782	-5.03	<0.0001
y 5	-2.2844	0.3922	-5.82	<0.0001

		S.E.	Wald Z	Pr(> Z)
y 7.5	-2.8457	0.4148	-6.86	<0.0001
y 10	-4.6674	0.5112	-9.13	<0.0001
y 12.5	-5.1894	0.5434	-9.55	<0.0001
y 15	-5.2932	0.5508	-9.61	<0.0001
y 17.5	-5.7973	0.5934	-9.77	<0.0001
type_obesity=35-40	0.9189	0.4822	1.91	0.0567
type_obesity=40-45	-0.0551	0.5524	-0.10	0.9205
type_obesity=45-50	2.0158	0.5164	3.90	<0.0001
type_obesity= 50	4.4561	0.5591	7.97	<0.0001

Odds ratio for type obesity in an univariable model:

Effects							
Response:							
atelectasis_percent							
	Low	High	Δ	Effect	S.E.	Lower 0.95	Upper 0.95
type_obesity --- 35---40:30---35	1	2		0.91890	0.4822	-0.02626	1.864
<i>Odds Ratio</i>	1	2		2.50600		0.97410	6.449
type_obesity --- 40---45:30---35	1	3		-0.05514	0.5524	-1.13800	1.028
<i>Odds Ratio</i>	1	3		0.94640		0.32050	2.794
type_obesity --- 45---50:30---35	1	4		2.01600	0.5164	1.00400	3.028
<i>Odds Ratio</i>	1	4		7.50700		2.72900	20.650
type_obesity --- 50:30---35	1	5		4.45600	0.5591	3.36000	5.552
<i>Odds Ratio</i>	1	5		86.15000		28.80000	257.800

Proportional odds assumption:

Wald			
Statistics for			
atelectasis_percent			
	²	d.f.	P
type_obesity	85.40	4	<0.0001
TOTAL	85.40	4	<0.0001

This shows that the proportional odds assumption is not met since $p < 0.05$ in the ANOVA test.

Will repeat the process described in Part 5:

0	5	10	15
170	47	7	12

Are subgroups better represented now?

	30-35	35-40	40-45	45-50	50
0	57	45	52	16	0
5	5	6	5	12	19
10	0	1	0	2	4
15	1	1	0	1	9

Some improvement.

Will now test the impact of not meeting the proportional odds assumption in a model adjusted for covariates:

```
impact_PO <- impactPO(
  atelectasis_percent ~ type_obesity + age + sex + altitude_cat,
  nonpo = ~ type_obesity,
  data = data,
  newdata = data_prop,
  relax = "multinomial"
)
```

I was not able to compare against the partial proportional odds (PPO). This can be corroborated by changing `relax = "multinomial"` to `relax = "both"` or `relax = "ppo"` in the above code. This was likely due to a problem in convergence of models with such small subgroups. Previously, I tried comparing models for posthoc analyses in the `VGAM` package and had problems in convergence. Thus, I am presenting the results for the comparison against a multinomial model only:

	PO	Multinomial
Deviance	354.2135	336.7042
d.f.	10	24
AIC	374.2135	384.7042
p	7	21
LR χ^2	109.2016	126.7109
LR - p	102.2016	105.7109
LR χ^2 test for PO		17.50931
d.f.		14
Pr(> χ^2)		0.230051
MCS R2	0.3704304	0.4154487

MCS R2 adj	0.3514770	0.3610492
McFadden R2	0.2356454	0.2734286
McFadden R2 adj	0.2054349	0.1827971
Mean difference from PO		0.02990068

The proportional odds model has a lower AIC and higher adjusted McFadden R2, meaning that the proportional odds model is a more parsimonious model that explains the relationship better than a multinomial model.

Thus, I will proceed to fit ordinal models.

Univariate models for covariates:

	Model Likelihood Ratio Test	Discrimination Indexes	Rank Discrim. Indexes
Obs 236	LR ² 58.42	R^2 0.244	0.539
Distinct Y 8	d.f. 1	$R^2_{1,236}$ 0.216	
$Y_{0.5}$ 1	Pr(> ²) <0.0001	$R^2_{1,163.1}$ 0.297	
max log	Score ² 70.39	Pr(Y	
L/ 2×10^{-5}		median)- ^{1/2} 0.275	
	Pr(> ²) <0.0001		

		S.E.	Wald Z	Pr(> Z)
y 2.5	-1.1960	0.1679	-7.12	<0.0001
y 5	-1.4963	0.1824	-8.20	<0.0001
y 7.5	-1.9260	0.2063	-9.34	<0.0001
y 10	-3.4122	0.3139	-10.87	<0.0001
y 12.5	-3.8627	0.3554	-10.87	<0.0001
y 15	-3.9514	0.3648	-10.83	<0.0001
y 17.5	-4.3894	0.4197	-10.46	<0.0001
sleep_apnea=Yes	2.7008	0.3665	7.37	<0.0001

Effects

Response:

atelectasis_percent

	Low	High	Δ	Effect	S.E.	Lower 0.95	Upper 0.95
sleep_apnea --- Yes:No	1	2		2.701	0.3665	1.982	3.419
<i>Odds Ratio</i>	1	2		14.890		7.260	30.540

Effects									
Response:									
atelectasis_percent									
		Low	High	Δ	Effect	S.E.	Lower 0.95	Upper 0.95	
<hr/>									
<hr/>									
		Model Likelihood			Discrimination		Rank Discrim.		
		Ratio Test			Indexes		Indexes		
Obs	236	LR ²	3.06		R^2	0.014			0.104
Distinct Y	8	d.f.	1		$R^2_{1,236}$	0.009			
$Y_{0.5}$	1	Pr(> ²)	0.0802		$R^2_{1,163.1}$	0.013			
max log		Score ²	2.69		Pr(Y				
L/	0.0001				median)- ¹ / ₂	0.175			
		Pr(> ²)	0.1007						

		S.E.	Wald Z	Pr(> Z)
y 2.5	-0.6588	0.1430	-4.61	<0.0001
y 5	-0.8826	0.1487	-5.94	<0.0001
y 7.5	-1.2022	0.1602	-7.50	<0.0001
y 10	-2.3759	0.2411	-9.85	<0.0001
y 12.5	-2.7833	0.2868	-9.70	<0.0001
y 15	-2.8681	0.2977	-9.63	<0.0001
y 17.5	-3.2923	0.3608	-9.12	<0.0001
asthma=Yes	-1.0153	0.6414	-1.58	0.1135

Effects									
Response:									
atelectasis_percent									
		Low	High	Δ	Effect	S.E.	Lower 0.95	Upper 0.95	
<hr/>									
asthma --- Yes:No		1	2	-1.0150	0.6414		-2.2720	0.2419	
<i>Odds Ratio</i>		1	2	0.3623			0.1031	1.2740	
<hr/>									
		Model Likelihood			Discrimination		Rank Discrim.		
		Ratio Test			Indexes		Indexes		
Obs	236	LR ²	2.08		R^2	0.010			0.096
Distinct Y	8	d.f.	1		$R^2_{1,236}$	0.005			
$Y_{0.5}$	1	Pr(> ²)	0.1490		$R^2_{1,163.1}$	0.007			

	Model Likelihood Ratio Test	Discrimination Indexes	Rank Discrim. Indexes
max log $L/ \mid 3 \times 10^{-7}$	Score ² 2.24 Pr(> ²) 0.1347	Pr(Y median)- ¹ / ₂ 0.173	

		S.E.	Wald Z	Pr(> Z)
y 2.5	-0.7883	0.1470	-5.36	<0.0001
y 5	-1.0111	0.1533	-6.60	<0.0001
y 7.5	-1.3318	0.1655	-8.05	<0.0001
y 10	-2.5139	0.2471	-10.17	<0.0001
y 12.5	-2.9216	0.2921	-10.00	<0.0001
y 15	-3.0059	0.3028	-9.93	<0.0001
y 17.5	-3.4289	0.3651	-9.39	<0.0001
sex=Man	0.6380	0.4315	1.48	0.1392

Effects

Response:

atelectasis_percent

	Low	High	Δ	Effect	S.E.	Lower 0.95	Upper 0.95
sex --- Man:Woman	1	2		0.638	0.4315	-0.2076	1.484
<i>Odds Ratio</i>	1	2		1.893		0.8125	4.409

	Model Likelihood Ratio Test	Discrimination Indexes	Rank Discrim. Indexes
Obs 236	LR ² 0.63	R^2 0.003	0.049
Distinct Y 8	d.f. 1	$R^2_{1,236}$ 0.000	
$Y_{0.5}$ 1	Pr(> ²) 0.4273	$R^2_{1,163.1}$ 0.000	
max log $L/ \mid 0.003$	Score ² 0.63 Pr(> ²) 0.4274	Pr(Y median)- ¹ / ₂ 0.173	

		S.E.	Wald Z	Pr(> Z)
y 2.5	-0.2840	0.5705	-0.50	0.6186
y 5	-0.5054	0.5715	-0.88	0.3765
y 7.5	-0.8240	0.5733	-1.44	0.1506

		S.E.	Wald Z	$\Pr(> Z)$
y 10	-1.9975	0.5986	-3.34	0.0008
y 12.5	-2.4040	0.6188	-3.89	0.0001
y 15	-2.4885	0.6239	-3.99	<0.0001
y 17.5	-2.9121	0.6559	-4.44	<0.0001
age	-0.0110	0.0138	-0.79	0.4278

Effects

Response:

atelectasis_percent

	Low	High	Δ	Effect	S.E.	Lower 0.95	Upper 0.95
age	32.75	48.25	15.5	-0.1702	0.2146	-0.5908	0.2504
<i>Odds Ratio</i>	32.75	48.25	15.5	0.8435		0.5539	1.2850

	Model Likelihood Ratio Test	Discrimination Indexes	Rank Discrim. Indexes
Obs 236	LR ² 0.06	R^2 0.000	0.016
Distinct Y 8	d.f. 1	$R^2_{1,236}$ 0.000	
$Y_{0.5}$ 1	$\Pr(>^2)$ 0.8060	$R^2_{1,163.1}$ 0.000	
max log	Score ² 0.06	$\Pr(Y$	
$L/ $ 3×10^{-6}	$\Pr(>^2)$ 0.8050	median)- $\frac{1}{2}$ 0.174	

		S.E.	Wald Z	$\Pr(> Z)$
y 2.5	-0.7383	0.1491	-4.95	<0.0001
y 5	-0.9595	0.1551	-6.19	<0.0001
y 7.5	-1.2772	0.1667	-7.66	<0.0001
y 10	-2.4491	0.2458	-9.96	<0.0001
y 12.5	-2.8556	0.2906	-9.83	<0.0001
y 15	-2.9401	0.3014	-9.76	<0.0001
y 17.5	-3.3631	0.3638	-9.24	<0.0001
altitude_cat=Moderate	0.0964	0.3906	0.25	0.8051

Effects							
Response:							
atelectasis_percent							
		Low	High	Δ	Effect	S.E.	Lower 0.95 Upper 0.95
altitude_cat --- Moderate:Low		1	2		0.09641	0.3906	-0.6692 0.862
Odds Ratio		1	2		1.10100		0.5121 2.368

Multivariable model

		Model Likelihood		Discrimination		Rank Discrim.	
		Ratio Test		Indexes		Indexes	
Obs	236	LR ²	114.10	R^2	0.426		0.562
Distinct Y	8	d.f.	7	$R^2_{7,236}$	0.365		
$Y_{0.5}$	1	Pr(> ²)	<0.0001	$R^2_{7,163.1}$	0.481		
max log		Score ²	133.56	Pr(Y			
L/	1×10^{-6}			median)- $\frac{1}{2}$	0.304		
		Pr(> ²)	<0.0001				

		S.E.	Wald Z	Pr(> Z)
y 2.5	-2.4485	0.7695	-3.18	0.0015
y 5	-2.8352	0.7786	-3.64	0.0003
y 7.5	-3.4015	0.7916	-4.30	<0.0001
y 10	-5.2495	0.8584	-6.12	<0.0001
y 12.5	-5.7752	0.8830	-6.54	<0.0001
y 15	-5.8795	0.8883	-6.62	<0.0001
y 17.5	-6.3861	0.9152	-6.98	<0.0001
type_obesity=35-40	0.9039	0.4844	1.87	0.0621
type_obesity=40-45	-0.1169	0.5595	-0.21	0.8344
type_obesity=45-50	2.0150	0.5227	3.85	0.0001
type_obesity= 50	4.5168	0.5691	7.94	<0.0001
sex=Man	0.8790	0.4960	1.77	0.0764
age	0.0117	0.0156	0.75	0.4548
altitude_cat=Moderate	-0.0048	0.4607	-0.01	0.9917

Effects							
Response:							
atelectasis_percent							
	Low	High	Δ	Effect	S.E.	Lower 0.95	Upper 0.95
age	32.75	48.25	15.5	0.180700	0.2417	-0.29300	0.6544
<i>Odds Ratio</i>	32.75	48.25	15.5	1.198000		0.74600	1.9240
type_obesity --- 35---40:30---35	1.00	2.00		0.903900	0.4844	-0.04558	1.8530
<i>Odds Ratio</i>	1.00	2.00		2.469000		0.95540	6.3810
type_obesity --- 40---45:30---35	1.00	3.00		-0.116900	0.5595	-1.21400	0.9797
<i>Odds Ratio</i>	1.00	3.00		0.889600		0.29710	2.6640
type_obesity --- 45---50:30---35	1.00	4.00		2.015000	0.5227	0.99050	3.0390
<i>Odds Ratio</i>	1.00	4.00		7.501000		2.69200	20.8900
type_obesity --- 50:30---35	1.00	5.00		4.517000	0.5691	3.40100	5.6320
<i>Odds Ratio</i>	1.00	5.00		91.540000		30.01000	279.3000
sex --- Man:Woman	1.00	2.00		0.879000	0.4960	-0.09320	1.8510
<i>Odds Ratio</i>	1.00	2.00		2.409000		0.91100	6.3680
altitude_cat --- Moderate:Low	1.00	2.00		-0.004817	0.4607	-0.90780	0.8981
<i>Odds Ratio</i>	1.00	2.00		0.995200		0.40340	2.4550

Package References

- Angelo Canty, B. D. Ripley (2024). *boot: Bootstrap R (S-Plus) Functions*. R package version 1.3-30. A. C. Davison, D. V. Hinkley (1997). *Bootstrap Methods and Their Applications*. Cambridge University Press, Cambridge. ISBN 0-521-57391-2, [doi:10.1017/CBO9780511802843](https://doi.org/10.1017/CBO9780511802843).
- Auguie B (2017). *gridExtra: Miscellaneous Functions for “Grid” Graphics*. R package version 2.3, <https://CRAN.R-project.org/package=gridExtra>.
- Fox J, Weisberg S (2019). *An R Companion to Applied Regression*, Third edition. Sage, Thousand Oaks CA. <https://socialsciences.mcmaster.ca/jfox/Books/Companion/>.
- Fox J, Weisberg S, Price B (2022). *carData: Companion to Applied Regression Data Sets*. R package version 3.0-5, <https://CRAN.R-project.org/package=carData>.
- Gohel D, Skintzos P (2024). *flextable: Functions for Tabular Reporting*. R package version 0.9.5, <https://CRAN.R-project.org/package=flextable>.
- Grolemund G, Wickham H (2011). “Dates and Times Made Easy with lubridate.” *Journal of Statistical Software*, 40(3), 1-25. <https://www.jstatsoft.org/v40/i03/>.
- Harrell Jr F (2024). *Hmisc: Harrell Miscellaneous*. R package version 5.1-2, <https://CRAN.R-project.org/package=Hmisc>.
- Harrell Jr FE (2024). *rms: Regression Modeling Strategies*. R package version 6.8-0, <https://CRAN.R-project.org/package=rms>.
- Iannone R, Cheng J, Schloerke B, Hughes E, Lauer A, Seo J (2024). *gt: Easily Create Presentation-Ready Display Tables*. R package version 0.10.1, <https://CRAN.R-project.org/package=gt>.
- Jeppson H, Hofmann H, Cook D (2021). *ggmosaic: Mosaic Plots in the ‘ggplot2’ Framework*. R package version 0.3.3, <https://CRAN.R-project.org/package=ggmosaic>.
- Makowski D, Lüdtke D, Patil I, Thériault R, Ben-Shachar M, Wiernik B (2023). “Automated Results Reporting as a Practical Tool to Improve Reproducibility and Methodological Best Practices Adoption.” *CRAN*. <https://easystats.github.io/report/>.
- Müller K, Wickham H (2023). *tibble: Simple Data Frames*. R package version 3.2.1, <https://CRAN.R-project.org/package=tibble>.
- Neuwirth E (2022). *RColorBrewer: ColorBrewer Palettes*. R package version 1.1-3, <https://CRAN.R-project.org/package=RColorBrewer>.
- Peng RD (2019). *simpleboot: Simple Bootstrap Routines*. R package version 1.1-7, <https://CRAN.R-project.org/package=simpleboot>.
- Pinheiro J, Bates D, R Core Team (2023). *nlme: Linear and Nonlinear Mixed Effects Models*. R package version 3.1-164, <https://CRAN.R-project.org/package=nlme>. Pinheiro JC, Bates DM (2000). *Mixed-Effects Models in S and S-PLUS*. Springer, New York. doi:10.1007/b98882 <https://doi.org/10.1007/b98882>.
- R Core Team (2024). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.
- Rich B (2023). *table1: Tables of Descriptive Statistics in HTML*. R package version 1.4.3, <https://CRAN.R-project.org/package=table1>.

- Rinker TW, Kurkiewicz D (2018). *pacman: Package Management for R*. version 0.5.0, <http://github.com/trinker/pacman>.
- Robinson D, Hayes A, Couch S (2023). *broom: Convert Statistical Objects into Tidy Tibbles*. R package version 1.0.5, <https://CRAN.R-project.org/package=broom>.
- VanderWeele TJ, Ding P (2011). “Sensitivity analysis in observational research: introducing the E-value.” *Annals of Internal Medicine*, 167(4), 268-274. Mathur MB, VanderWeele TJ (2019). “Sensitivity analysis for unmeasured confounding in meta-analyses.” *Journal of the American Statistical Association*. Smith LH, VanderWeele TJ (2019). “Bounding bias due to selection.” *Epidemiology*.
- Wickham H (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York. ISBN 978-3-319-24277-4, <https://ggplot2.tidyverse.org>.
- Wickham H (2023). *forcats: Tools for Working with Categorical Variables (Factors)*. R package version 1.0.0, <https://CRAN.R-project.org/package=forcats>.
- Wickham H (2023). *stringr: Simple, Consistent Wrappers for Common String Operations*. R package version 1.5.1, <https://CRAN.R-project.org/package=stringr>.
- Wickham H, Averick M, Bryan J, Chang W, McGowan LD, François R, Golemund G, Hayes A, Henry L, Hester J, Kuhn M, Pedersen TL, Miller E, Bache SM, Müller K, Ooms J, Robinson D, Seidel DP, Spinu V, Takahashi K, Vaughan D, Wilke C, Woo K, Yutani H (2019). “Welcome to the tidyverse.” *Journal of Open Source Software*, 4(43), 1686. doi:10.21105/joss.01686 <https://doi.org/10.21105/joss.01686>.
- Wickham H, François R, Henry L, Müller K, Vaughan D (2023). *dplyr: A Grammar of Data Manipulation*. R package version 1.1.4, <https://CRAN.R-project.org/package=dplyr>.
- Wickham H, Henry L (2023). *purrr: Functional Programming Tools*. R package version 1.0.2, <https://CRAN.R-project.org/package=purrr>.
- Wickham H, Hester J, Bryan J (2024). *readr: Read Rectangular Text Data*. R package version 2.1.5, <https://CRAN.R-project.org/package=readr>.
- Wickham H, Vaughan D, Girlich M (2024). *tidyr: Tidy Messy Data*. R package version 1.3.1, <https://CRAN.R-project.org/package=tidyr>.
- Wood SN (2011). “Fast stable restricted maximum likelihood and marginal likelihood estimation of semiparametric generalized linear models.” *Journal of the Royal Statistical Society (B)*, 73(1), 3-36. Wood S, N., Pya, S”afken B (2016). “Smoothing parameter and model selection for general smooth models (with discussion).” *Journal of the American Statistical Association*, 111, 1548-1575. Wood SN (2004). “Stable and efficient multiple smoothing parameter estimation for generalized additive models.” *Journal of the American Statistical Association*, 99(467), 673-686. Wood S (2017). *Generalized Additive Models: An Introduction with R*, 2 edition. Chapman and Hall/CRC. Wood SN (2003). “Thin-plate regression splines.” *Journal of the Royal Statistical Society (B)*, 65(1), 95-114.
- Yee TW (2015). *Vector Generalized Linear and Additive Models: With an Implementation in R*. Springer, New York, USA. Yee TW, Wild CJ (1996). “Vector Generalized Additive Models.” *Journal of Royal Statistical Society, Series B*, 58(3), 481-493. Yee TW (2010). “The VGAM Package for Categorical Data Analysis.” *Journal of Statistical*

- Software*, 32(10), 1-34. doi:10.18637/jss.v032.i10 <https://doi.org/10.18637/jss.v032.i10>. Yee TW, Hadi AF (2014). “Row-column interaction models, with an R implementation.” *Computational Statistics*, 29(6), 1427-1445. Yee TW (2024). *VGAM: Vector Generalized Linear and Additive Models*. R package version 1.1-10, <https://CRAN.R-project.org/package=VGAM>. Yee TW (2013). “Two-parameter reduced-rank vector generalized linear models.” *Computational Statistics and Data Analysis*, 71, 889-902. Yee TW, Stoklosa J, Huggins RM (2015). “The VGAM Package for Capture-Recapture Data Using the Conditional Likelihood.” *Journal of Statistical Software*, 65(5), 1-33. doi:10.18637/jss.v065.i05 <https://doi.org/10.18637/jss.v065.i05>. Yee TW (2020). “The VGAM package for negative binomial regression.” *Australian and New Zealand Journal of Statistics*, 62(1), 116-131.
- Zeileis A, Köll S, Graham N (2020). “Various Versatile Variances: An Object-Oriented Implementation of Clustered Covariances in R.” *Journal of Statistical Software*, 95(1), 1-36. doi:10.18637/jss.v095.i01 <https://doi.org/10.18637/jss.v095.i01>. Zeileis A (2004). “Econometric Computing with HC and HAC Covariance Matrix Estimators.” *Journal of Statistical Software*, 11(10), 1-17. doi:10.18637/jss.v011.i10 <https://doi.org/10.18637/jss.v011.i10>. Zeileis A (2006). “Object-Oriented Computation of Sandwich Estimators.” *Journal of Statistical Software*, 16(9), 1-16. doi:10.18637/jss.v016.i09 <https://doi.org/10.18637/jss.v016.i09>.