

Wilczyńska Agnieszka, Wawryków Agata, Korabiusz Katarzyna, Niedzielska Mirela, Janik Inga, Fabian Danielewska Anna, Maciejewska Martyna, Gutowski Piotr. The use of complex decongestive therapy in the process of rehabilitation in case of primary and secondary lymphatic insufficiency. *Journal of Education, Health and Sport*. 2018;8(01):86-94. eISSN 2391-8306. DOI <http://dx.doi.org/10.5281/zenodo.1154929> <http://ojs.ukw.edu.pl/index.php/johs/article/view/5211> <https://pbn.nauka.gov.pl/sedno-webapp/works/848545>

The journal has had 7 points in Ministry of Science and Higher Education parametric evaluation. Part B item 1223 (26.01.2017).
1223 Journal of Education, Health and Sport eISSN 2391-8306 7

© The Authors 2018;
This article is published with open access at Licensee Open Journal Systems of Kazimierz Wielki University in Bydgoszcz, Poland

Open Access. This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited. This is an open access article licensed under the terms of the Creative Commons Attribution Non Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted, non commercial use, distribution and reproduction in any medium, provided the work is properly cited.

This is an open access article licensed under the terms of the Creative Commons Attribution Non Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted, non commercial use, distribution and reproduction in any medium, provided the work is properly cited.

The authors declare that there is no conflict of interests regarding the publication of this paper.
Received: 05.12.2017. Revised: 15.12.2017. Accepted: 18.01.2018.

The use of complex decongestive therapy in the process of rehabilitation in case of primary and secondary lymphatic insufficiency

Agnieszka Wilczyńska¹, Agata Wawryków¹, Katarzyna Korabiusz¹, Mirela Niedzielska¹, Inga Janik¹, Anna Fabian-Danielewska², Martyna Maciejewska¹, Piotr Gutowski³

¹ Pomorski Uniwersytet Medyczny w Szczecinie, Studium Doktoranckie Wydziału Nauk o Zdrowiu, ul. Żołnierska 54, 71-210 Szczecin

² Pomorski Uniwersytet Medyczny w Szczecinie, Studium Doktoranckie Wydziału Lekarskiego, ul. Żołnierska 54, 71-210 Szczecin

³ Pomorski Uniwersytet Medyczny w Szczecinie, Klinika Chirurgii Naczyniowej, Ogólnej i Angiologii ul. Powstańców Wlkp. 72, 70-111 Szczecin

¹Pomeranian Medical University in Szczecin, PhD Studies at the Faculty of Health Sciences

²Pomeranian Medical University in Szczecin, PhD Studies at the Faculty of Medicine

³Pomeranian Medical University in Szczecin, Department of Vascular, General and Angiology Surgery

Abstract

The purpose of this study is to draw attention to the problems connected with the failure of the lymphatic system. In particular the primary and secondary swelling. The reference in this work to the scientific literature indicates a growing need for the development of Lymphology in Poland. It indicates a significant problem, which is the lack of specific standards and procedures of treatment and methods of treatment of disorders of lymph in the educational process of medical schools in Poland.

Key Words: lymphedema, physiotherapy

Introduction

Diseases associated with the failure of the lymphatic system are among the diseases in which diagnosis and treatment are still struggling with not formally established method of prevention work and decongestants. It is also worth noting that the Polish nomenclature of medical specializations does not include Lymphology, which will determine the standards and procedures of treatment and methods of treatment of disorders of the lymph. As a not popularized area it is still struggling with not enough support of the medical services, which further lowers its essence in medical rehabilitation.

Lymphology in Europe is practiced in an unconventional way since 1936. Until its mode of action and therapeutic efficiency has been scientifically proven. It is proven by the popularity of specialized clinics, eg. In Germany (Hinteranter, Fachklinik für Lymphologie-Földi Klinik, Dr. Feldbergklinik Asdonk in the Black Forest). Lymphoedema treatment precursor was the Danish philologist, sociologist and later physiotherapist Dr Emil Vodder (1893-1986). He introduced the term „manual lymphatic drainage” (MLD- manual lymphatic drainage) (45). Also deserving recognition is German physician Dr. Johannes Asdonk (1910-2003). He enriched therapy by additional special grips and compression therapy. He published his results in 1972 and introduced the method of German clinics under the name „Physiotherapy edema By Asdonk and Barteczko”. The effectiveness of this method has been also shown by the work conducted by prof. Kuhnke Bonn thanks to them from 1974., rehabilitation of lymphatic insufficiency is recognized and reimbursed by the German health insurance funds.

Current situation of Lymphology in Poland also points to the growing interest in the discipline. This is proven by the analysis of medical publications conducted by the National Board of Physiotherapy (2). The first position on the market concerning the matter of Lymphology written by dr n. Med. Janusz Dosia entitled „Lymphology" also deserves credit(3). The author refers to issues related to disorders of the lymphatic system. He points to various factors of developing lymphatic insufficiency, diagnosis and the process procedure in the

reduction of disorders of lymph. Established in Chorzów Polish Society of Lymphology substantially affected the development of Lymphology in Poland. It is one of the most important centers of rehabilitation and training in our country. It also contributes to the development of new centers, ie. In Warsaw, Krakow and Poznan.

Despite such extensive resources for antioedema rehabilitation, there is a noticeable trend in education (at universities) directed only toward the side effects of mastectomy rather than the failure and diseases of the lymphatic system. Only interdisciplinary collaboration / and change in education can contribute to the consolidation of knowledge and development of Lymphology in Poland.

The lymphatic system is a semi-open system of vessels and ducts, in which lymph flows (3). The main source of lymph is the filtrate of the blood plasma (through the vessel wall), secretion of cells and the gastrointestinal tract. The composition of lymph is similar to blood plasma. It contains proteins, fats, water, minerals and large amounts of white blood cells (which shows the dense structure and slow movement). About 90% of the system is subcutaneous parenchyma, and only 10% of the deep vasculature system.

The pathophysiology of lymphedema

Lymphedema is often defined as a chronic condition (3), which is the result of mechanical failure (aplasia, valvular dysplasia), dysfunctional vessels, or both. Mainly due to the accumulation of high protein fluid in parenchyma and lymphatic vessels. The migrating and resident immune cells, metabolic products, apoptosing cells cause, as a result of standstill, proliferation of keratinocytes, fibroblasts and thus collagen deposition (5,6,7,8) binding water molecules. Simplified pathophysiology of oedema, enabled their classification by dividing them into primary and secondary (3).

Primary oedema is frequently caused by genetic factors determining the abnormal growth of lymphatic vessels (97-99% spontaneous mutation; 1-3% change hereditary). This form is substantially asymmetrical and mostly concerns the lower limbs. Moreover, an oedema of parenchymatous organs (liver, lung, intestine) and CNS. Depending on the age of the patients clinical symptoms, characterized by the primary lymphatic insufficiency can be divided as follows:

- 1) Lymphoedema congenital - current clinical picture at birth or shortly after birth (up to 1 month of age).
- 2) The early lymphatic swelling - appeared before 35 years, most often affects people puberty.

3) Late Lymphedema swelling- appeared after 35 years

Secondary oedema mainly caused by damage to the lymphatic system. There is noticeable lack of information about frequency of lymphatic problems. It is estimated that 1.8% of the population (2% of women and 1.5% men) suffer from the lymphatic oedema. The main causes are:

1) Malignant tumors (approximately 4%) as a growing form of cancer usually causes stasis, of lymph and as a lead to intratissue exudates, such as: malignant melanoma, prostate cancer, malignant gynecological tumors (11)

2) Infection (bacteria, parasites, viruses, fungi) (9). Especially recurrent episodes of bacterial inflammation of the lymphatic vessels. The most common inflammatory pathogens include Streptococcus, nematodes (which cause filariasis). According to WHO, about 90 million of the people worldwide may be infected by nematodes (12,13,14,15,16).

3) Chronic skin diseases (acne rosacea, acne vulgaris, psoriasis, atopic dermatitis) (17, 18, 19,20)

4) Rheumatic diseases (rheumatoid arthritis, chronic articular and periarticular inflammations) by autoimmune destruction of lymph vessels. However, the etiological factor is unknown (21).

5) Tissue damage due to burns, crushing, fracture (22)

6) Iatrogenic damage caused by surgery, radiation (23, 24, 25, 26, 27, 28, 29, 30)

7) Vascular diseases (recurrent vasculitis syndrome, post-thrombotic syndroms (31, 32)

8) Pharmacotherapy, mainly as a side effect after hydrochloride pentazocine application. It can lead to lymph vessels sclerosis and obstruction (10).

9) Pregnancy, as a result of expanding uterus, and therefore the pressure on the vessel around the inferior vena cava and the iliac vein (33, 34)

In each sort of swelling (primary or secondary) is a huge noticeable impact on the physical and mental health of the patient. Chronic swelling significantly impedes the proper functioning and comfort. In the worst cases contributes to inflammation (erysipelas), structural changes within the osteoarticular system, vascular (felt stiffness, rest pain, local problem of mobility) (46, 47, 48).

Referring to the guidelines of International Society of Lymphology and the British Lymphology Society (BLS) (1) in the case of lymphatic insufficiency it is recommended to use the Complex Decongestive Therapy. Developed by renowned specialists include:

1. Manual Lymphatic Drainage- delicate hand-made system of tricks designed to activate lymphatic vessels, relaxation of interstitial fibrosis, thus affecting the aid penetration of the interstitial fluid to the lymphatic vessels (49).

2. Compression (multilayer bandage or use of compression). The use of external pressure is an essential component of the treatment of oedema. It supports preferably the effect of lymphatic drainage by increasing resorption of lymphatic vessels and veins. It is an essential component of work with edema in its initial stage of formation. Then replaced with a compression device designed individually for the corresponding pressure dependent upon the problem of lymphatic insufficiency (36,37,38).

3. Specialized kinesiotherapy. Exercises to improve lymph drainage - focusing on active work diaphragm (cardio-pulmonary exercises) and the mobilization of the muscle pump (for upper limb oedema) and triceps calf muscle (in the case of lower extremity edema). The literature indicates that these exercises are carried out with the appropriate compression products using a multilayer compression bandage or clothing (39,41) that reduce the risk of oedema recurrence. Particularly effective is the scientific progress taking into account the use of such forms of activity as PNF, Pilates, Yoga (40,42,43)

4. Patient education in the aspect of home care. It is an indispensable parameter therapy and thus is an element of cooperation of patient and therapist. It is based on substantive and practical preparation of the patient to work at home. The patient must be aware of the essence of his chronic disease, and must follow the basic principles relating to the skin care. Must protect his skin against damage, use appropriate clothing, cosmetics (44). This also applies to self lymphatic massage and home gymnastics.

References

- 1) Lymphoedema Framework. Best Practice for Management of Lymphoedema. International Consensus. MEP Ltd., London 2006;14
- 2) Taradaj J.: „Analiza skuteczności poszczególnych procedur fizjoterapeutycznych w leczeniu obrzęku limfatycznego: rekomendacje w świetle Evidence Based Medicine (EBM)”, Grupa Ekspertów ds. Nauki przy Krajowej Izbie Fizjoterapeutów, www.kif.info.pl
- 3) Gültig O., Miller A., Zöltzer H., pod red. Wyd. Pol. Doś J.: „Limfologia“, wyd. Edra, Wrocław 2017
- 4) Ochalek K., Grądalski T.: Zastosowanie ręcznego drenażu limfatycznego w chorobach naczyń, *Acta Angiology*, 2011, vol. 17(3): 189-198
- 5) Daroczy J.: Pathology of Lymphoedema. *Clin. Dermatol* 1995;13:433-44
- 6) Wiktor M.: Patofizjologia obrzęku chłonnego, [w:] *Obrzęk chłonny*. Red. Chęciński P., Termedia Wydawnictwa Medyczne, Poznań 2010
- 7) Piotrowicz R., Ciecierski M., Jawień A.: Obrzęki limfatyczne – patomechanizm i diagnostyka, *Przewodnik Lekarza*, 2000, 7, 70-72
- 8) Kozikowska J., Łuczak J.: Obrzęk limfatyczny - patomechanizm, podział, zasady leczenia, *Praktyka Medyczna, Przewodnik Lekarza*, 48 – 54
- 9) Schirger A.: lymphedema. *Cardiovasc Clin* 13:293, 1983
- 10) M. A. Creager, V. J. Dzau, Loscalzo J.: *Choroby naczyń*. Redakcja naukowa wydania polskiego Adamiec R., Wydawnictwo Czelej, Lublin 2008
- 11) Gashev A: Physiologic aspects of lymphatic contractile function: Current perspectives. *Ann N Y Acta Sci* 979:178,2002
- 12) Lymphatic filariasis-tropical medicine's origin will not go awal. *Lancet (Editorial)* 1:1409,1987.
- 13) Dandapat MC, Mahapatro SK, Dash DM: Management of chronic manifestations of filariasis. *J Indian Med Assoc* 84:219,1986.
- 14) Baird J, K, Alpert LI, Friedman R. et al: North American brugian filariasis: Report of nine infections of humans. *Am J Trop Med Hgz* 35:1205, 1986
- 15) Clark WR, Lieber MM, Genital filariasis in Minnesota. *Urology* 28:518,1986
- 16) Leonard JC, Humphrey GB, Basmadjian G: Lymphadema secondary to filariasis. *Clin Nucl Med* 10:203, 1985
- 17) Abe R, Kimura M, Airosaki A: Retroperitoneal lymphangiomyomatosis with lymphodema of the legs. *Lymphology* 13:62, 1980

- 18) Lynde CW, Mitchell IC: Unusual complication of allergic contact dermatitis of hands-recurrent lymphangitis and persistent lymphoedema. *Contact Dermatitis* 8:279, 1982.
- 19) Kyle VM, DeSilva M, Hurts G: Rheumatoid lymphoedema. *Clin Rheumatol* 1:126,1982
- 20) Reed Burke S, Bozeman M, et al: Lymphoedema of the lower abdominal wall in pregnancy. *J Am Acta Dermatol* 12:930, 1985
- 21) Majerski I: Lymphoedema tarda. *Cutis* 38:105, 1985
- 22) Brunning I, Gibson AG, Peery M: Oedema bleu: A reappraisal. *Lancet* 1:810, 1980
- 23) Szuba A, Rockson S: Lymphedema: A review of diagnostic techniques and therapeutic options. *Vasc Med* 3:145, 1998
- 24) Rockson SG, Precipitating factor in lymphedema: Myths and realities *Cancer* 83:2814,1998
- 25) Hojris I, Andersen J, Overgaard M, Overgaard J.:Late treatment related morbidity in breast cancer patients randomized to postmastectomy radiotherapy and systemic treatment versus systemic treatment alone. *Acta Oncol* 39:355, 2000
- 26) Tengrup I, Tennval-Nittby L, Christiansson I, Laurin M.: Arm morbidity after breast-conserving therapy for breast cancer. *Acta Oncol* 39:393, 2000
- 27) Pettek JA, Heelan MC: Incidence of breast carcinoma-related lymphoedema. *Cancer* 83 (suppl 12):2776, 1998
- 28) Fiorica JV, Roberts WS, Greenberg H, et al: Morbidity and survival patterns in patients after radical hysterectomy and postoperative adjuvant pelvic radiotherapy. *Gynecol Oncol* 36:343, 1990
- 29) Werngren-Elgstrom M, Lidman D: Lymphoedema of the lower extremities after surgery and radiotherapy for cancer of the cervix. *Scand J Plast Reconstr Surg Hand Surg* 28:289, 1994
- 30) Soisson AP, Soper JT, Clarke- Peterson DL, et al: Adjuvant radiotherapy following radical hysterectomy for patients with stage IB and IIA cervical cancer. *Gynecol Oncol* 37:390, 1990
- 31) Drażkiewicz M, Góral M, Główska B, Pietrzak K.: Infekcje protezy aortalno-dwuudowej-trudności w rozpoznaniu. *Chirurgia Polska* 2013, 15,1, 98-103, ISSN 1507-5524
- 32) Niedźwiadek J, Mazur E, Terlecki P et al.: Czynniki etiologiczne zakażeń przeszczepów naczyniowych i ocena ich lekooporności. *Pol Merk Lek.* 125:423-426, 2006
- 33) Gunther H, Kohlrausch W, Teirich-Leube H.: *Gimnastyka w ginekologii i położnictwie.* Państwowy Zakład Wydawnictw Lekarskich, Warszawa 1968, 101-105
- 34) Wawryków A, Korabiusz K, Torbe A, Lubkowska A.: Edema In pregnant women-possibility of physiotherapeutic treatment. *Journal od Education, Health and Sport.*2017:7(6):136-144, eISSN 2391-8306. DOI

- 35) Ochalek K.:The role of compression therapy in venous-lymphatic disturbances — case report, *Acta Angiol*, vol. 18, No. 4 pp. 169–176
- 36) Ochalek K, Gradalski T, Partsch H.: Preventing early postoperative arm swelling and lymphadema manifestation by compression sleeves after axillary lymph node interventions in breast cancer patients: A randomized controlled trial. *J Pain Symptom Manage*. 2017 Aug 7. Pii: S0885-3924(17)30342-1 [Epub ahead of print]
- 37) Yuksel A, Gurbuz O, Velioglu Y, Kumtepe G, Senol S.: Management of Lymphoedema. *Vasa*. 2016; 45(4):283-91
- 38) Chang DW, Masia J, Garza R 3 rd, Skoracki R, Neligan PC.Lymphedema: Surgical and Medical Therapy. *Plast Reconstr Surg*. 2016 Sep; 138 (3 Suppl):209S-18S
- 39) NLN Medical Advisory Committee. The diagnosis and the treatment of lymphedema. Position statement of the national lymphedema network. Feb.2011 (online)
- 40) Hwang WT, Jeong YJ, Kim SY, Jeong YG.: Effects of proprioceptive neuromuscular facilitation stretching and deep-breathing exercises on upper extremity lymphedema in stroke patients. *J Phys Ther Sci*. 2016 Dec; 28(12):3276-3278
- 41) Fukushima T, Tsuji T, Sano Y, Miyata C, Kamisako M, Hohri H, Yoshimura C, Asakura M, Okitsu T, Muraoka K, Liu M.: Immediate effect of active exercise with compression therapy on lower-limb lymphedema. *Support Care Cancer*. 2017 Aug; 25(8):2603-2610
- 42) Sener HO, Malkoc M, Ergin G, Karadibak D, Yavuzsen T.:Effect of clinical Pilates Exercises on Patients Developing Lymphedema after breast cancer treatment: A randomized Clinical Trial. *J Breast Health (2013)*. 2017 Jan 1; 13(1):16-22
- 43) Loudon A, Barnett T, Williams AD, Visentin D, Imminik MA, Piller N.:Guidelines for teaching yoga to women with breast cancer related lymphedema: an evidence-based approach. *Int J Yoga Therap*. 2017 Aug 1. Doi:10.17761/IJYT2017 [Epub ahead of print]
- 44) Biały S, Chmielewska D.: Aktywny udział pacjenta w terapii obrzęku limfatycznego- niezbędne informacje dla terapeuty i pacjenta. *Rehabilitacja w Praktyce, dodatek: Limfologia w Praktyce*, 1/2012. Wydawnictwo Elamed.
- 45) 17. Lymphoedema Framework. Best Practice for Management of Lymphodema. International consensus. MEP Ltd., London 2006;14
- 46) Piotrowska R., Dobosz M., Książek J., Halena G., Jakość życia chorych z miażdżycą naczyń obwodowych, przegląd piśmiennictwa. *Ann. Acad.Medd. Gedan*. 2011,41,89-95
- 47) Drażkiewicz T. Własne doświadczenia w zwalczaniu zakażeń miejscowych i ogólnych u chorych z zawnosowanym przewlekłym niedokrwieniem kończyn dolnych. *Pol Przegl Chir*. 1993; 65: 163–70.

- 48) Cnotliwy M. Powikłania chłonne w pachwinach po operacjach naprawczych tętnic. Rozprawa Habilitacyjna. PUM, Szczecin 2003.
- 49) Piotrowicz R, Ciecierski M, Jawień A.: Obrzęk limfatyczny-leczenie, Termedia Przewodnik Lekarza 7/2000