


Changing-look NLS1s?

*Victor L. Oknyansky,
K. L. Malanchev,
(Moscow State University)*

*Martin Gaskell
(University of California at Santa Cruz)*



A common cartoon of a thermal AGN:



Two of the biggest challenges to simple AGN unification schemes:

1. **NLS1s**
2. **“Changing-look” (CL) AGNs.**

∴ important to look at the relationship between these two phenomena.


(After Urry & Padovani 1995; see Keel 1980; Krolik & Begelman 1988; Antonucci 1993)

A Unified Picture for *All Super-Eddington NLS1s*

Super-Eddington Narrow-Line Seyfert 1s

'Simple' NLS1s
 (e.g. RX J0439.6-5311, PG 1244+026, RE J1034+396,
 RX J1140.1+0307, Ton S180)

'Complex' NLS1s
 (e.g. 1H 0707-495, Mrk 335,
 PG 1211+143)



(Jin et al., 2017b, submitted)

Different types of CL AGNs

1. Vanishing or strong decreasing of broad-line components over periods from months to years.

Examples:

NGC 3516 (Andrillat, 1968)

NGC 4151 (Lyutyj, Oknyanskij, Chuvaev 1984; Penston and Perez, 1984; Oknyansky, Lyuty & Chuvaev 1991)

NGC7603, Mrk 372, 3C390.3. Mrk993, NGC7469
(Chuvaev, Lyutyi & Doroshenko 1990)

NGC 6814 (Sekiguch & Menzies, 1990),

NGC 5548 (Shapovalova et al. 2004).

- There are more than 20 known CL QSOs (Ruan et al. 2016; MacLeod et al. 2016;)
- [See references in Shappee et al. (2014) and Koay et al. (2015); Yang et al. 2017].

2. Appearance of broad-line components in AGNs which are usually previously seen as type-2 AGNs.

Examples:

Mrk 6, Mrk 1018, NGC 1097, NGC 7582, Mrk 590, NGC 2617.

In some cases they return to their usual low state (a type-2 appearance) after few years (e.g., NGC 7582) or after very long time (e.g., Mrk 6 and Mrk 590).

For some of these objects the changing look is connected with appearing of a blue shifted broad emission components (e.g., Mrk 6, 3C 390.3)

**NOTE: CL AGN ARE NOT A RARE PHENOMENON!
ALMOST EVERY WELL-MONITORED AGN SHOWS
THE PHENOMENON!**

NGC 4151: a Seyfert 2 in a deep photometric minimum

V. M. Lyutyĭ, V. L. Oknyanskiĭ, and K. K. Chuvaev






FIG. 1. Comparative variations in the optical continuum (U magnitude) of the NGC 4151 nucleus and the ratio $W_{\lambda}(H\beta)/W_{\lambda}(N_2)$. The lower envelopes outline the slow fluctuations.



Fairall 9, 1981-4



Kollatschny & Fricke 1985


Additional types of CL AGNs

3. More than 20 AGNs are changing-look AGNs in X-ray properties (see references in [Reachi et al. 2016](#)). Some had also been noted previously as being changing-look AGNs in the optical (NGC4151 and Mrk6). **NLS1s ARE FAMOUS FOR THEIR STRONG X-RAY VARIABILITY!**
4. Other changing-look AGNs which do not fit in categories 1 – 3. Objects with strong continuum variation of factors of hundred without significant variation of lines. *E.g.*, changing from a QSO to BL Lac state (QSO 1256+295, [Wills et. al. 1983](#)).
5. TDEs observed in non-AGN galaxies
6. CL NLS1? (few cases of CL for each of these types are suspected)

Are all CLQs part of the low Eddington ratio tail, or are some one-off events with different physics, *e.g.*, TDEs? ([MacLeod, 2017](#))

NGC 2617

“The man behind the curtain...” Shappee et al. (2014)




Multi-Wavelength Monitoring of the Changing-Look AGN NGC 2617 During State Changes


V.L. Oknyansky (M.V. Lomonosov Moscow State University, Moscow, Russia), V.M. Lipunov, N.I. Shatsky, E.S. Gorbovskoy, A.M. Tatarnikov, V.G. Metlov, M.A. Burlak, K.L. Malanchev, C.M. Gaskell (UCSC, USA), N.A. Huseynov (Shamakhy Astrophysical Observatory, Azerbaijan), I.A. Alikberov, Kh.M. Mikailov, S.S. Tsygankov (Tuorla Observatory, Piikkiö, Finland), M.B. Brotherton (University of Wyoming, Laramie, USA), D. Kasper, P. Du (Institute of High Energy Physics, Chinese Academy of Sciences, Beijing, China), X. Chen (Handong University, Shandong, China), D.A.H. Buckley (South African Astronomical Observatory, Observatory, South Africa), M. Gromadzki (University of Warsaw, Poland), H. Winkler (University of Johannesburg, Auckland, South Africa), B. Paul, F. van Wyk, R. Rebolo (Instituto de Astrofisica de Canarias, Tenerife, Spain), M. Serra-Ricart, R. Podesta (OFA, San Juan, Argentina), H. Levato (ICATE, San Juan, Argentina)



Our observations of the NGC 2617


- IR *JHK* – 2.5m, CMO SAI, (2016 – 2017)
- *BVRI* – AZT-5, CS SAI MSU (2016 – 2017)
- *BVRI* – 0.6m, CS SAI MSU (2016 – 2017)
- *BVRI* – 0.6m, ShAO, (2016 – 2017)
- *BV* – 1m, China, Shandong (2016 – 2017)
- *WBV* – MASTER (2010 – 2017)
- Optical spectra (2 nights) – 1.9m, SAAO (2017)
- Optical spectra (18 nights) – 2m, ShAO (2016 – 2018)
- Optical spectra (68 nights) – 2.3m, WIRO WU (2017 – 2018)
- Optical spectra (4 nights) – 11m , SALT, SAAO (2017–2018)
- X-Ray, UV, *BV* – *Swift* (2013 – 2018)





2013

2016




The curtain remains open: NGC 2617 continues in a high state

V. L. Oknyansky ✉; C. M. Gaskell; N. A. Huseynov; V. M. Lipunov; N. I. Shatsky; S. S. Tsygankov; E. S. Gorbovskoy; Kh. M. Mikailov; A. M. Tatarnikov;

Table 1.


Lags from the reverberation mapping analysis with the MCCF and JAVELIN methods in days. MCCF and JAVELIN 1σ confidence limits are presented.

	MCCF		JAVELIN	Time
	τ_{peak}	τ_{cent}	τ_{JAV}	
<i>J</i> from <i>B</i>	$2.8^{+1.2}_{-0.8}$	$3.0^{+1.0}_{-1.2}$	$2.7^{+1.3}_{-1.4}$	2016
<i>H</i> from <i>J+B</i>	$18.2^{+3.0}_{-4.0}$	$13.9^{+4.0}_{-3.0}$	$10.7^{+1.2}_{-1.2}$	-
<i>K</i> from <i>J</i>	$21.5^{+2.4}_{-2.6}$	$20.5^{+2.2}_{-2.4}$	$19.9^{+1.4}_{-1.6}$	-
<i>K</i> from <i>J+B</i>	$25.3^{+3.4}_{-2.3}$	$24.5^{+3.1}_{-2.6}$	$26.0^{+2.1}_{-2.1}$	-
<i>UV</i> from <i>Xray</i>	$2.0^{+0.7}_{-0.5}$	$1.5^{+0.7}_{-0.5}$	$2.7^{+0.4}_{-0.2}$	-
<i>UV</i> from Γ X	$0.8^{+0.5}_{-0.5}$	$0.9^{+0.4}_{-0.2}$	$0.8^{+0.3}_{-0.3}$	-
<i>BW</i> from <i>UV</i>	$1.0^{+0.5}_{-0.5}$	$0.5^{+0.5}_{-0.5}$	$0.6^{+0.3}_{-0.5}$	-
<i>UV</i> from <i>Xray</i>	$2.5^{+0.5}_{-0.7}$	$2.9^{+0.5}_{-0.8}$	$2.5^{+0.6}_{-0.8}$	2013–2014
<i>BS</i> from <i>UV</i>	$0.4^{+0.4}_{-0.4}$	$0.6^{+0.3}_{-0.3}$	$0.4^{+0.2}_{-0.2}$	-




Volume 467, Issue 2

May 2017



SUMMARY (just for NGC2617)


- NGC 2617 remains in a high (Seyfert1) state, but from the end of the March 2017 till the April 2018 the object was in very low level of brightness and variability. **The X-ray flux was at an all-time low in December 2017 (lowest since monitoring began in 1982.)** The spectral type was **Sy 1.7**. We suspect that this is the transient state of the object and **we expect the changing of its type back to Sy 1.8 soon.**
- Light-travel-time delays (X-ray  UV, Opt., IR) increase with wavelength, but **hard X-rays lag to soft ones ~0.1 day.**
- New IR data from October 2016 to May 2017 confirm our published (2016) result: **K band lags ~ 25 days relative to UV/optical. This lag is probably smaller now (~15 days) and we suspect it is result of dust recovering in the minimum state.**
- The reason for a significant change in the luminosity of the object is not a change in absorption alone. **We propose that a change in the energy-generation rate led to a change in absorption.** What caused the change in energy generation remains the main mystery.

Proposed model of IR emission

(Oknyansky & Gaskell 2015;
Oknyansky et al. 2015,2 017;
Gaskell & Harrington 2018)

Sublimation and recovering of dust on the line of sight (following the strong luminosity variations) can explain CL events.

*[Thanks to graphics designer
Natalia Sinugina]*



CL NLS1 ?

Optical spectra

1) Narrow line

NLS1  BLS1?

2) Strong Fe II lines

NLS1  NLS2 ?

few cases of the broad emission lines
appearance :

PS16dtm (Blanchard et al. 2017) TDE? PS16dtm
CSS100217 (Drake et al. 2011) SN? NLS1 CLQ (MacLeod, 2017)

3) HeII $\lambda 4686$ nearly disappeared in NGC 4051 (Peterson et al. 2000)



X-ray spectra

1) Compton-thin  Compton-thick
NGC 4051 (Guainazziet al. 1998)

2) Soft weak  Not weak
PG 1535+547 (Ballo et al. 2017)

NLS1 CLQ

- $\log(L/L_{\text{Edd}}) = -1.057$, Strong FeII line change
- Similar to TDE candidate (Blanchard+17)



From MacLeod et al. (2016)

Post Script

- Tidal disruption events occur when a star passes too close to a massive black hole and is totally ripped apart by tidal forces. It may also happen that the star is not close enough to the black hole to be totally disrupted and a less dramatic event might happen. If the stellar orbit is bound and highly eccentric, just like some stars in the centre of our own Galaxy, repeated flares should occur. (Campana et al., 2015; Ivanov & Chernyakova, 2006).

It is my pleasure to thank the organizers of the conference for their invitation and all of you for attention.

*Thanks to **Pavel Ivanov** and **Nikolay Shakura** for useful discussions, thanks to the **Swift** and **MASTER** teams for organizing the observations.*

RadioNet has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 730562