

# Ecto- and endo-parasitic monogeneans (Platyhelminthes) on cultured freshwater exotic fish species in the state of Morelos, South-Central Mexico

Edgar F. Mendoza-Franco<sup>1</sup>, Juan Manuel Caspeta-Mandujano<sup>2</sup>,  
Marina Tapia Osorio<sup>3</sup>

**1** Instituto de Ecología, Pesquerías y Oceanografía del Golfo de México (EPOMEX), Universidad Autónoma de Campeche, Campeche, México **2** Facultad de Ciencias Biológicas y Centro de Investigaciones Biológicas, Universidad Autónoma del Estado de Morelos, Cuernavaca, Morelos, Mexico **3** Centro Nacional de Servicios de Constatación en Salud Animal Jiutepec, Morelos, México

Corresponding author: *Edgar F. Mendoza-Franco* ([efmendoz@uacam.mx](mailto:efmendoz@uacam.mx))

---

Academic editor: *D. Gibson* | Received 24 April 2018 | Accepted 6 July 2018 | Published 26 July 2018

---

<http://zoobank.org/2ACDA693-4496-4E91-AA67-D36232377E7D>

---

**Citation:** Mendoza-Franco EF, Caspeta-Mandujano JM, Osorio MT (2018) Ecto- and endo-parasitic monogeneans (Platyhelminthes) on cultured freshwater exotic fish species in the state of Morelos, South-Central Mexico. ZooKeys 776: 1–12. <https://doi.org/10.3897/zookeys.776.26149>

---

## Abstract

An extensive parasitological study of 365 freshwater exotic fish specimens belonging to 13 species of seven families (Cichlidae, Cyprinidae, Osphronemidae, Pangasidae, Poeciliidae, Characidae, and Loricariidae) collected from 31 Aquaculture Production Units (APU) from Central Mexico revealed the occurrence of 29 ecto- and endo-parasitic monogeneans found on gills and stomachs: *Cichlidogyrus sclerosus*, *C. thurstonae*, *C. tilapiae*, *Cichlidogyrus* sp. 1, *Cichlidogyrus* sp. 2, *Enterogyrus coronatus*, *E. malmbergi*, *Gussevius spirallocirra*, *Sciadicleithrum iphthimum*, *Sciadicleithrum* sp., *Scutogyrus longicornis* (all Dactylogyridae), *Gyrodactylus cichlidarum*, and *G. yacatli* (Gyrodactylidae) on *Oreochromis niloticus*, *Pterophyllum scalare* and *Hemichromis* sp. (Cichlidae); *Dactylogyrus baueri*, *D. formosus*, *D. intermedius*, *D. vastator*, *D. extensus*, *Dactylogyrus* sp. (all Dactylogyridae), and *G. kobayashii* on *Carassius auratus*, *Cyprinus carpio* and *Ctenopharyngodon idella* (Cyprinidae); *Trianchoratus acleithrium* and *T. trichogasterium* (Dactylogyridae) on *Trichogaster trichopterus* (Osphronemidae); *Thaparocleidus caecus*, *T. siamensis* (Dactylogyridae), and Dactylogyridae sp. on *Pangasianodon hypophthalmus* (Pangasidae); *G. poeciliae* on *Poecilia reticulata* (Poeciliidae); *Diaphorocleidus armillatus* (Dactylogyridae) on *Gymnocorymbus ternetzy* (Characidae); *Unilatus unilatus* (Dactylogyridae) and Gyrodactylidae sp. on *Hypostomus* sp. (Loricariidae). The paramount importance of the establishment of these monogeneans due to the importation/exportation of non-native ornamental and other exotic host fish species cultured for food in Mexico is briefly discussed. Quarantine is recommended for all transferred host species.

## Keywords

characids, cichlids, cyprinids, fish introductions, loriciariids, Monogenea, Morelos state, non-native ornamental fish, osphronemids, pangasids, parasites, poeciliids, quarantine, tilapia

## Introduction

At a global level, increasing attention is being paid to generate useful ecological indicators that favor invasiveness and geographic range expansion by introduced species (Lavergne and Molofsky 2007, Blackburn and Ewen 2017). Conjointly, introductions of species are rising sharply because of increased trade, transport, travel, and tourism associated with globalization (IPPC Secretariat 2005). Within this context, trade of the non-native ornamental fish industry and/or fish farms for food production, has been the main cause of introductions of fish and their parasites around the world (Barroso de Magalhães and Jacobi 2013, Mendoza et al. 2015). Furthermore, the same industries pose a growing threat to native wildlife if non-native fishes are later released into the wild (see Mendoza-Franco et al. 2012). Culture of non-native ornamental and food fishes represents major activities in the state of Morelos (south-central Mexico) since these fishes are commercially distributed within and outside of Mexico in large quantities (Martínez et al. 2010).

Although non-native aquatic organisms are important to Morelos aquaculture and the economy of the state of Morelos, the aquaculture industry should be made aware of the considerable local, state, and national concern over the potential ecological or economic problems arising from non-native fish introductions and their parasites in natural environments (i.e., parasite transfer and/or fish competition with native species) (Barroso de Magalhães and Jacobi 2013). Recently, a total of 44 helminth species on introduced freshwater fishes were listed for Mexico, of which five are invasive species, i.e., *Cichlidogyrus sclerosus* Paperna & Thurston, 1969 *Dactylogyrus extensus* Mueller & Van Cleave, 1932 and *Gyrodactylus cichlidarum* (Paperna 1968) García-Vasquez & Hansen, 2007 (Monogenea); *Centrocestus formosanus* (Nishigori 1924) Price, 1932 (Digenea) and *Schyzocotyle acheilognathi* Yamaguti, 1934 (Cestoda), all of them introduced with their Asian and African hosts (Tapia Osorio et al. 2014). The present study was conducted to identify the most common ecto- and endo-parasitic monogeneans inhabiting commercially important ornamental and/or food fish species that have been imported into Mexico.

## Materials and methods

Ornamental fish species were collected from 2010 to 2014 from different municipalities (Axochiapan, Ayala, Cuautla, Jiutepec, Jojutla, Tlaltizapan, Tlaquiltenango, Xochitepec, and Zacatepec) located in the state of Morelos. Live fish were examined thoroughly externally under a stereo-microscope before opening the visceral cavity.

Fish were sacrificed by puncturing the brain region and the gills of each fish were removed and placed in vials containing hot 4–5% formalin solution to fix any of the ectoparasites that might be present and labeled with data of each collection site. The internal cavity of each fish was exposed by an incision made along the venter from the anus to mouth. The entire alimentary canal was removed; the interior of the gut was thoroughly examined in situ, then placed in a Petri dish containing hot formalin solution 4–5%, where it was searched for monogeneans (Salgado-Maldonado et al. 2014). Subsequently, all monogeneans specimens were isolated and stained with Gomori's trichrome and mounted in Canada balsam. In addition, some specimens were mounted in a mixture of lactic-acid (LA) and glycerin- ammonium picrate (GAP) and then re-mounted in Canada balsam as permanent preparations (Mendoza-Franco et al. 2013). Parasite identifications were made using a Leica microscope DM2500 with Nomarski interference contrast and based on descriptions provided in the following references: García-Vásquez et al. 2007, 2015, Jogunoori et al. 2004, Kritsky et al. 1989, Lim 1996, Mendoza-Palmero et al. 2012, Pariselle and Euzet 1995, Yamaguti 1963. Reference specimens were deposited in the National Helminthological Collection of Mexico (CNHE). Prevalence (percent of hosts infected), mean abundance (mean number of parasites per examined fish), and intensity range for each monogenean species follows Bush et al. (1997). Host species and common names follow those in the FishBase (Froese and Pauly 2017).

## Results

A total of 365 fish specimens of 13 species belonging to 7 families was examined for monogeneans: Cichlidae, Characidae, Cyprinidae, Loricariidae, Osphronemidae, Pangasidae, and Poeciliidae. Twenty-nine monogenean species infecting gills and/or stomachs were identified from hosts species of all families mentioned above from a total of 31 Aquaculture Production Units (APU) from different municipalities located in the state of Morelos (see Table 1 and Figure 1). The prevalence, mean abundance, and mean intensity of infections at each APU of individual species from different hosts are provided in Tables 2–4.


## Discussion

Currently, 31 species of exotic monogeneans have been registered in the state of Morelos due to the introduction of their hosts that are cultured either for food or aquariums (present data; Caspeta-Mandujano et al. 2009). This current study on cultured exotic fish species revealed that cichlids (i.e., species of *Oreochromis*, *Hemichromis*, and *Pterophyllum*), harbored the highest number of monogeneans (14 species) followed by cyprinids with seven species of which *Dactylogyrus baueri* Gussev, 1955, *Dactylogyrus formosus* Kulwiec, 1927, *Dactylogyrus intermedius* Wegener, 1909, and

**Table 1.** Ecto- and endo-parasitic monogeneans (Platyhelminthes) on cultured exotic fish from several Aquaculture Production Units (APU) in the state of Morelos, South-Central Mexico.

Host species/Family	Monogeneans/CNHE	APU	Municipalities
<i>Oreochromis niloticus</i> (Cichlidae)	<i>Cichlidogyrus sclerosus</i> † /10743	Acuícola Jaloxtoc El Cifón 7 Hermanos La cascada Acuícola Ayala Maricultura Argos	Ayala Zacatepec Cuautla Tlaltizapan Ayala Zacatepec
	<i>Cichlidogyrus thurstonae</i> † /10744	La Cascada	Tlaltizapan
	<i>Cichlidogyrus tilapiae</i> † /10745	Acuícola Ayala Maricultura Argos	Ayala Zacatepec
<i>Oreochromis niloticus</i>	* <i>Cichlidogyrus</i> sp. 1 † /10746 * <i>Cichlidogyrus</i> sp. 2 † /10747	Acuícola Ayala	Ayala
<i>Hemichromis</i> sp.	<i>Enterogyrus coronatus</i> ‡ /10748	Maleny	Zacatepec
<i>Oreochromis niloticus</i>	<i>Enterogyrus malmbergi</i> ‡ /10749-10750	Acuícola Ayala Adilene Marisol San Tilapia Acuícola de Jiutepec Pliego	Ayala Ayala Tlaltizapan Juitepec Ayala
		San Tilapia La buena Fortuna	Tlaltizapan Jojutla
<i>Oreochromis niloticus</i>	<i>Gyrodactylus cichlidarum</i> £ /10756 <i>Gyrodactylus yacatlí</i> £ /10757	Acuícola Jaloxtoc Centro Zacatepec	Ayala Zacatepec
		Centro Zacatepec	Zacatepec
<i>Pterophyllum scalare</i>	* <i>Gussevía spiralo-cirra</i> † /10752	Jesús Madariaga	Zacatepec
	* <i>Sciadicleithrum ipthimum</i> † /10753	El Chino	Juitepec
	* <i>Sciadicleithrum</i> sp. † /10754	Olascoaga	Juitepec
<i>Oreochromis niloticus</i>	<i>Scutogyrus longicornis</i> † /10755	La Cascada	Tlaltizapan
<i>Carassius auratus</i> (Cyprinidae)	* <i>Dactylogyrus baueri</i> † /10758	Centro de acopio La Perla	Tlaltizapan
	* <i>Dactylogyrus formosus</i> † /10759	El Invernadero	Ayala
	* <i>Dactylogyrus intermedius</i> † /10760	Los Huajes	Ayala
	<i>Dactylogyrus vastator</i> † /10761-10762	Linda Vista Platanar	Ayala Ayala
	* <i>Gyrodactylus kobayashii</i> £ /10765-10767	Grupo Carsal	Ayala
		Linda Vista Los Huajes Grupo Carsal El Invernadero	Ayala Ayala Ayala Ayala
<i>Cyprinus carpio</i>	* <i>Dactylogyrus extensus</i> † /10763	Ornapez	Ayala
<i>Ctenopharyngodon idella</i>	* <i>Dactylogyrus</i> sp. † /10764	Centro Zacatepec	Zacatepec
<i>Trichogaster trichopterus</i> (Osphronemidae)	* <i>Trianchonatus acleithrium</i> † /10768	Consorcio Lugo-Galeana Granja Acuícola Foras	Jiutepec Axochiapan
	* <i>Trianchonatus trichogasterium</i> † /10769	Consorcio Lugo-Galeana	Jiutepec
<i>Pangasianodon hypophthalmus</i> (Pangasidae)	* <i>Thaparocleidus caecus</i> † /10770	Betta Fish	Xochitepec
	* <i>Thaparocleidus siamensis</i> † /10771-10772	Betta Fish La buena Fortuna	Xochitepec Jojutla
	*Dactylogyridae sp. †	La buena Fortuna	Jojutla
<i>Poecilia reticulata</i> (Poeciliidae)	* <i>Gyrodactylus poeciliae</i> £ /10773	Huertas de Cuatla Exopez Agua Fría	Ayala Tlaltizapan Tlaquiltenengo
<i>Gymnocorymbus ternetzy</i> (Characidae)	* <i>Diaphorocleidus armillatus</i> † /10774-10775	Aquafish Tropipez	Zacatepec
<i>Hypostomus</i> sp. (Loricariidae)	*Gyrodactylidae sp. £ /10777	Consorcio Lugo-Galeana	Jiutepec
	* <i>Unilatus unilatus</i> † /10776	Consorcio Lugo-Galeana	Jiutepec

\* = new record in Mexico. Site of infection on host: † = gills lamellae; ‡ = stomach; £ = fins.



**Figure 1.** Map of the state of Morelos, Mexico showing position of each APU: **1** 7 Hermanos (18°51'49.82132"N; 98°58'01.20211"W) **2** Acuícola Ayala (18°45'11.59525"N; 98°56'58.87989"W) **3** Acuícola de Jiutepec (18°52'29.84116"N; 99°09'24.49751"W) **4** Acuícola Jaloxtoc (18°43'56.72740"N; 98°55'20.14003"W) **5** Adilene Marisol (18°35'43.94208"N; 99°01'43.49419"W) **6** Agua Fría (18°33'22.41096"N; 99°00'57.44948"W) **7** Aquafish (18°38'53.20757"N; 99°13'13.80019"W) **8** Betta Fish (18°46'15.00012"N; 99°12'05.44263"W) **9** Centro Zacatepec (18°39'22.70079"N; 99°12'02.36030"W) **10** Centro de Acopio La Perla (18°38'18.23968"N; 99°00'32.15165"W) **11** Consorcio Lugo-Galeana (18°53'48.34681"N; 99°11'13.92251"W) **12** El Chino (18°54'03.35178"N; 99°12'10.27438"W) **13** El Cifón (18°40'42.68111"N; 99°11'26.16448"W) **14** El Invernadero (18°37'11.86468"N; 98°59'37.85120"W) **15** Exopez (18°41'41.78829"N; 99°06'07.81780"W) **16** Granja Acuicola Foras (18°31'07.09460"N; 98°47'54.39963"W); **17**. Grupo Carsal (18°37'21.23567"N; 99°00'05.49462"W) **18** Huertas de Cuatla (18°45'41.45252"N; 98°54'57.10516"W) **19** Jesús Madariaga (18°39'59.91903"N; 99°12'05.85187"W) **20** La Buena Fortuna (18°38'07.31312"N; 99°10'58.58424"W) **21** La Cascada (18°41'06.91860"N; 99°09'05.97650"W) **22** Linda Vista (18°38'11.27728"N; 98°59'41.36454"W) **23** Los Huajes (18°38'01.06064"N; 98°59'39.86312"W) **24** Maleny (18°39'43.43675"N; 99°11'52.86078"W) **25** Maricultura Argos (18°35'50.18775"N; 99°12'16.44262"W) **26** Olascoaga (18°55'43.39346"N; 99°10'40.92078"W) **27** Ornapez (18°45'06.02177"N; 98°59'14.37030"W) **28** Platanar (18°43'30.25259"N; 98°54'30.22690"W) **29** Pliego (18°37'45.93123"N; 98°59'53.99321"W) **30** San Tilapia (18°39'09.51796"N; 99°11'36.53955"W) **31** Tropipez (18°46'10.83544"N; 99°12'05.47184"W).

**Table 2.** Parameters of infection of monogeneans on cichlids (APU: Aquaculture Production Unit; P%: Prevalence; MA: mean abundance; RI: range of infection; MI: mean intensity; IH: infected hosts).

APU	Hosts	Monogeneans	Inds.	P%	MA	RI	MI	IH
Maleny	<i>Hemichromis</i> sp.	<i>Enterogyrus coronatus</i>	36	50	5.14	1–13	3.6	10/20
7 hermanos	<i>Oreochromis niloticus</i>	<i>Cichlidogyrus sclerosus</i>	12	57	1.71	2–4	3.0	4/7
Acuícola de Jiutepec	<i>Oreochromis niloticus</i>	<i>Enterogyrus malmbergi</i>	18	50	2.57	2–5	3.6	5/10
Acuícola Jaloxtoc	<i>Oreochromis niloticus</i>	<i>Gyrodactylus cichlidarum</i>	18	20	2.57	18	18	1/5
	<i>Oreochromis niloticus</i>	<i>Cichlidogyrus sclerosus</i>	13	100	2.60	1–7	2.6	5/5
Adilene Marisol	<i>Oreochromis niloticus</i>	<i>Enterogyrus malmbergi</i>	53	100	7.57	2–13	5.3	10/10
Centro Zacatepec	<i>Oreochromis niloticus</i>	<i>Gyrodactylus yacatli</i>	15	10	2.14	15	15	1/10
El Cifón	<i>Oreochromis niloticus</i>	<i>Cichlidogyrus sclerosus</i>	7	40	1.00	3–4	3.5	2/5
Acuícola Ayala	<i>Oreochromis niloticus</i>	<i>Cichlidogyrus tilapiae</i>	159	100	22.71	3–37	15.9	10/10
	<i>Oreochromis niloticus</i>	<i>Enterogyrus malmbergi</i>	6	50	0.86	1–2	1.2	5/10
	<i>Oreochromis niloticus</i>	<i>Enterogyrus malmbergi</i>	1	10	0.14	1	1.0	1/10
Pliego	<i>Oreochromis niloticus</i>	<i>Enterogyrus malmbergi</i>	2	25	0.29	2	2.0	1/4
San Tilapia	<i>Oreochromis niloticus</i>	<i>Enterogyrus malmbergi</i>	34	100	4.86	1–17	8.5	4/4
	<i>Oreochromis</i> sp.	<i>Enterogyrus malmbergi</i>	23	60	3.29	1–7	3.83	6/10
La Buena Fortuna	<i>Oreochromis</i> sp.	<i>Enterogyrus malmbergi</i>	76	76.9	10.86	2–19	7.6	10/13
Jesús Madariaga	<i>Pterophyllum scalare</i>	<i>Gussevia spiralicirra</i>	5	10	0.71	5	5.0	1/10
El Chino	<i>Pterophyllum scalare</i>	<i>Sciadicleithrum</i> spp.	6	83.3	1.00	1–2	1.2	5/6
Olascoaga	<i>Pterophyllum scalare</i>	<i>Sciadicleithrum</i> sp.	9	75	1.29	1–4	3.0	3/4

**Table 3.** Parameters of infection of monogeneans on hosts of the Cyprinidae (APU: Aquaculture Production Unit; P%: Prevalence; MA: mean abundance; RI: range of infection; MI: mean intensity; IH: infected hosts).

APU	Hosts	Monogeneans	Inds.	P%	MA	RI	MI	IH
Consorcio Lugo-Galeana	<i>Carassius auratus</i>	<i>Dactylogyrus</i> sp.	520	100	52.0	13–154	86.7	10/10
El invernadero	<i>Carassius auratus</i>	<i>Gyrodactylus kobayashii</i>	525	100	87.5	5–314	87.5	6/6
	<i>Carassius auratus</i>	<i>Dactylogyrus formosus</i>	1	17	0.17	1–8	1.0	1/6
Grupo Carsal	<i>Carassius auratus</i>	<i>Gyrodactylus kobayashii</i>	28	100	20	0.3–54	26.7	3/3
	<i>Carassius auratus</i>	<i>Dactylogyrus vastator</i>	5	33	1.7	5	5.0	1/3
Linda vista	<i>Carassius auratus</i>	<i>Gyrodactylus kobayashii</i>	12	20	1.2	2–10	6	2/10
	<i>Carassius auratus</i>	<i>Dactylogyrus vastator</i>	3	10	0.3	3	3.0	1/10
Los Huajes	<i>Carassius auratus</i>	<i>Dactylogyrus baueri</i>	1	20	0.2	1	1.0	5/5
	<i>Carassius auratus</i>	<i>Dactylogyrus</i> spp.	38	100	7.6	2–25	7.6	5/5
	<i>Carassius auratus</i>	<i>Gyrodactylus kobayashii</i>	102	100	20.4	2–58	20.4	5/5
Centro Zacatepec	<i>Ctenopharyngodon idella</i>	<i>Dactylogyrus</i> sp.	100	14	14.3	100	100.0	1/7
Ornapez	<i>Cyprinus carpio</i>	<i>Dactylogyrus extensus</i>	5	20	0.5	2–3	2.5	2/10

**Table 4.** Parameters of infection of monogeneans on characids, loricariids, osphronemids, pangasids, and poeciliids (APU: Aquaculture Production Unit; P%: Prevalence; MA: mean abundance; RI: range of infection; MI: mean intensity; IH: infected hosts).

APU	Host	Monogeneans	Inds.	P%	MA	RI	MI	IH
Aquafish	<i>Gymnocorymbus ternetzi</i>	<i>Diaphorocleidus armillatus</i>	131	100	13.1	2–24	13.1	10/10
Tropipez	<i>Gymnocorymbus ternetzi</i>	<i>Diaphorocleidus armillatus</i>	698	100	69.8	7–217	69.8	10/10
Consorcio Lugo-Galeana	<i>Hypostomus</i> sp.	<i>Unilatus unilatus</i>	15	60	1.5	1–11	2.5	6/10
	<i>Hypostomus</i> sp.	<i>Gyrodactylus</i> sp.	14	60	1.4	1–8	2.3	6/10
	<i>Trichogaster trichopterus</i>	<i>Trianchoratus</i> spp.	80	75	20	03–54	26.7	3/4
	<i>Trichogaster trichopterus</i>	<i>Trianchoratus trichogasterium</i>	250	80	25	16–61	31.3	8/10
Granja Acuicola Foras	<i>Trichogaster trichopterus</i>	<i>Trianchoratus trichogasterium</i>	564	90	56.4	1–262	62.7	9/10
Betta fish	<i>Pangasianodon hypophthalmus</i>	<i>Thaparocleidus</i> spp.	536	40	26.8	1–125	67.0	8/20
La Buena Fortuna	<i>Pangasianodon hypophthalmus</i>	<i>Thaparocleidus siamensis</i>	1000	100	200	130–300	200.0	5/5
	<i>Pangasianodon hypophthalmus</i>	Dactylogyridae sp.	10400	100	2080	1000–3000	1733.3	5/5
Exopez	<i>Poecilia reticulata</i>	<i>Gyrodactylus poeciliae</i>	4	33	0.67	2	2.0	2/6
Agua fría	<i>Poecilia reticulata</i>	<i>Gyrodactylus poeciliae</i>	75	90	7.5	1–37	8.3	9/10
Huertas de Cuautla	<i>Poecilia reticulata</i>	<i>Gyrodactylus poeciliae</i>	1	12.5	0.125	1	1.0	1/8

*Gyrodactylus kobayashii* Hukuda, 1940 are new geographical records in Mexico (see Tables 1 and 3). Despite the great number of parasitological studies on native and/or introduced species of Cichlidae in Mexico (Vidal-Martínez et al. 2001), studies on the parasite fauna of other exotic freshwater fishes, especially on their monogeneans, are relatively scarce. Exceptionally, there have been many reports of species of *Cichlidogyrus* on species of *Oreochromis* (often called tilapia) (see Kritsky et al. 1994, Jiménez-García et al. 2001). Even so, intensity of infection is comparatively high as well as the number of new records of these monogeneans, the latter which continues to grow each year (see Table 3, Mendoza-Franco et al. 2015b). In the present study, the angelfish *P. scalare* (Schultze) and *Hemichromis* sp. were studied for the first time and are shown to be parasitized with *G. spirallocirra* Kohn & Paperna, 1964, *S. iphthimum* Kritsky, Thatcher & Boeger, 1989, *Sciadicleithrum* sp. (new geographical records), and *E. coronatus* Pariselle, Lambert & Euzet, 1991.

Monogeneans usually exhibit high host specificity in comparison with other parasite groups, parasitizing a single or few closely related host species. The only zoogeographic range expansion of exotic monogeneans on native hosts is the discovery of species of *Cichlidogyrus* and *G. cichlidarum* from tilapia on native cichlids and poeciliids, respectively, in natural environments of Mexico (Jiménez-García et al. 2001, García-Vásquez



et al. 2007, 2017). The present study revealed the highest intensity of infection with *G. cichlidarum* (identified as a tilapia pathogen by García-Vásquez et al. 2017) and *Cichlidogyrus* spp. on *Oreochromis* spp. (see Table 2). Therefore, preventing escape of these tilapia from culture systems due to their monogeneans' ability to infest and persist on other non- or related wild fish is urgently required. Another example of the persistence of monogeneans is seen with the dactylogyrid *Urocleidoides vaginoclastrum* Jogunoori, Kritsky & Venkatanarasaiah 2004. This monogenean was originally described from fishes introduced to India via the aquarium trade. Its type host, the green swordtail *Xiphophorus hellerii* (Heckel) (Poeciliidae), is naturally distributed in southern Mexico and Central America, where the native profundulid *Profundulus labialis* (Günther) also hosts *U. vaginoclastrum*. The problem is that *X. hellerii* has been artificially introduced along with *U. vaginoclastrum* to other hydrological systems such as India and northern Mexico (Jogunoori et al. 2004, Mendoza-Palmero and Aguilar-Aguilar 2008, Mendoza-Franco et al. 2015a) from which other cyprinodontiform hosts could potentially become infected with this parasite. Additionally, in the present study the black tetra *G. ternetzi* (Boulenger) (Characidae) was studied for the first time and is revealed to be highly infested with *D. armillatus* Jogunoori, Kritsky & Venkatanarasaiah, 2004 (Dactylogyridae) (see Table 4). *Gymnocorymbus ternetzi* is native to South America and has been introduced via the aquarium trade to India and Mexico. Currently, there are nine species of *Diaphorocleidus* dispersed on native bryconid and characid (Characiformes) hosts in the neotropics (South and Central America) (Santos et al. 2018). The transfer and/or evidence of extensive cryptic speciation of other monogenean groups from exotic to native or vice versa on closely related hosts in Mexico remains unknown, but that potential exists.

Similarly to the introduced tilapia in Mexico, cyprinids (i.e., *C. idella*) are also widely distributed in the country including habitats located within areas protected for conservation (see Salgado Maldonado et al. 2014). These fishes were introduced to Central America (i.e., Mexico and Honduras) for aquaculture purposes from 1965-1980s (Salgado-Maldonado and Rubio-Godoy 2014, Salgado-Maldonado et al. 2015) and the presence of species of *Dactylogyrus* and *G. kobayashii* (see Table 1, 3) in Morelos might be originally related to these introductions. Poeciliids (known as guppies, mollies, platies, and swordtails) have been studied for ectoparasitic monogeneans in Mexico and mainly gyrodactylids have been reported on the skin and/or gills on these fishes (García-Vásquez et al. 2015). Currently, there are 11 gyrodactylid species described and/or reported from poeciliids. Only species of *Urocleidoides* (Dactylogyridae) have been reported on the gills of the poeciliids of the two-spot livebearers *Pseudoxiphophorus bimaculata* (Heckel), *X. hellerii*, and *Poeciliopsis retropinna* (Regan) from Mexico and Panama (Mendoza-Franco et al. 2015). In the present study, *G. poeciliae* Harris & Cable, 2000 was found for the first time on the guppy *Poecilia reticulata* Peters from Mexico (see Tables 1, 4). This monogenean species has been reported on *Poecilia caucana* (Steindachner) and *P. reticulata* from their natural ranges of distribution (Venezuela and Trinidad, respectively). Among all species of *Gyrodactylus* mentioned above, only *G. bullatarudis* Turnbull, 1956 and *G. turnbulli* Harris, 1986 have been reported on six poeciliid host species (*Gambusia*



*bolbrooki* Girard, *Poecilia sphenops* Valenciennes, *P. reticulata*, *P. bimaculata*, *Poeciliopsis* sp., and *X. hellerii*) from Mexico, Canada, Costa Rica, Peru, Trinidad, Australia, and Singapore (see García-Vásquez et al. 2015). Given the low host specificity of both gyro-dactylid species and the invasive characteristic of poeciliids, the potential transfer of these gyro-dactylids to native poeciliids and other ecologically-associated hosts in Mexico is high (see García-Vásquez et al. 2017, Mendoza-Franco et al. 2015).

The African tilapia (Cichlidae) and the Asian catfish (Pangasiidae) are both freshwater whitefish aquaculture species that potentially compete for similar markets. In fact, in 2013 Mexico was recognized as the second largest importer of pangasius fillet in the world (Martínez et al. 2016). No analysis concerning the environmental impact of the introduction of these latter fishes and their parasites from Vietnam into Mexican aquaculture and/or in wild habitats (Martínez et al. 2016) has been made. *Pangasianodon hypophthalmus* (Sauvage) was studied for the first time in the present study and it revealed to be parasitized with three monogenean species: *Thaparocleidus caecus* (Mizelle & Kritsky, 1969) Lim 1996, *T. siamensis* (Lim 1990) Lim, 1996, and Dactylogyridae sp. (Table 4). Finally, Loricariids, otherwise known as plecos (species of *Hyposomus*) are very popular ornamental freshwater fish naturally found in tropical South America, Panama, and Costa Rica. In Mexico, *Hyposomus plecostomus* (L.) was introduced into the Balsas Basin (see geographic position in Figure 1) to control macrophytes and algae, and are now established in multiple water bodies (Ramírez-Morales and Ayala-Pérez 2009). The only report of a gill monogenean species on an introduced pleco to Mexico is that of *Heteropriapulius* sp. (Dactylogyridae) on the Amazon sail-fin *Pterygoplichthys pardalis* Castelnau from the Reserva de la Biosfera Montes Azules (BRMA) in the state of Chiapas (Mendoza-Franco et al. 2012). The present study provides two new monogenean records for Mexico, Gyrodactylidae sp. and *Unilatus unilatus*, the latter belonging to the Dactylogyridae which was previously reported on the snow pleco *P. anisitsi* Eigenmann and Kennedy and on *Plecostomus* sp., from Brazil and Peru, respectively (Mendoza-Palmero et al. 2012).

The fish examined in the present study are ornamental and/or for food production that are commercialized in Mexico. Results clearly show that importation of these fish can carry several monogeneans, both ecto- and endo-parasitic species, which could infect other related fish in systems they invade. Therefore, determining the occurrence of parasitic species will help provide better aquaculture conditions and will help to solve some of the problems faced by fish farmers. In the literature, there are a number of reports dealing with the introduction of parasites by ornamental fish from which the consequences of parasite introduction can be detrimental to native fish. For example, epizootics that may lead to extensive mortality (i.e., *D. vastator* on cyprinids, see Cone 1999) as shown for several species of monogeneans introduced into farms or aquariums, and from there to natural populations (Bakke et al. 2002, 2007; García-Vásquez et al. 2017). In addition to the identification of invasive host fish species, it is recommended that all freshwater fish imported into the country for food (farmed) or ornamental purposes must comply, at least, with quarantine regulations.

## Acknowledgements

We thank the owners of the Aquaculture Production Units (APU) located in the municipalities of the state of Morelos, Mexico. MTO was supported by a Master student fellowship (scholarship number 301041) from the Consejo Nacional de Ciencia y Tecnología (CONACyT), Mexico. This study was completed during a search and training visit of MTO to EPOMEX from the Universidad Autónoma de Campeche, Mexico; the visit was financially supported by the Fondo para Elevar la Calidad en la Educación Superior (FECES) 2012 (May–June 2014) in Mexico and FECES 2014 (March–April 2015).

## References

- Bakke TA, Harris PD, Cable J (2002) Host specificity dynamics: observations on gyrodactylid monogeneans. *International Journal for Parasitology* 32: 281–308. [https://doi.org/10.1016/S00207519\(01\)00331-9](https://doi.org/10.1016/S00207519(01)00331-9)
- Bakke TA, Cable J, Harris PD (2007) The Biology of Gyrodactylid Monogeneans: The “Russian-Doll Killers”. *Advances in Parasitology* 64: 161–376. [https://doi.org/10.1016/S0065-308X\(06\)64003-7](https://doi.org/10.1016/S0065-308X(06)64003-7)
- Barroso de Magalhães AL, Jacobi CM (2013) Invasion risks posed by ornamental freshwater fish trade to southeastern Brazilian rivers. *Neotropical Ichthyology* 11: 433–441. <http://dx.doi.org/10.1590/S1679-62252013005000003>
- Bush AO, Lafferty KD, Lotz JM, Shostak AW (1997) Parasitology meets ecology on its own terms: Margolis et al. revisited. *Journal of Parasitology* 83: 575–583. <https://doi.org/10.2307/3284227>
- Blackburn TM, Ewen JG (2017) Parasites as drivers and passengers of human-mediated biological invasions. *EcoHealth* 14: S61–S73. <https://doi.org/10.1007/s10393-015-1092-6>
- Caspeta-Mandujano J, Cabañas-Carranza G, Mendoza-Franco EF (2009) Helmintos parásitos de peces dulceacuícolas mexicanos, caso Morelos. AGT, Editor, S.A. Primera Edición, Progreso 202 Planta Alta, Col. Escandón, Mexico, 129 pp.
- Cone DK (1999). Monogenea (Phylum: Platyhelminthes). In: Woo PTK (Ed.) *Fish Diseases and disorders*. CABI Publishing, 289–327.
- Froese R, Pauly D (Eds) (2017) FishBase. <http://www.fishbase.org> [accessed October 2017]
- García-Vásquez A, Hansen H, Shinn AP (2007) A revised description of *Gyrodactylus cichlidarum* Paperna, 1968 (Gyrodactylidae) from the Nile tilapia, *Oreochromis niloticus niloticus* (Cichlidae), and its synonymy with *G. niloticus* Cone, Arthur et Bondad-Reantaso, 1995. *Folia Parasitologica* 54: 129–140. <https://doi.org/10.14411/fp.2007.018>
- García-Vásquez A, Razo-Mendivil U, Rubio-Godoy M (2015) Morphological and molecular description of eight new species of *Gyrodactylus* von Nordmann, 1832 (Platyhelminthes: Monogenea) from poeciliid fishes, collected in their natural distribution range in the Gulf of Mexico slope, Mexico. *Parasitology Research* 114: 3337–3355. <https://doi.org/10.1007/s00436-015-4559-z>
- García-Vásquez A, Razo-Mendivil U, Rubio-Godoy M (2017) Triple trouble? Invasive poeciliid fishes carry the introduced tilapia pathogen *Gyrodactylus cichlidarum* in the Mexican highlands. *Veterinary Parasitology* 235: 37–40. <https://doi.org/10.1016/j.vetpar.2017.01.014>

- IPPC Secretariat (2005) Identification of risks and management of invasive alien species using the IPPC framework. Proceedings of the workshop on invasive alien species and the International Plant Protection Convention, Braunschweig, Germany, 22–26 September 2003. Rome, Italy, FAO, 301 pp.
- Jiménez-García MI, Vidal-Martínez VM, Lopez-Jiménez S (2001) Monogeneans in introduced and native cichlids in Mexico: evidence for transfer. *Journal of Parasitology* 84: 907–909. [https://doi.org/10.1645/0022-3395\(2001\)087\[0907:MIIANC\]2.0.CO;2](https://doi.org/10.1645/0022-3395(2001)087[0907:MIIANC]2.0.CO;2)
- Jogunoori W, Kritsky DC, Venkatanarasaiiah J (2004) Neotropical Monogenoidea. 46. Three new species from the gills of introduced aquarium fishes in India, the proposal of *Heterotylus* n. g. and *Diaphorocleidus* n. g., and the reassignment of some previously described species of *Urocleidoidea* Mizelle & Price, 1964 (Polyonchoinea: Dactylogyridae). *Systematic Parasitology* 58: 115–124. <https://doi.org/10.1023/B:SYPA.0000029422.16712.9a>
- Kritsky DC, Thatcher VE, Boeger WA (1989) Neotropical Monogenea. 15 Dactylogyrids from gills of Brazilian Cichlidae with proposal of *Sciadicleithrum* gen. n. (Dactylogyridae). *Proceedings of the Helminthological Society of Washington* 56: 128–140.
- Kritsky DC, Vidal-Martínez VM, Rodríguez-Canul R (1994) Neotropical Monogenoidea. 19. Dactylogyridae of cichlids (Perciformes) from the Yucatán Peninsula, with descriptions of three new species of *Sciadicleithrum* Kritsky, Thatcher, and Boeger, 1989. *Journal Helminthological Society of Washington* 61: 26–33.
- Lavergne S, Molofsky J (2007) Increased genetic variation and evolutionary potential drive the success of an invasive grass. *PNAS* 104: 3883–3888. <https://doi.org/10.1073/pnas.0607324104>
- Lim LHS (1996) *Thaparocleidus* Jain, 1952, the senior synonym of *Silurodiscoides* Gussev, 1976 (Monogenea: Ancylo-discoidinae). *Systematic Parasitology* 35(3): 207–215. <https://doi.org/10.1007/BF00009640>
- Martínez ED, Malpica SA, Hernández AJ (2010) Estructura de la producción de la piscicultura de ornato del estado de Morelos y su relación con la diversidad de la oferta. *Sociedades rurales, producción y medio ambiente* 10: 16–36.
- Martínez CA, Ramírez HM (2016) Catálogo de peces ornamentales producidos en Morelos con capacidad de ser Especies Exóticas Invasoras (EEI). Elaborado dentro del proyecto GEF 00089333 “Aumentar las capacidades de México para manejar especies exóticas invasoras a través de la implementación de la Estrategia Nacional de Especies Invasoras”. Morelos, México, 170 pp.
- Mendoza-Franco EF, Caspeta-Mandujano JM, Salgado-Maldonado G (2012) Primer reporte de *Heteropriapulius* sp. (Platelmintos, Monogenoidea) infectando al pez diablo *Pterygoplichthys pardalis* (Siluriformes, Loricariidae) introducido en la cuenca del Rio Lacantún de la Reserva de la biosfera montes azules, Chiapas, México. *Jaina* 23: 1–6.
- Mendoza-Franco EF, Caspeta-Mandujano JM, Salgado-Maldonado G (2013) New species of *Cacatuocotyle* (Monogenoidea, Dactylogyridae) parasitising the anus and the gill lamellae of *Astyanax aeneus* (Pisces, Ostariophysi: Characidae) from the Rio Lacantún basin in the Biosphere Reserve of Montes Azules, Chiapas, Mexico. *Parasitology Research* 112: 199–205. <https://doi.org/10.1007/s00436-012-3126-0>
- Mendoza-Franco EF, Caspeta-Mandujano JM, Salgado-Maldonado G, Matamoros WA (2015a) Two new species of *Urocleidoidea* Mizelle et Price, 1964 (Monogenoidea) from the

- gill lamellae of profundulids and poeciliids from Central America and southern Mexico. *Folia Parasitologica* 62: 059. <https://doi.org/10.14411/fp.2015.059>
- Mendoza-Franco EF, Quintal Méndez JR, Laffón Leal SM, del Rio Rodríguez RE (2015b) Ectoparásitos (Platelmintos: Monogenea) exóticos de peces cultivados en zonas aledañas a la Laguna de Términos, municipio del Carmen, Campeche, México” In: Aspectos Socio-Ambientales de la Región de la Laguna de Términos Campeche. Editor-Universidad Autónoma de Campeche, México, 210 pp.
- Mendoza-Palmero CA, Scholz T, Mendoza-Franco EF, Kuchta R (2012) New species and geographical records of dactylogyrids (Monogenea) of catfish (Siluriformes) from the Peruvian Amazonia. *Journal of Parasitology* 98: 484–497. <http://www.journalofparasitology.org/doi/pdf/10.1645/GE-2941.1>
- Mendoza R, Luna S, Aguilera C (2015) Risk assessment of the ornamental fish trade in Mexico: analysis of freshwater species and effectiveness of the FISK (Fish Invasiveness Screening Kit). *Biological Invasions* 17: 3491–3502. <https://doi.org/10.1007/s10530-015-0973-5>
- Pariselle A, Euzet L (1995) *Scutogyrus* gen. n. (Monogenea: Ancyrocephalinae) for *Cichlodygryrus longicornis minus* Dossou, 1982. *C. l. longicornis*, and *C. l. gravivaginus* Paperna and Thurston, 1969, with description of three new species parasitic on African cichlids. *Journal Helminthological Society of Washington* 62: 157–173.
- Ramírez-Morales S, Ayala-Pérez LA (2009) “Plecos” en la presa “Infiernillo”. *Jaina* 20: 24–35.
- Salgado Maldonado G, Caspeta Mandujano JM, Ramírez Martínez C, Lozano Vilano L, García Ramírez ME, Mendoza Franco EF (2014) Helmintos de parásitos de los peces del río Lacantún en la reserva de la Biosfera Montes Azules, Chiapas. Editor: Universidad Autónoma de Nuevo León con Financiamiento de Natura y Ecosistemas Mexicanos, Asociación Civil, Mexico, 147 pp.
- Salgado-Maldonado G, Matamoros WA, Kreiser BR, Caspeta-Mandujano JM, Mendoza-Franco EF (2015) First record of the invasive Asian fish tapeworm, *Bothriocephalus acheilognathi* Yamaguti, 1934, in Honduras Central America. *Parasite* 22: 1–6. <https://doi.org/10.1051/parasite/2015007>
- Salgado-Maldonado G, Rubio-Godoy M (2014) Helmintos parásitos de peces de agua dulce introducidos. In: Mendoza R, Koleff P (Coords) Especies acuáticas invasoras en México. Comisión Nacional para el Conocimiento y Uso de la Biodiversidad, México, 269–285.
- Santos NJE, Costa NGS, Soares JB, Domingues MV (2018) Monogenoidean parasites of *Acestrorhynchus falcatus* (Characiformes: Acestrorhynchidae) from Parána, Brazil: species of *Diaphorocleidus* and *Rhinoxenoides* n. gen. (Monogenoidea: Dactylogyridae). *Journal of Helminthology* 7: 1–12. <https://doi.org/10.1017/S0022149X18000093>
- Tapia Osorio M, Mendoza Franco EF, Caspeta-Mandujano JM (2014) Species exóticas en ecosistemas acuáticos: ¿Invasiones biológicas silenciosas? *Jaina* 25: 25–29.
- Vidal-Martínez VM, Aguirre-Macedo ML, Scholz T, González-Solís D, Mendoza-Franco EF (2001) Atlas of the helminth parasites of cichlid fish of Mexico. Academia, Prague, Czech Republic, 185 pp.
- Yamaguti S (1963) *Systema helminthum*. Volume IV. Interscience Publishers, New York, 699 pp.