

Addition of three new species of Xylariomycetidae fungi on bamboo from Southern China

Xin Zhou^{1,2}, Kamran Habib^{1,2,3}, Wenyu Zeng¹, Yulin Ren¹, Xiangchun Shen^{1,2}, Jichuan Kang⁴, Qirui Li^{1,2}

1 State Key Laboratory of Functions and Applications of Medicinal Plants, Guizhou Medical University, Gui'an, Guizhou, 561113, China

2 The High Efficacy Application of Natural Medicinal Resources Engineering Centre of Guizhou Province (The Key Laboratory of Optimal Utilization of Natural Medicine Resources), School of Pharmaceutical Sciences, Guizhou Medical University, Gui'an, Guizhou, 561113, China

3 Department of Botany, Khushal Khan Khattak University, Karak, KP, 27200 Pakistan

4 Engineering and Research Centre for Southwest Bio-Pharmaceutical Resources of National Education Ministry of China, Guizhou University, Guiyang, Guizhou, 550025, China

Corresponding authors: Qirui Li (lqrnd2008@163.com); Jichuan Kang (jckang@gzu.edu.cn)

Abstract

In our ongoing research on bambusicolous Xylariomycetidae fungi, three new microfungi taxa were collected and identified as members of the genera *Amphibambusa*, *Arecophila*, and *Nigropunctata*. *Amphibambusa aureae* sp. nov., *Arecophila gaofengensis* sp. nov., and *Nigropunctata xiaohensis* sp. nov. are introduced based on morphological comparisons and phylogenetic analyses using combined ITS, LSU, *tub2*, and *tef1a* loci. Comprehensive morphological descriptions, illustrations, and a phylogenetic tree showcasing the placement of these new taxa are provided. Additionally, keys to *Amphibambusa* and *Nigropunctata* are provided.

Key words: Cainiaceae, Guizhou, systematics, Xylariales



Academic editor:

Nalin Wijayawardene

Received: 21 May 2024

Accepted: 3 September 2024

Published: 2 October 2024

Citation: Zhou X, Habib K, Zeng W, Ren Y, Shen X, Kang J, Li Q (2024) Addition of three new species of Xylariomycetidae fungi on bamboo from Southern China. MycoKeys 109: 109–129. <https://doi.org/10.3897/mycokeys.109.128020>

Copyright: © Xin Zhou et al.

This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

Introduction

Bamboo, as the largest member of the grass family Poaceae, plays an important role in local economies worldwide, being distributed across diverse climates, from cold mountainous regions to hot tropical areas. Bamboos exhibit high diversity and are particularly abundant in Asia, notably in China. China boasts plentiful bamboo resources, with its bamboo species constituting of more than 50% of the world's total (Liu et al. 2018). Due to their low natural toxicity, bamboos are susceptible to fungi and insect infestations resulting in abundant microfungi inhabiting their culms and leaves (Dai et al. 2018; Wang et al. 2018; Jiang et al. 2022; Hyde et al. 2023a, b).

Dai et al. (2018) reported an association of more than 1300 fungi with bamboo, consisting of 150 basidiomycetes and 800 ascomycetes species. Among these, 240 and 110 taxa have been reported as hyphomycetous and coelomycetous, respectively. The taxonomic placements of bamboo-associated ascomycetous fungi are highly diverse, comprising more than 1150 species, in 120 families and 400 genera (Dai et al. 2018). Among these families, Xylariaceae

and Hypocreaceae are the most abundant, with 74 (belonging to 18 genera) and 63 (belonging to 14 genera) species, respectively (Dai et al. 2018; Wijayawardene et al. 2022). The genus *Phyllachora* holds the highest number of species occurring on bamboo, followed by *Nectria* and *Hypoxyylon* (Dai et al. 2018). Most bambusicolous ascomycetous taxa in China are known from Taiwan, with 144 species, followed by Hong Kong with 139 species, Yunnan with 133 species, Guangdong with 53 species, Zhejiang with 37 species, Jiangsu with 36 species, and Sichuan with 35 species (Jiang et al. 2022). Jiang et al. (2022) reported 512 bambusicolous ascomycetous taxa from China, associated with 16 bamboo genera. These species are distributed across 50 orders, 116 families, and 279 genera (including 45 genera without any higher rank) and represent more than one-third of the known bambusicolous ascomycetes in the world. Most reported bambusicolous fungi lack detailed morphology or sequence data thus, still require further study (Dai et al. 2018).

During the investigation of bambusicolous Xylariomycetidae fungi, we observed specimens that could not be readily assigned to any known species. To better understand their taxonomic position, we conducted a phylogenetic analysis using a multi-marker approach (internal transcribed spacer ITS, large subunit LSU, β -tubulin *tub2*, and translation elongation factor *tef1a*). Their distinct morphological characteristics distinguish them from the known species. As a result, we propose these specimens as new species.

Materials and methods

Sample collection and morphological study

The specimens were collected during surveys conducted in Guizhou province, and Guangxi Zhuang Autonomous Region in China during 2023. All related habitat information was recorded. The photos of the collected materials were taken using a Canon G15 camera (Canon Corporation, Tokyo, Japan). Materials were placed in paper bags and taken to the lab for morphological characterization and isolation. To preserve the freshness of the specimens, they were dried using a portable fan drier. The dried specimens were carefully labeled and stored. After this preparation, the specimens were ready for both morphological and molecular studies. All specimens were deposited at the Herbarium of Guizhou Medical University (**GMB**) and the Herbarium of Cryptogams, Herbarium of Kunming Institute of Botany, Chinese Academy of Sciences (**KUN-HKAS**), living cultures were deposited at the Guizhou Medical University Culture Collection (**GMBC**).

Morphological characterization and isolation

Macroscopic features (ostiole, clypeus, etc.) of the specimens were examined using an Olympus SZ61 stereomicroscope and photographed using a Canon 700D digital camera. Microscopic morphological features (ascocarps, peridium, paraphyses, asci, ascospore, etc.) were observed using an optical microscope (Nikon Ni) and photographed using a Canon 700D digital cam-

era attached. Melzer's iodine reagent was used to test the apical apparatus structures for amyloid reaction. Ascii and ascospores of the samples were measured using Tarosoft Image Framework (v. 0.9.0.7). Images were polished using Adobe Photoshop CS6 (Adobe Systems, USA). Pure cultures were obtained by single-ascospore isolation (Long et al. 2019) and maintained at 25 °C for 1–5 weeks on PDA (potato dextrose agar) and oatmeal-agar (OA) medium.

DNA extraction, PCR amplification and sequencing

Mycelium was scraped from pure culture plates using a sterilized scalpel and was used for DNA extraction with the methods following the manufacturer's instructions of the BIOMIGA fungus genomic DNA extraction kit. For some specimens where the ascospores did not germinate, we used a method of directly extracting DNA from the contents of the peritheciun. The DNA samples were kept at –20 °C. Internal transcribed spacers (ITS), large subunit LSU, β-tubulin (*tub2*), and translation elongation factor (*tef1a*) were amplified by PCR with primers ITS1/ITS4 (White et al. 1990; Gardes and Bruns 1993), LR0R/LR5 (Vilgalys and Hester 1990), Bt2a/Bt2b (Glass and Donaldson 1995), and EF1-983F/EF1-2218R (Rehner and Buckley 2005), respectively. The components of a 25 µL volume PCR mixture was: 9.5 µL of double distilled water, 12.5 µL of PCR Master Mix, 1 µL of each primer, and 1 µL of template DNA. Qualified PCR products were checked through 1.5% agarose gel electrophoresis stained with Golden View, and were sent to Sangon Co., China, for sequencing (Xie et al. 2020).

Sequence alignments and phylogenetic analyses

All the obtained sequences were deposited in the GenBank (Tables 1, 2). These sequences were compared with each other and all the known sequences in GenBank using the BLASTN algorithm for precise identification. The molecular phylogeny was inferred from a combined dataset of ITS, LSU, *tub2* and *tef1a* sequences. The reference sequences retrieved from open databases originated from recent published literature, and the Blastn results of close matches. Sequences were aligned using the MAFFT v.7.110 online program (Katoh et al. 2019) with the default settings, respectively. The alignment was adjusted manually using BioEdit v.7.0.5.3 (Hall 1999) where necessary. The maximum likelihood (ML) analysis was implemented in RAxML v.8.2.12 using the GTRGAMMA substitution model with 1,000 bootstrap replicates (Stamatakis 2014). The phylogenetic analyses were also performed for Bayesian inference in MrBayes v. 3.2.1 (Ronquist et al. 2012) online. The Markov Chain Monte Carlo (MCMC) sampling in MrBayes v.3.2.2 (Ronquist et al. 2012) was used to determine the posterior probabilities (PP). Six simultaneous Markov chains were run for 1,000,000 generations, and trees were sampled every 1,000th generation. The phylogenetic tree was visualized in FIGTREE v.1.4.4 (Rambaut 2018). All analyses were run on the CIPRES Science Gateway v 3.3 webportal (Miller et al. 2010).

Table 1. Taxa and corresponding GenBank accession numbers of sequences used in the phylogenetic analysis of Fig. 1.

Species	Strain number	GenBank Accession Numbers		References
		ITS	LSU	
<i>Amphibambusa aureae</i>	GMB4550^T	PQ066508	PQ066514	The study
<i>Amphibambusa aureae</i>	GMB4561	PQ066509	PQ066515	The study
<i>Amphibambusa bambusicola</i>	MFLUCC 11-0617 ^T	KP744433	KP744474	Liu et al. (2015)
<i>Amphibambusa hongheensis</i>	KUN-HKAS 112723 ^T	MW892971	MW892969	Jiang et al. (2021a)
<i>Amphibambusa hongheensis</i>	KUNMCC 20-0334 ^T	MW892972	MW892970	Jiang et al. (2021a)
<i>Arecophila australis</i>	GZUCC0124	MT742125	MT742132	Li et al. (2022)
<i>Arecophila australis</i>	GZUCC0112 ^T	MT742126	MT742133	Li et al. (2022)
<i>Arecophila bambusae</i>	HKUCC 4794	NA	AF452038	Kang et al. (1999)
<i>Arecophila clypeata</i>	GZUCC0127	MT742128	MT742135	Li et al. (2022)
<i>Arecophila clypeata</i>	GZUCC0110 ^T	MT742129	MT742136	Li et al. (2022)
<i>Arecophila gaofengensis</i>	GMB4541^T	PQ066512	PQ066516	The study
<i>Arecophila gaofengensis</i>	GMB4559	PQ066513	PQ066517	The study
<i>Arecophila miscanthii</i>	MFLU 19-2333 ^T	NR171235	NG088086	Hyde et al. (2020a)
<i>Arecophila miscanthii</i>	FU31025	MK503821	MK503827	Hyde et al. (2020a)
<i>Arecophila muroiana</i>	GZUCC0122	MT742127	MT742134	Li et al. (2022)
<i>Arecophila zhaotongensis</i>	ZHKU 23-0260	OR995738	OR995745	Han et al. (2024)
<i>Arecophila zhaotongensis</i>	ZHKU 23-0259	OR995735	OR995742	Han et al. (2024)
<i>Arecophila zhaotongensis</i>	GMBCC1145 ^T	OR995740	OR995747	Han et al. (2024)
<i>Arecophila</i> sp.	HKUCC 6487	NA	AF452039	Jeewon et al. (2003)
<i>Arecophila xishuangbannaensis</i>	ZHKU 23-0280	OR995737	OR995744	Han et al. (2024)
<i>Arecophila xishuangbannaensis</i>	GMB-W1283 ^T	OR995736	OR995743	Han et al. (2024)
<i>Atrotorquata lineata</i>	Mt25	AF009807	NA	Kang et al. (1998)
<i>Barrmaelia macrospora</i>	CBS 142768 ^T	NR167684	NA	Jaklitsch et al. (2014)
<i>Barrmaelia rhamnicola</i>	CBS 142772 ^T	NR153497	NA	Voglmayr et al. (2018)
<i>Cainia anthoxanthis</i>	MFLUCC 15-0539 ^T	NR138407	NG070382	Senanayake et al. (2015)
<i>Cainia desmazieri</i>	CAI	KT949896	NA	Jaklitsch et al. (2016)
<i>Cainia globosa</i>	MFLUCC 13-0663 ^T	NR171724	KX822123	Hyde et al. (2016)
<i>Cainia graminis</i>	CBS 136.62	MH858123	MH869701	Vu et al. (2019)
<i>Endocalyx cinctus</i>	JCM 7946	LC228648	LC228704	Delgado et al. (2022)
<i>Endocalyx cinctus</i>	NBRC 31306	MZ313191	MZ313152	Delgado et al. (2022)
<i>Endocalyx grossus</i>	JCM 5164 ^T	MZ313160	MZ313138	Delgado et al. (2022)
<i>Endocalyx grossus</i>	JCM 5165	MZ313159	MZ313158	Delgado et al. (2022)
<i>Endocalyx grossus</i>	JCM 5166	MZ313179	MZ313171	Delgado et al. (2022)
<i>Endocalyx indumentum</i>	JCM 5171 ^T	MZ313153	MZ313161	Delgado et al. (2022)
<i>Endocalyx indumentum</i>	JCM 8042	MZ313162	MZ313157	Delgado et al. (2022)
<i>Endocalyx melanoxanthus</i>	CBS 147393	MW718204	NA	Delgado et al. (2022)
<i>Endocalyx melanoxanthus</i>	CBS 147394	MW718203	NA	Delgado et al. (2022)
<i>Endocalyx metroxylī</i>	MFLUCC 15-0723B	MT929163	MT929314	Konta et al. (2021)
<i>Endocalyx metroxylī</i>	MFLUCC 15-0723A ^T	NR176745	MT929313	Konta et al. (2021)
<i>Endocalyx metroxylī</i>	MFLUCC 15-0723C	NA	MT929315	Konta et al. (2021)
<i>Endocalyx ptychospermatis</i>	ZHKUCC 21 0008 ^T	MZ493352	OK569894	Phukhamsakda et al. (2022)
<i>Endocalyx ptychospermatis</i>	ZHKUCC 21 0009 ^T	MZ493353	OK569895	Phukhamsakda et al. (2022)
<i>Endocalyx ptychospermatis</i>	ZHKUCC 21 0010 ^T	MZ493354	OK569896	Phukhamsakda et al. (2022)
<i>Longiappendispore chromolaenae</i>	MFLUCC 17-1485 ^T	MT214370	MT214464	Mapook et al. (2020)
<i>Requienella fraxini</i>	RS7	KT949911	NA	Jaklitsch et al. (2016)
<i>Requienella fraxini</i>	CBS 140475	NR138415	NA	Jaklitsch et al. (2016)
<i>Requienella seminuda</i>	CBS 140502 ^T	NR154630	MH878683	Jaklitsch et al. (2016)
<i>Seynesia erumpens</i>	SMH 1291	NA	AF279410	Bhattacharya et al. (2000)

Notes: Type specimens are marked with T; "NA": indicates no sequence available in GenBank; newly generated sequences are indicated in bold.

Table 2. Taxa and corresponding GenBank accession numbers of sequences used in the phylogenetic analysis of Fig. 2.

Species	Strain number	GenBank Accession Numbers				References
		ITS	LSU	tub2	tef1α	
<i>Alloanthostomella rubicola</i>	MFLUCC 16-0479	KX533455	KX533456	NA	NA	Daranagama et al. (2016)
<i>Anthostomella obesa</i>	MFLUCC 14-0171	KP297405	KP340546	KP406616	NA	Daranagama et al. (2015)
<i>Melanographium phoenicis</i>	MFLUCC 18-1481 ^T	MN482677	MN482678	NA	MN481518	Hyde et al. (2020b)
<i>Melanographium smilacis</i>	MFLU 21-0075	MZ538514	MZ538548	NA	NA	Boonmee et al. (2021)
<i>Nigropunctata bambusicola</i>	MFLU 19-2134 ^T	MW240662	MW240592	NA	MW759547	Samarakoon et al. (2022)
<i>Nigropunctata bambusicola</i>	MFLU 19-2145 ^T	MW240664	MW240594	NA	MW759548	Samarakoon et al. (2022)
<i>Nigropunctata complanata</i>	HHUF 30674 ^T	LC760560	LC760580	NA	LC760613	Sugita et al. (2024)
<i>Nigropunctata complanata</i>	HHUF 30675 ^T	LC760561	LC760581	NA	LC760614	Sugita et al. (2024)
<i>Nigropunctata complanata</i>	HHUF 30676 ^T	LC760562	LC760582	NA	LC760615	Sugita et al. (2024)
<i>Nigropunctata complanata</i>	HHUF 30677 ^T	LC760563	LC760583	NA	LC760616	Sugita et al. (2024)
<i>Nigropunctata hydei</i>	CMUB 40018 ^T	OR507150	OR507163	NA	NA	Samarakoon et al. (2023)
<i>Nigropunctata hydei</i>	MFLU 23-0410 ^T	OR507151	OR507164	NA	NA	Samarakoon et al. (2023)
<i>Nigropunctata khalidii</i>	GMB1156 ^T	PP153389	NA	PP209114	NA	Li et al. (2024)
<i>Nigropunctata nigrocircularis</i>	MFLU 19-2130 ^T	MW240661	MW240591	MW775612	MW759546	Samarakoon et al. (2022)
<i>Nigropunctata saccata</i>	MFLU 19-2144 ^T	MW240663	MW240593	MW775613	NA	Samarakoon et al. (2023)
<i>Nigropunctata saccata</i>	MFLU 18-0804	MW240658	MW240588	MW775611	NA	Samarakoon et al. (2023)
<i>Nigropunctata thailandica</i>	MFLU 19-2118 ^T	MW240659	MW240589	NA	MW759544	Samarakoon et al. (2022)
<i>Nigropunctata thailandica</i>	HKAS 106975	MW240660	MW240590	NA	MW759545	Samarakoon et al. (2022)
<i>Nigropunctata xiaohensis</i>	GMB4503^T	PQ066510	PQ066518	PQ083530	PQ083532	The study
<i>Nigropunctata xiaohensis</i>	GMB4552	PQ066511	PQ066519	PQ083531	PQ083533	The study
<i>Pseudoanthostomella conorum</i>	CBS 119333	EU552099	EU552099	NA	NA	Daranagama et al. (2016)
<i>Pseudoanthostomella delitescens</i>	MFLUCC 16-0477	KX533451	KX533452	KX789490	NA	Daranagama et al. (2016)
<i>Pseudoanthostomella pini-nigrae</i>	MFLUCC 16-0478 ^T	KX533453	KX533454	NA	NA	Daranagama et al. (2016)
<i>Pseudoanthostomella pini-nigrae</i>	MFLU 18-0877	MW240654	MW240584	MW820918	MW759541	Daranagama et al. (2016)
<i>Pseudoanthostomella pini-nigrae</i>	MFLU 15-3608	MW240655	MW240585	MW820919	MW759542	Daranagama et al. (2016)
<i>Pseudoanthostomella pini-nigrae</i>	HKAS 102309	MW240656	MW240586	MW820920	NA	Daranagama et al. (2016)
<i>Pseudoanthostomella senecionica</i>	MFLUCC 15-0013	MW240674	MW240604	MW820913	MW759554	Daranagama et al. (2016)
<i>Virgaria nigra</i>	CBS 128006	MH864744	MH876180	NA	NA	Vu et al. (2019)

Notes: Type specimens are marked with T; "NA": indicates no sequence available in GenBank; newly generated sequences are indicated in bold.

Results

Phylogeny

Analyses 1: Placements of *Amphibambusa* and *Arecophila*

The aligned data set of phylogram (Fig. 1) comprised 1250 (ITS/LSU) characters, after the exclusion of ambiguously aligned regions and long gaps. *Barrmaelia macrospora* (Nitschke) Rappaz and *B. rhamnicola* Rappaz were chosen as the outgroup taxa. The sequences of our collection *Amphibambusa aureae* formed a clade, exhibiting a firmly established sister relationship with *Amphibambusa bambusicola* D.Q. Dai & K.D. Hyde (94/1.00 ML/BI, Fig. 1). Newly generated sequences from *Arecophila gaofengensis* strains formed a sister branch to those of *A. xishuangbannaensis* L.S. Han & D.Q. Dai with a low support value (69/0.86 ML/BI, Fig. 1). *Amphibambusa aureae* and *Arecophila gaofengensis* are described as two new species.




Figure 1. Molecular phylogenetic analysis of *Amphibambusa aureae*, *Arecophila gaofengensis* and related taxa based on a combined ITS and LSU sequences. Bootstrap support values for maximum likelihood (ML) greater than 75% and Bayesian posterior probabilities (BPP) greater than 0.95 are displayed above or below the respective branches (ML/BP). The newly described species are marked red. Holotype and ex-type materials are in bold.

Analyses 2: Placement of *Nigropunctata*

The aligned dataset of *Nigropunctata* (Fig. 2) comprised 2730 (ITS/LSU/tub2/tef1α) characters, after exclusion of ambiguously aligned regions and long gaps. *Virgaria nigra* (Link) Nees was chosen as the outgroup taxon. In the phylogram (Fig. 2), the sequences of our collection *Nigropunctata xiaohensis* formed a well-supported (100/1.00, ML/BI) distinct clade on the basal of *Nigropunctata*. *Nigropunctata xiaohensis* is described as a new species.




Figure 2. Molecular phylogenetic analysis of *Nigropunctata* and related taxa based on a combined ITS, LSU, *tub2* and *tef1a* sequences. Bootstrap support values for maximum likelihood (ML) greater than 75% and Bayesian posterior probabilities (BPP) greater than 0.95 are displayed above or below the respective branches (ML/BI). The newly described species are marked red. Holotype and ex-type materials are in bold.

Taxonomy

Amphibambusa D.Q. Dai & K.D. Hyde *Fungal Diversity* 72: 9, 2015.
MycoBank No: 550940

Notes. The genus *Amphibambusa* was introduced by Liu et al. (2015) which is characterized by immersed, solitary, scattered, globose to subglobose ascomata, ostiole at the centre, surrounded by white margin, unitunicate, cylindrical, short-pedicellate ascospores with a J+, subapical ring, and fusiform, subhyaline, longitudinally striated, 1-septate ascospores surrounded by a gelatinous

sheath. Currently, the genus comprises two species: *A. hongheensis* H.B. Jiang & Phookamsak and *A. bambusicola* D.Q. Dai & K.D. Hyde (Liu et al. 2015, Jiang et al. 2021a). In this study, we introduce a new species of *Amphibambusa* from Guangxi Zhuang Autonomous Region, China.

***Amphibambusa aureae* X. Zhou, K. Habib & Q. R. Li, sp. nov.**

Mycobank No: 853721

Fig. 3

Etymology. Named after the host-specific epithet “*Phyllostachys aureae* Rivière & C. Rivière” from which the fungus was isolated.

Type. CHINA • Guangxi Zhuang Autonomous Region, Liangfengjiang Forest Park (22°43'24.91"N, 108°26'56.39"E), altitude: 99 m, on *Phyllostachys aureae*, 15 August 2023, Xin Zhou, Wenyu Zeng, 2023LFJ9 (GMB4550, holotype; GMBC4550, ex-type); *ibid* KUN-HKAS 134919, isotype.

Description. Saprobic on dead culms of bamboo, forming black circular spots on the host surface. **Sexual morph:** Ascomata 660–860 µm wide, 520–630 µm high, immersed under host epidermis, solitary, scattered, globose to subglobose, visible as a black dot, ostiole at the center, with a neck, with an underdeveloped clypeus. Ostioles are centrally located, black, surrounded by white margin. Peridium 13–30 µm thick, outer brown to hyaline inner, cells ***textura angularis***. Paraphyses 2–4.8 µm ($\bar{x} = 3.7 \mu\text{m}$, n = 20) wide, longer than the ascii, numerous, filamentous, colorless, branched, septate. Ascii 90–190 × 9–18 µm ($\bar{x} = 148.5 \times 13.1 \mu\text{m}$, n = 20), 8-spored, unitunicate, cylindrical, short-pediculate, apically rounded, with a J+ subapical ring, 1.4–1.9 × 2.5–3.6 µm ($\bar{x} = 1.7 \times 3.1 \mu\text{m}$, n = 6). Ascospores 15–22.5 × 5–7.9 µm ($\bar{x} = 19 \times 6.6 \mu\text{m}$, n = 40), L/W 3.4, 1–2 seriate, fusiform, subhyaline, 1-septate in the middle, slight constricted at the septum, with round ends, with longitudinal striations along the entire length of the ascospore, and enveloped by a gelatinous sheath 2.5–7 µm ($\bar{x} = 5.2 \mu\text{m}$, n = 20), lacking appendage. **Asexual morph:** Undetermined.

Culture characteristics. Cultured on PDA medium at 27 °C for 4–5 weeks, the colony diameter measures 4–4.5 cm, round, slightly raised in the center, with a neat margin. The mycelium at the colony edge is degraded, appearing white and glossy. A portion of the colony center is brown.

Paratype. CHINA • Guangxi Zhuang Autonomous Region, Liangfengjiang Forest Park (22°43'20.90"N, 108°26'33.52"E), altitude: 99 m, on *Phyllostachys aureae*, 15 August 2023, Xin Zhou, Wenyu Zeng, 2023LFJ190 (GMB4561; paratype; GMBC4561, ex-paratype).

Notes. In the phylogram, *Amphibambusa aureae* (ex-type: GMBC4550) clustered in a distinct clade close to *A. bambusicola* D.Q. Dai & K.D. Hyde (ex-type: MFLLUCC 11–0617). The genus *Amphibambusa* is represented by two species, *A. hongheensis* H.B. Jiang & Phookamsak and *A. bambusicola*. *Amphibambusa aureae* shares similarities with both species, such as ascocarps immersed in a black clypeus, ostiolar openings surrounded by a white margin, cylindrical ascii with a J+ subapical ring, and fusiform, longitudinally striated ascospores enveloped by a distinct mucilaginous sheath (Liu et al. 2015, Jiang et al. 2021a). However, *A. aureae* can be distinguished from *A. bambusicola* by its smaller ascospores (15–22.5 × 5–7.9 µm compared




Figure 3. *Amphibambusa aureae* (GMB4550, holotype) **A** type material **B** ascoma immersed under the surface of host **C** cross-section of ascoma **D**, **E** longitudinal sections of ascomata **F** peridium **G** paraphyses **H**, **I** asci **J** a J+ subapical ring bluing in Melzer's reagent **K–M** ascospores **N** culture on PDA. Scale bars: 0.5 mm (**C**, **D**); 100 µm (**E**); 10 µm (**F–M**).

to $25\text{--}27 \times 5.5\text{--}6 \mu\text{m}$ in *A. bambusicola*) (Liu et al. 2015). Additionally, ascospores of *A. aureae* have rounded ends and are slightly constricted at the septum, whereas those of *A. bambusicola* have pointed end cells and are deeply constricted at the septum. *Amphibambusa hongheensis* differs from *A. aureae* by having smaller asci ($118\text{--}160 \times 14\text{--}18 \mu\text{m}$ vs. $90\text{--}190 \times 9\text{--}18 \mu\text{m}$) and larger ascospores ($25.5\text{--}33 \times 5.5\text{--}7.2 \mu\text{m}$ vs. $15\text{--}22.5 \times 5\text{--}7.9 \mu\text{m}$) (Jiang et al. 2021a).

Key to the *Amphibambusa* species

- 1 Ascospore > 22 µm long.....3
- 2 14.7–21.47 µm long ascospore*A. aureae*
- 3 Ascospore 25–27 µm long, with pointed end cells, deeply constricted at the septum.....*A. bambusicola*
- Ascospore 25.5–33 µm long, with round end cells, and slightly constricted at the septum.....*A. hongheensis*

***Arecophila* K.D. Hyde, Nova Hedwigia 63(1–2): 82 (1996)**

Mycobank No: 27653

Notes. The genus *Arecophila* was introduced by Hyde (1996). The genus is characterized by immersed ascomata with blackened clypeus, ostiole at the centre, unitunicate, long-cylindrical asci with a J+, apical ring, and 1-septate ascospores with striations, and covered with a thick mucilaginous sheath (Hyde 1996; Li et al. 2022; Han et al. 2024). In this study, we introduce a new species of *Arecophila* from Guizhou Province, China.

***Arecophila gaofengensis* X. Zhou, K. Habib & Q. R. Li, sp. nov.**

Mycobank No: 853722

Fig. 4

Etymology. The specific epithet “gaofengensis” refers to the geographical location, Gaofeng Village, where the holotype specimen was collected.

Type. CHINA • Guizhou Province, Anshun City, Pingba District, Gaofeng Town, 26°33'96.54"N, 106°54'20.37"E, altitude: 1250 m, on dead culms of bamboo, 30 October 2023, Yulin Ren, 2023GFZ15 (GMB4541, holotype; GMBC4541, ex-type); *ibid* KUN-HKAS 134920, isotype.

Description. Saprobic on the surface of dead bamboo culms, forming black round spots. **Sexual morph:** Ascomata 400–600 µm high, 600–900 µm diam, globose to subglobose, solitary, scattered, sometimes gregarious, immersed beneath blackened clypeus; clypeus well developed, black, coriaceous, ostiole at the center, weakly papillate. Peridium 13–20 µm wide, composed of thick walled, hyaline to brown cells, **texture angularis**. Paraphyses 2–3 µm ($\bar{x} = 2.6$ µm, n = 20) wide, hyaline, numerous, filamentous, branched, septate. Asci 126–210 × 10–13.5 µm ($\bar{x} = 165 \times 12.5$ µm, n = 20), 8-spored, unitunicate, long-cylindrical, short-pedicellate, with a J+, trapezoidal shape apical ring, bluing in Melzer's reagent, 2.2–3.4 µm high, 3.6–4.2 µm diam. Ascospores 19–24.5 × 7–9.5 µm ($\bar{x} = 21.6 \times 7.8$ µm, n = 30), uniseriate, fusiform, brown, 1-septate, septate at the center, slightly constricted septum, tapering at the ends, with longitudinal and sulcate striations, covered with a thick mucilaginous sheath measuring 3–8 µm ($\bar{x} = 6.3$ µm, n = 10). **Asexual morph:** Undetermined.

Culture characteristics. Ascospores germinating on PDA within 36 hours and germ tubes produced from upper cells. Colonies growing fast on PDA, reaching 2 cm in 1 week at 28 °C, effuse, velvety to hairy, nearly circular, irregular at the margin, white from above, pale yellowish white from below. Mycelium




Figure 4. *Arecophila gaofengensis* (GMB4541, holotype) **A, B** ascomata immersed in bamboo host **C** Cross-section of ascoma **D, E** longitudinal sections of ascomata **F** peridium **G** paraphyses **H** a J+ subapical ring staining by Melzer's reagent **I–K** asci with ascospores **L–N** ascospores surrounded by a gelatinous sheath. Scale bars: 0.5 mm (**B–D**); 100 µm (**E**); 10 µm (**F–N**).

immersed in the media, composed of branched, septate, smooth-walled, hyaline, hyphae.

Paratype. CHINA • Guizhou Province, Anshun City, Pingba District, Gaofeng Town, 26°33'95.44"N, 106°54'30.27"E, altitude: 1250 m, on dead culms of bamboo, 30 October 2023, Yulin Ren, 2023GFZ530 (GMB 4559; paratype; GMBC4559, ex-paratype).

Notes. In the phylogram (Fig. 1), *Arecophila gaofengensis* formed a sister branch with *A. xishuangbannaensis* L.S. Han & D.Q. Dai with a low bootstrap values (69/0.86 ML/BI, Fig. 1). *Arecophila gaofengensis* differs from *A. xishuangbannaensis* by its smaller ascospores ($19\text{--}24.5 \times 7\text{--}9.5 \mu\text{m}$ vs. $23\text{--}27 \times 8.5\text{--}9.5 \mu\text{m}$) and smaller asci ($126\text{--}210 \times 10\text{--}13.5 \mu\text{m}$ vs. $180\text{--}270 \times 12\text{--}14 \mu\text{m}$) (Han et al. 2024). The analysis of ITS sequences for these two species reveals a sequence length of 471 base pairs, with a 92.8% similarity, and a 2.1% gap presence, indicating 437 matching positions. Morphologically, the new taxon is close to *A. bambusae*, but can be distinguished from *A. bambusae* by having larger asci ($126\text{--}210 \times 10.3\text{--}13.7 \mu\text{m}$ vs. $132.5\text{--}140 \times 7.5\text{--}8 \mu\text{m}$) and wider ascospores ($19\text{--}24.5 \times 7.1\text{--}9.5 \mu\text{m}$ vs. $19\text{--}22.5 \times 5.5\text{--}7 \mu\text{m}$) (Umali et al. 1999). Morphologically, the new species also resembles *A. muroiana* (I. Hino & Katum.) You Z. Wang et al. However, clypeus is absent in *A. muroiana* (Li et al. 2022), while blackened clypeus was observed in *A. gaofengensis*. So, here we introduced it as a new species of *Arecophila*.

***Nigropunctata* Samarak. & K.D. Hyde, Fungal Diversity 112: 68, 2022.**

MycoBank No: 558737

Notes. The genus *Nigropunctata*, typified by *N. bambusicola* Samarak. & K.D. Hyde, has recently been classified into Xylariales. The genus is characterized by immersed, solitary or scattered ascomata appearing as small black dots, unitunicate, cylindrical asci with a J+, discoid apical ring (Samarakoon et al. 2022). The genus is represented by seven species (<https://www.indexfungorum.org/Names/Names.asp>; Accessed June 21, 2024). In this study, we introduce a new species of *Nigropunctata* from China.

***Nigropunctata xiaohensis* X. Zhou, K. Habib & Q. R. Li, sp. nov.**

MycoBank No: 853723

Fig. 5

Etymology. The specific epithet “*xiaohensis*” refers to the geographical location, Xiaohe Village, where the holotype specimen was collected.

Type. CHINA • Guizhou Province, Guiyang City, Huaxi District, Xiaohe Village, ($25^{\circ}33'10.46''\text{N}$, $105^{\circ}38'22.57''\text{E}$), altitude: 120 m, on bamboo, 1 June 2023, Xin Zhou, Wenyu Zeng, 2023XHC1 (GMB4503, holotype, no culture was obtained); *ibid* KUN-HKAS 134921, isotype.

Description. Saprobic on decaying bamboo culms. **Sexual morph:** Ascomata $320\text{--}380 \times 340\text{--}400 \mu\text{m}$ ($\bar{x} = 352.7 \times 360 \mu\text{m}$, $n = 10$), immersed, solitary or scattered, appearing as small black dots, solitary, in cross-section globose to subglobose with a flattened base. Ostioles centrally, slightly, papillate, black, flush with the surface of the host. Peridium $15\text{--}25 \mu\text{m}$ thick, comprised of several layers, composed of thick-walled, dense, brown to hyaline, cells of ***textura angularis***. Paraphyses $2.8\text{--}4.3 \mu\text{m}$ ($\bar{x} = 3.6 \mu\text{m}$, $n = 20$) wide, longer than the asci, numerous, filamentous, curving, contain white intracellular material. Asci $85.5\text{--}140 \times 11\text{--}18.5 \mu\text{m}$ ($\bar{x} = 120.2 \times 15.5 \mu\text{m}$, $n = 20$) 8-spored, unitunicate, cylindrical, short-pedicellate, apically rounded, with a J+, discoid




Figure 5. *Nigropunctata xiaohensis* (GMB4503, holotype) **A** material **B** ascoma on the surface of host **C** cross-section of ascoma **D**, **E** longitudinal sections of ascomata **F** peridium **G** paraphyses **H–J** asci **K** a wedge-shaped, **J+** apical ring bluing in Melzer's reagent **L**, **M** ascospores **N** ascospores with germ slits. Scale bars: 0.5 mm (**C**, **D**); 100 µm (**E**); 10 µm (**F–N**).

apical ring, measures 1.3–2.4 μm high, 3.5–5.0 μm wide ($\bar{x} = 1.8 \times 4.4 \mu\text{m}$, $n = 10$). Ascospores 11–21 \times 6.5–10.5 μm ($\bar{x} = 17.8 \times 8.1 \mu\text{m}$, $n = 30$), L/W 2.2, uniseriate, unicellular, ellipsoid to broadly ellipsoid, dark brown to black, with rounded ends, covered with a thick mucilaginous sheath measuring 5–8 μm ($\bar{x} = 6.2 \mu\text{m}$, $n = 10$), with a germ slit extending across the entire spore. **Asexual morph:** Undetermined.

Paratype. CHINA • Guizhou Province, Guiyang City Huaxi District, Xiaohe Village (25°33'20.34"N, 105°38'32.23"E), altitude: 120 m, on bamboo, 4 June 2023, Xin Zhou, Wenyu Zeng, 2023XHC340 (GMB4552, paratype).

Notes. In the phylogram (Fig. 2), *Nigropunctata xiaohensis* formed a separate clade in *Nigropunctata* s. str. Morphologically, *N. xiaohensis* resembles *N. complanata* R. Sugita & Kaz. Tanaka (Sugita et al. 2024) as both share similar size ascospore. However, *N. complanata* is distinguished by thick clypeus (75–90 μm high, 270–410 μm diam.), larger asci (130–175 \times 13–20 μm), and an inverted hat-shaped apical ring. The ITS sequences analysis of *N. complanata* and *N. xiaohensis* reveals a sequence length of 496 base pairs, with an 84.3% identity, and 9.1% gap presence. *Nigropunctata nigrocircularis* Samarak. & K.D. Hyde differs in having larger ascomata (450–535 \times 455–560 μm), longer asci (125–170 μm) and smaller ascospore averaging 15.5 \times 6.4 μm with a 3–4.5 μm mucilaginous sheath (Samarakoon et al. 2022). The type species of the genus, *N. bambusicola* Samarak. & K.D. Hyde differs in having smaller ascomata measuring 285–315 \times 260–340 μm , smaller discoid-inverted hat-shaped ascal apical rings (1.7–2 \times 4–4.8 μm), and ascospores measuring 13.5–17 \times 5.5–9.5 μm , with a 2–6 μm mucilaginous sheath (Samarakoon et al. 2022). A recently reported new species from China, *N. khalidii* Y. P. Wu & Q. R. Li, differs by possessing larger ascomata (608–782 \times 762–830 μm vs. 320–380 \times 340–400 μm in *N. xiaohensis*), larger asci (146–173 \times 8.6–13.6 μm vs. 85.5–140 \times 11–18.5 μm in *N. xiaohensis*), and slightly smaller ascospores (14.8–18 \times 6.3–9 μm) lacking a germ slit (Li et al. 2024).

Key to the *Nigropunctata* species

- | | | |
|----|--|----------------------------------|
| 1 | Ascospores lacking a germ slit | 4 |
| 2 | Ascospores with germ slit | 5 |
| 3 | Lacking mucilaginous sheath around ascospores | <i>N. saccata</i> |
| 4a | peridium 11–16 μm wide, ascomata 606–782 \times 762–830 μm | <i>N. khalidii</i> |
| 4b | Peridium 16.5–31 μm wide, ascomata 400–520 \times 485–575 μm | <i>N. hydei</i> |
| 5a | Ascomata > 450 μm diam..... | 6 |
| 5b | Ascomata 260–340 μm diam, asci 95–140 μm long, ascal apical apparatus 1.7–2 \times 4–4.8 μm | <i>N. bambusicola</i> |
| 5c | Ascomata 390–450 μm diam, asci 130–175 μm long, ascal apical apparatus 2.5–3 \times 4.5–5 μm | <i>N. complanata</i> |
| 5d | Ascomata 340–400 μm diam, asci 85.5–140 μm long, ascal apical apparatus 1.3–2.4 \times 3.5–5 μm | <i>N. xiaohensis</i> |
| 6a | Ascomata 450–535 \times 455–560 μm , ascal apical apparatus 3.2–3.6 μm wide | <i>N. nigrocircularis</i> |
| 6b | Ascomata 615–830 \times 770–965 μm , ascal apical apparatus 4.5–6 μm wide | <i>N. thailandica</i> |

Discussion

In this paper, three new species of *Amphibambusa*, *Arecophila*, and *Nigropunctata* associated with bamboo were introduced, which were collected from karst areas of China. Recent studies have expanded our understanding of bambusicolous fungi from southern China. Han et al. (2024) introduced three new species from the family Cainiaceae, including a novel genus *Paramphibambusa* and two new *Arecophila* species. Jiang et al. (2021b) described two new species, *Occultibambusa hongheensis* and *Seriascoma bambusae*, and reported *Occultibambusa kunmingensis* from new habitats. Yu et al. (2023) identified three new species in the Savoryellaceae family and two new records from Sichuan Province. Our discoveries have enriched the research on bambusicolous fungal diversity in southern China.

Amphibambusa has a widespread distribution, reported in both Thailand and China. All known species of *Amphibambusa* have been found exclusively on decorating bamboo, indicating a potential host specificity (Liu et al. 2015; Jiang et al. 2021a). Phylogenetic analysis conducted in this study reveals a close relationship between *Amphibambusa* and *Arecophila*. However, *Amphibambusa* possesses hyaline ascospores pointed at both ends, which distinguishes it from *Arecophila* (Liu et al. 2015). The longitudinal stripes on the surface of *Amphibambusa* ascospores are not easily visible under an optical microscope and can be easily overlooked. Special attention should be paid when observing and describing morphology. Here one new species of *Amphibambusa aureae* was introduced as the third species of the genus.

Currently, there are 20 *Arecophila* epithets in Index Fungorum (<http://www.indexfungorum.org/Names/Names.asp>, July 2024), but only six species and one strain of *Arecophila* sp. have molecular data on Genbank. *Arecophila* clustered into two clades through phylogenetic analysis (Li et al. 2022, Han et al. 2024). Our study also identifies *Arecophila* as comprising two clades. Morphologically, we cannot find a clear difference between these two clades. At the same time, the morphological characteristics of the species in both branches conform to definitions of *Arecophila* (Hyde 1996). This may indicate that *Arecophila* is a polyphyletic group that has undergone convergent evolution. It may also indicate that the genes currently used to construct phylogenetic trees cannot serve as good DNA barcoding for distinguishing *Arecophila* from its approximate genera. In summary, the use of more samples, gene sequences, and morphological features is essential for the future accurate identification of *Arecophila*.

Ascospores are the main identifying feature of ascomycetous fungi (Webster and Weber 2007). Currently, there are eight *Nigropunctata* species published including our new introduction. However, the shapes, dimensions, and colors of the ascospores of all species in the genus *Nigropunctata* are similar with very little variation (Samarakoon et al. 2022, 2023; Li et al. 2024; Sugita et al. 2024). The presence or absence of germ slits and mucilaginous sheaths of ascospores is used as the main basis for distinguishing *Nigropunctata khalidii*, *N. hydei*, *N. saccata* from similar species (Samarakoon et al. 2023; Li et al. 2024). In terms of ascospores' size, the mean value of ascospores of all eight species was 15–18 µm. For example, the ascospores of *N. thailandica* measure 15–18.5 × 7–11.5 µm (mean = 17 × 9 µm, n = 25), while those of *N. complanata* measure 14.5–19.5 × 7.5–10 µm (Samarakoon et al. 2022; Sugita et al. 2024). The averages of the as-

cospore sizes of these two species differed by only 0.1 µm. The ascospore colors of all eight species are brown to dark brown, and the ascospore shapes of all eight species are ellipsoidal (Samarakoon et al. 2022; Samarakoon et al. 2023; Li et al. 2024; Sugita et al. 2024). Except for the ascospores, there are also relatively small morphological differences among the *Nigropunctata* species (Samarakoon et al. 2022; Samarakoon et al. 2023; Li et al. 2024; Sugita et al. 2024). However, there are significant differences in their DNA sequences. Hence, we believe that DNA sequence should be a primary feature for the species identification of *Nigropunctata*.

Acknowledgements

This research was supported by the National Natural Science Foundation of China (32170019 and 31960005); the Guizhou Medical University High-Level Talent Launch Fund Project (2023-058); the Guizhou Provincial Scientific and Technologic Innovation Base (No. [2023]003); the High-level Innovation Talents of Guizhou (No. GCC [2023]048); National Natural Science Foundation of China (12132006); the Guizhou Provincial Natural Science Foundation for High-Level Innovative Talents and Teams (2016-5676, 2015-4021).

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Author contributions

Conceptualization: Jichuan Kang, Qirui Li, Xiangchun Shen. Collection and morphological examinations: Xin Zhou, Wenyu Zen, Yulin Ren. Molecular sequencing, and phylogenetic analyses: Xin Zhou, Kamran Habib. Specimen identification: Xin Zhou, Qirui Li. Original draft preparation: Xin Zhou, Qirui Li. Review and editing, supervision: Xiangchun Shen, Jichuan Kang, Kamran Habib. All authors have read and agreed to the published version of the manuscript.

Author ORCIDs

Kamran Habib  <https://orcid.org/0000-0003-2572-0306>

Yulin Ren  <https://orcid.org/0009-0003-9063-425X>

Jichuan Kang  <https://orcid.org/0000-0002-6294-5793>

Qirui Li  <https://orcid.org/0000-0001-8735-2890>

Data availability

The datasets generated during and/or analyzed during the current study are available in the MycoBank repository (included in the manuscript), and GenBank (included in Tables 1, 2). And also, the datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

References

- Bhattacharya D, Lutzoni F, Reeb V, Simon D, Nason J, Fernandez F (2000) Widespread occurrence of spliceosomal introns in the rDNA genes of ascomycetes. *Molecular Biology and Evolution* 17(12): 1971–1984. <https://doi.org/10.1093/oxfordjournals.molbev.a026298>
- Boonmee S, Wanasinghe DN, Calabon MS, Huanraluek N, Chandrasiri SKU, Jones GEB, Rossi W, Leonardi M, Singh SK, Rana S, Singh PN, Maurya DK, Lagashetti AC, Choudhary D, Dai Y-C, Zhao C-L, Mu Y-H, Yuan H-S, He S-H, Phookamsak R, Jiang H-B, Martín MP, Dueñas M, Telleria MT, Kałucka IL, Jagodziński AM, Liimatainen K, Pereira DS, Phillips AJL, Suwannarach N, Kumla J, Khuna S, Lumyong S, Potter TB, Shivas RG, Sparks AH, Vaghefi N, Abdel-Wahab MA, Abdel-Aziz FA, Li G-J, Lin W-F, Singh U, Bhatt RP, Lee HB, Nguyen TTT, Kirk PM, Dutta AK, Acharya K, Sarma VV, Niranjan M, Rajeshkumar KC, Ashtekar N, Lad S, Wijayawardene NN, Bhat DJ, Xu R-J, Wijesinghe SN, Shen H-W, Luo Z-L, Zhang J-Y, Sysouphanthong P, Thongklang N, Bao D-F, Aluthmuhandiram JVS, Abdollahzadeh J, Javadi A, Dovana F, Usman M, Khalid AN, Dissanayake AJ, Telagathoti A, Probst M, Peintner U, Garrido-Benavent I, Bóna L, Merényi Z, Boros L, Zoltán B, Stielow JB, Jiang N, Tian C-M, Shams E, Dehghanizadeh F, Pordel A, Javan-Nikkhah M, Denchev TT, Denchev CM, Kemler M, Begerow D, Deng C-Y, Harrower E, Bozorov T, Kholmuradova T, Gafforov Y, Abdurazakov A, Xu J-C, Mortimer PE, Ren G-C, Jeewon R, Maharachchikumbura SSN, Phukhamsakda C, Mapook A, Hyde KD (2021) Fungal diversity notes 1387–1511: Taxonomic and phylogenetic contributions on genera and species of fungal taxa. *Fungal Diversity* 111(1): 1–335. <https://doi.org/10.1007/s13225-021-00489-3>
- Dai DQ, Tang LZ, Wang HB (2018) A review of bambusicolous ascomycetes. Bamboo—current and future prospects. <https://doi.org/10.5772/intechopen.76463>
- Daranagama DA, Camporesi E, Tian Q, Liu X, Chamuyang S, Stadler M, Hyde KD (2015) *Anthostomella* is polyphyletic comprising several genera in Xylariaceae. *Fungal Diversity* 73(1): 203–238. <https://doi.org/10.1007/s13225-015-0329-6>
- Daranagama DA, Camporesi E, Jeewon R, Liu X, Stadler M, Lumyong S, Hyde KD (2016) Taxonomic rearrangement of *Anthostomella* (Xylariaceae) based on a multigene phylogeny and morphology. *Cryptogamie. Mycologie* 37(4): 509–538. <https://doi.org/10.7872/crym/v37.iss4.2016.509>
- Delgado G, Miller AN, Hashimoto A, Iida T, Ohkuma M, Okada G (2022) A phylogenetic assessment of *Endocalyx* (Cainiaceae, Xylariales) with *E. grossus* comb. et stat. nov. *Mycological Progress* 21(1): 221–242. <https://doi.org/10.1007/s11557-021-01759-9>
- Gardes M, Bruns TD (1993) ITS primers with enhanced specificity for basidiomycetes—Application to the identification of mycorrhizae and rusts. *Molecular Ecology* 2(2): 113–118. <https://doi.org/10.1111/j.1365-294X.1993.tb00005.x>
- Glass NL, Donaldson GC (1995) Development of primer sets designed for use with the PCR to amplify conserved genes from filamentous ascomycetes. *Applied and Environmental Microbiology* 61(4): 1323–1330. <https://doi.org/10.1128/aem.61.4.1323-1330.1995>
- Hall TA (1999) BioEdit: A user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series* 41: 95–98.
- Han LS, Wijayawardene NN, Liu C, Han LH, Promputtha I, Dai DQ (2024) *Paramphibambusa bambusicola* gen. et. sp. nov., *Arecophila xishuangbannaensis* and *A. zhaotongensis* spp. nov. in Cainiaceae from Yunnan, China. *MycoKeys* 104: 113–132. <https://doi.org/10.3897/mycokeys.104.117872>

- Hyde KD (1996) Fungi from palms. XXIX. *Arecophila* gen. nov. (Amphisphaeriaceae, Ascomycota), with five new species and two new combinations. *Nova Hedwigia* 63: 81–100.
- Hyde KD, Hongsanan S, Jeewon R (2016) Fungal diversity notes 367–490: Taxonomic and phylogenetic contributions to fungal taxa. *Fungal Diversity* 80: 1–270. <https://doi.org/10.1007/s13225-016-0373-x>
- Hyde KD, Norphanphoun C, Maharachchikumbura SSN, Bhat DJ, Jones EBG, Bundhun D, Chen YJ, Bao DF, Boonmee S, Calabon MS, Chaiwan N, Chethana KWT, Dai DQ, Dayarathne MC, Devadatha B, Dissanayake AJ, Dissanayake LS, Doilom M, Dong W, Fan XL, Goonasekara ID, Hongsanan S, Huang SK, Jayawardena RS, Jeewon R, Karunarathna A, Konta S, Kumar V, Lin CG, Liu JK, Liu NG, Luangsa-ard J, Lumyong S, Luo ZL, Marasinghe DS, McKenzie EHC, Niego AGT, Niranjan M, Perera RH, Phukham-sakda C, Rathnayaka AR, Samarakoon MC, Samarakoon SMBC, Sarma VV, Senanayake IC, Shang QJ, Stadler M, Tibpromma S, Wanasinghe DN, Wei DP, Wijayawardene NN, Xiao YP, Yang J, Zeng XY, Zhang SN, Xiang MM (2020a) Refined families of Sordariomycetes. *Mycosphere* 11(1): 305–1059. <https://doi.org/10.5943/mycosphere/11/1/7>
- Hyde KD, Dong Y, Phookamsak R, Jeewon R, Bhat DJ, Jones EBG, Liu N-G, Abeywickrama PD, Mapook A, Wei D, Perera RH, Manawasinghe IS, Pem D, Bundhun D, Karunarathna A, Ekanayaka AH, Bao D-F, Li J, Samarakoon MC, Chaiwan N, Lin C-G, Phutthacharoen K, Zhang S-N, Senanayake IC, Goonasekara ID, Thambugala KM, Phukhamsakda C, Tennakoon DS, Jiang H-B, Yang J, Zeng M, Huanraluek N, Liu J-KJ, Wijesinghe SN, Tian Q, Tibpromma S, Brahmanage RS, Boonmee S, Huang S-K, Thiyagaraja V, Lu Y-Z, Jayawardena RS, Dong W, Yang E-F, Singh SK, Singh SM, Rana S, Lad SS, Anand G, Devadatha B, Niranjan M, Sarma VV, Liimatainen K, Aguirre-Hudson B, Niskanen T, Overall A, Alvarenga RLM, Gibertoni TB, Pfliegler WP, Horváth E, Imre A, Alves AL, da Silva Santos AC, Tiago PV, Bulgakov TS, Wanasinghe DN, Bahkali AH, Doilom M, Elgorban AM, Maharachchikumbura SSN, Rajeshkumar KC, Haelewaters D, Mortimer PE, Zhao Q, Lumyong S, Xu J, Sheng J (2020b) Fungal diversity notes 1151–1276: Taxonomic and phylogenetic contributions on genera and species of fungal taxa. *Fungal Diversity* 100(1): 5–277. <https://doi.org/10.1007/s13225-020-00439-5>
- Hyde KD, Norphanphoun C, Ma J, Yang HD, Zhang JY, Du TY, Gao Y, Gomes de Farias AR, He SC, He YK, Li CJY, Li JY, Liu XF, Lu L, Su HL, Tang X, Tian XG, Wang SY, Wei DP, Xu RF, Xu RJ, Yang YY, Zhang F, Zhang Q, Bahkali AH, Boonmee S, Chethana KWT, Jayawardena RS, Lu YZ, Karunarathna SC, Tibpromma S, Wang Y, Zhao Q (2023a) Mycosphere notes 387–412: Novel species of fungal taxa from around the world. *Mycosphere* 14(1): 663–744. <https://doi.org/10.5943/mycosphere/14/1/8>
- Hyde KD, Amuhenage TB, Apurillo CCS, Asghari R, Aumentado HD, Bahkali AH, Bera I, Bhunjun CS, Calabon MS, Chandrasiri S, Chethana KWT, Doilom M, Dong W, Fallahi M, Gajanayake AJ, Gomdola D, Gunaratne A, Hongsanan S, Huanraluek N, Jayawardena RS, Kapov SA, Khyaju S, Le L, Li CJY, Li QR, Li YX, Lin CG, Linn MM, Liu JK, Liu NG, Luangharn T, Madagammana AD, Manawasinghe IS, Marasinghe DS, McKenzie EHC, Meakala N, Meng QF, Mukhopadhyay S, Norphanphoun C, Pem D, Phukhamsakda C, Sarma VV, Selcuk F, Senanayake IC, Shah S, Shu YX, Silva HVS, Su HL, Tavakol M, Thakshila SAD, Thiyagaraja V, Thongklang N, Tian Q, Tibpromma S, Tun ZL, Ulukapi M, Wang Y, Wannasawang N, Wijayawardene NN, Wimalasena SDMK, Xiao Y, Xiong YR, Yasanthika WAE, Li Q, Dai DQ (2023b) Fungalpedia, an illustrated compendium for the fungi and fungus-like taxa. *Mycosphere : Journal of Fungal Biology* 14(1): 1835–1959. <https://doi.org/10.5943/mycosphere/14/1/22>

- Jaklitsch WM, Fournier J, Rogers JD, Voglmayr H (2014) Phylogenetic and taxonomic revision of *Lopadostoma*. Persoonia. Persoonia 32(1): 52–82. <https://doi.org/10.3767/003158514X679272>
- Jaklitsch WM, Gardiennet A, Voglmayr H (2016) Resolution of morphology-based taxonomic delusions: *Acrocordiella*, *Basiseptospora*, *Blogiascospora*, *Clypeosphaeria*, *Hymenoplectella*, *Lepteutypa*, *Pseudapiospora*, *Requienella*, *Seiridium* and *Strickeria*. Persoonia – Molecular Phylogeny and Evolution of Fungi 37(24): 82–105. <https://doi.org/10.3767/003158516X690475>
- Jeewon R, Liew EC, Hyde KD (2003) Molecular systematics of the Amphisphaeriaceae based on cladistic analyses of partial LSU rDNA gene sequences. Mycological Research 107(Pt 12): 1392–1402. <https://doi.org/10.1017/S095375620300875X>
- Jiang H, Zhang S, Phookamsak R, Promputtha I, Kakumyan P, Xu J (2021a) *Amphibambusa hongheensis* sp. nov., a novel bambusicolous ascomycete from Yunnan, China. Phytotaxa 505(2): 201–212. <https://doi.org/10.11646/phytotaxa.505.2.6>
- Jiang H-B, Phookamsak R, Hyde KD, Mortimer PE, Xu J-C, Kakumyan P, Karunaratna SC, Kumla J (2021b) A taxonomic appraisal of bambusicolous fungi in Occultibambusaceae (Pleosporales, Dothideomycetes) with new collections from Yunnan province, China. Life 11(9): 932. <https://doi.org/10.3390/life11090932>
- Jiang H, Phookamsak R, Hongsanan S, Bhat DJ, Mortimer PE, Suwannarach N, Kakumyan P, Xu J (2022) A review of bambusicolous Ascomycota in China with an emphasis on species richness in southwest China. Studies in Fungi 7(1): 20. <https://doi.org/10.48130/SIF-2022-0020>
- Kang JC, Kong RYC, Hyde K (1998) Studies on the Amphisphaerales 1. Amphisphaeriaceae (sensu stricto) and its phylogenetic relationships inferred from 5.8S rDNA and ITS2 sequences. Fungal Diversity 1: 147–157.
- Kang JC, Hyde KD, Kong RYC (1999) Studies on Amphisphaerales: The Cainiaceae. Mycological Research 103(12): 1621–1627. <https://doi.org/10.1017/S0953756299001264>
- Katoh K, Rozewicki J, Yamada KD (2019) MAFFT online service: Multiple sequence alignment, interactive sequence choice and visualization. Briefings in Bioinformatics 20(4): 1160–1166. <https://doi.org/10.1093/bib/bbx108>
- Konta S, Hyde KD, Eungwanichayapant PD, Karunaratna SC, Samarakoon MC, Xu J, Dauner LAP, Aluthwattha ST, Lumyong S, Tibpromma S (2021) Multigene phylogeny reveals *Haploanthostomella elaeidis* gen. et sp. nov. and familial replacement of *Endocalyx* (Xylariales, Sordariomycetes, Ascomycota). Life 11(486): 1–25. <https://doi.org/10.3390/life11060486>
- Li QR, Zhang X, Lin Y, Samarakoon MC, Hyde KD, She XC, Liao WQ, Karunaratna A, Long SH, Kang YQ, Kang JC (2022) Morpho-molecular characterisation of *Arecophila*, with *A australis* and *A. clypeata* sp. nov. and *A. miscanthi* comb. nov. MycoKeys 88: 123–149. <https://doi.org/10.3897/mycokeys.88.79475>
- Li QR, Habib K, Long SH, Wu YP, Zhang X, Hu HM, Kang JC (2024) Unveiling fungal diversity in China: New species and records within the Xylariaceae. Mycosphere : Journal of Fungal Biology 15(1): 275–364. <https://doi.org/10.5943/mycosphere/15/1/2>
- Liu JK, Hyde KD, Jones EBG, Ariyawansa HA, Bhat DJ, Boonmee S, Maharachchikumbura SSN, McKenzie EHC, Phookamsak R, Phukhamsakda C, Shenoy BD, Abdel-Wahab MA, Buyck B, Chen J, Chethana KWT, Singtripop C, Dai DQ, Dai YC, Daranagama DA, Dissanayake AJ, Doilom M, D'souza MJ, Fan XL, Goonasekara ID, Hirayama K, Hongsanan S, Jayasiri SC, Jayawardena RS, Karunaratna SC, Li WJ, Mapook A, Norphanphoun C, Pang KL, Perera RH, Persoh D, Pinruan U, Senanayake IC, Som-

- rithipol S, Suetrong S, Tanaka K, Thambugala KM, Tian Q, Tibpromma S, Udayanga D, Wijayawardene NN, Wanasinghe D, Wisitrasameewong K, Zeng XY, Abdel-Aziz FA, Adamčík S, Bahkali AH, Boonyuen N, Bulgakov T, Callac P, Chomnunti P, Greiner K, Hashimoto A, Hofstetter V, Kang JC, Lewis D, Li XH, Liu XZ, Liu ZY, Matsumura M, Mortimer PE, Rambold G, Randrianjohany E, Sato G, Sri-Indrasutdi V, Tian CM, Verbeeken A, von Brackel W, Wang Y, Wen TC, Xu JC, Yan JY, Zhao RL, Camporesi E (2015) Fungal diversity notes 1–110: Taxonomic and phylogenetic contributions to fungal species. *Fungal Diversity* 72(1): 1–197. <https://doi.org/10.1007/s13225-015-0324-y>
- Liu W, Hui C, Wang F, Wang M, Liu G (2018) Review of the resources and utilization of bamboo in China. Bamboo-current and future prospects, InTech 133–142. <https://doi.org/10.5772/intechopen.76485>
- Long QD, Liu LL, Zhang X, Wen TC, Kang JC, Hyde KD, Shen XC, Li QR (2019) Contributions to species of Xylariales in China-1. *Durotheca* species. *Mycological Progress* 18(3): 495–510. <https://doi.org/10.1007/s11557-018-1458-6>
- Mapook A, Hyde KD, Mckenzie EHC, Jones GD, Bhat J, Jeewon R, Stadler M, Samarakoon MC, Malaithong M, Tanunchai B, Buscot F, Wubet T, Purahong W (2020) Taxonomic and phylogenetic contributions to fungi associated with the invasive weed *Chromolaena odorata* (Siam weed). *Fungal Diversity* 101(1): 1–175. <https://doi.org/10.1007/s13225-020-00444-8>
- Miller M, Pfeiffer WT, Schwartz T (2010) Creating the CIPRES science gateway for inference of large phylogenetic trees. Proceedings of the Gateway Computing Environments Workshop: 14 Nov 2010. 14: 1–8. <https://doi.org/10.1109/GCE.2010.5676129>
- Phukhamsakda C, Nilsson RH, Bhunjun CS, de Farias ARG, Sun Y-R, Wijesinghe SN, Raza M, Bao D-F, Lu L, Tibpromma S, Dong W, Tennakoon DS, Tian X-G, Xiong Y-R, Karunaratna SC, Cai L, Luo Z-L, Wang Y, Manawasinghe IS, Camporesi E, Kirk PM, Promputtha I, Kuo C-H, Su H-Y, Doilom M, Li Y, Fu Y-P, Hyde KD (2022) The numbers of fungi: Contributions from traditional taxonomic studies and challenges of metabarcoding. *Fungal Diversity* 114(1): 327–386. <https://doi.org/10.1007/s13225-022-00502-3>
- Rambaut A (2018) FigTree—Tree Figure Drawing Tool Version v. 1.4.4. Institute of Evolutionary Biology, University of Edinburgh, Edinburgh.
- Rehner SA, Buckley E (2005) A *Beauveria* phylogeny inferred from nuclear ITS and EF1-alpha sequences: Evidence for cryptic diversification and links to *Cordyceps* telomorphs. *Mycologia* 97(1): 84–98. <https://doi.org/10.3852/mycologia.97.1.84>
- Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) MrBayes 3.2: Efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* 61(3): 539–542. <https://doi.org/10.1093/sysbio/sys029>
- Samarakoon MC, Hyde KD, Maharanachikumbura SSN, Stadler M, Gareth Jones EB, Promputtha I, Suwannarach N, Camporesi E, Bulgakov TS, Liu J-K (2022) Taxonomy, phylogeny, molecular dating and ancestral state reconstruction of Xylariomycetidae (Sordariomycetes). *Fungal Diversity* 112(1): 1–88. <https://doi.org/10.1007/s13225-021-00495-5>
- Samarakoon MC, Lumyong S, Manawasinghe IS, Suwannarach N, Cheewangkoon R (2023) Addition of five novel fungal flora to the Xylariomycetidae (Sordariomycetes, Ascomycota) in Northern Thailand. *Journal of Fungi* 9(11): 1065. <https://doi.org/10.3390/jof9111065>
- Senanayake IC, Maharanachikumbura SSN, Hyde KD, Bhat JD, Jones EBG, McKenzie EHC, Dai DQ, Daranagama DA, Dayarathne MC, Goonasekara ID, Konta S, Li WJ, Shang QJ, Stadler M, Wijayawardene NN, Xiao YP, Norphanphoun C, Li Q, Liu XZ, Bahkali AH, Kang JC, Wang Y, Wen TC, Wendt L, Xu JC, Camporesi E (2015) Towards unraveling

- relationships in Xylariomycetidae (Sordariomycetes). *Fungal Diversity* 73(1): 73–144. <https://doi.org/10.1007/s13225-015-0340-y>
- Stamatakis A (2014) RAxML version 8: A tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 30(9): 1312–1313. <https://doi.org/10.1093/bioinformatics/btu033>
- Sugita R, Yoshioka R, Tanaka K (2024) *Anthostomella*-like fungi on bamboo: Four new genera belonging to a new family Pallidoperidiaceae (Xylariales). *Mycoscience* 65(1): 28–46. <https://doi.org/10.47371/mycosci.2023.11.005>
- Umalí TE, Hyde KD, Quimio TH (1999) *Arecophila bambusae* sp. nov. and *A. coronata* comb. nov., from dead culms of bamboo. *Mycoscience* 40(2): 185–188. <https://doi.org/10.1007/BF02464296>
- Vilgalys R, Hester M (1990) Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. *Journal of Bacteriology* 172(8): 4238–4246. <https://doi.org/10.1128/jb.172.8.4238-4246.1990>
- Voglmayr H, Friebes G, Gardiennet A, Jaklitsch WM (2018) *Barrmaelia* and *Entosordaria* in Barrmaeliaceae (fam. nov., Xylariales) and critical notes on *Anthostomella*-like genera based on multigene phylogenies. *Mycological Progress* 17(1): 155–177. <https://doi.org/10.1007/s11557-017-1329-6>
- Vu D, Groenewald M, de Vries M, Gehrmann T, Stielow B, Eberhardt U, Al-Hatmi A, Groenewald JZ, Cardinali G, Houbraken J, Boekhout T, Crous PW, Robert V, Verkley GJM (2019) Large-scale generation and analysis of filamentous fungal DNA barcodes boosts coverage for kingdom fungi and reveals thresholds for fungal species and higher taxon delimitation. *Studies in Mycology* 92(1): 135–154. <https://doi.org/10.1016/j.simyco.2018.05.001>
- Wang J, Li J, Zhuang X, Pan X, Yu H, Sun F, Song J, Jin C, Jiang Y (2018) Improved mould resistance and antibacterial activity of bamboo coated with ZnO/graphene. *Royal Society Open Science* 5(8): 180173. <https://doi.org/10.1098/rsos.180173>
- Webster J, Weber R (2007) Introduction to fungi. Cambridge university press. <https://doi.org/10.1017/CBO9780511809026>
- White TJ, Bruns T, Lee SJWT, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. PCR protocols: a guide to methods and applications 18(1): 315–322. <https://doi.org/10.1016/B978-0-12-372180-8.50042-1>
- Wijayawardene NN, Hyde KD, Dai DQ, Sánchez-García M, Goto BT, Saxena RK, Erdoğdu M, Selçuk F, Rajeshkumar KC, Aptroot A, Błaszkowski J, Boonyuen N, da Silva GA, de Souza FA, Dong W, Ertz D, Haelewaters D, Jones EBG, Karunaratna SC, Kirk PM, Kukwa M, Kumla J, Leontyev DV, Lumbsch HT, Maharachchikumbura SSN, Marguno F, Martínez-Rodríguez P, Mešić A, Monteiro JS, Oehl F, Pawłowska J, Pem D, Pfliegler WP, Phillips AJL, Pošta A, He MQ, Li JX, Raza M, Sruthi OP, Suetrong S, Suwannarach N, Tedersoo L, Thiagaraja V, Tibpromma S, Tkalčec Z, Tokarev YS, Wanasinghe DN, Wijesundara DSA, Wimalaseana SDMK, Madrid H, Zhang GQ, Gao Y, Sánchez-Castro I, Tang LZ, Stadler M, Yurkov A, Thines M (2022) Outline of Fungi and fungus-like taxa – 2021. *Mycosphere* 13(1): 53–453. <https://doi.org/10.5943/mycosphere/13/1/2>
- Xie X, Liu LL, Shen XC, Kang YQ, Hyde KD, Kang JC (2020) Contributions to species of Xylariales in China-3. *Collodiscula tubulosa* (Xylariaceae). *Phytotaxa* 428(2): 122 Mycoscience 130. <https://doi.org/10.11646/phytotaxa.428.2.6>
- Yu X-D, Zhang S-N, Liu J-K (2023) Additions to Bambusicolous fungi of Savoryellaceae from Southwest China. *Journal of Fungi* 9(5): 571. <https://doi.org/10.3390/jof9050571>