



Review article

Biomarkers of exposure in environment-wide association studies – Opportunities to decode the exposome using human biomonitoring data



Nadine Steckling^{a,b,*}, Alberto Gotti^c, Stephan Bose-O'Reilly^{a,b}, Dimitris Chapizanis^c, Danae Costopoulou^d, Frank De Vocht^e, Mercè Garí^{a,f}, Joan O. Grimalt^f, Ester Heath^g, Rosemary Hiscock^h, Marta Jagodic^g, Spyros P. Karakitsios^c, Kleopatra Kedikoglou^d, Tina Kosjek^g, Leondios Leondiadis^d, Thomas Maggos^d, Darja Mazej^g, Kinga Polańskaⁱ, Andrew Povey^e, Joaquim Rovira^j, Julia Schoierer^a, Marta Schuhmacher^j, Zdravko Špirić^k, Anja Stajniko^g, Rob Stierum^l, Janja Snoj Tratnik^g, Irene Vassiliadou^d, Isabella Annesi-Maesano^m, Milena Horvat^g, Dimosthenis A. Sarigiannis^c

^a University Hospital Munich, WHO Collaborating Centre for Occupational Health, Institute and Outpatient Clinic for Occupational, Social and Environmental Medicine, Unit Global Environmental Health, Ziemssenstr. 1, D-80336 Munich, Germany

^b Department of Public Health and Health Technology Assessment, University for Health Sciences, Medical Computer Science and Technology, Eduard-Wallnöfer-Zentrum 1, A-6060 Hall in Tirol, Austria

^c Aristotle University of Thessaloniki, School of Engineering, Building D, University Campus, GR-54124, Greece

^d National Centre for Scientific Research "Demokritos", Neapoleos 27, 15310 Athens, Greece

^e Centre for Occupational and Environmental Health, Centre for Epidemiology, Division of Population Health, Health Services Research and Primary Care, School of Health

Abbreviations: β -HCH, β -hexachlorocyclohexane; $\mu\text{g/L}$, microgram per liter; $\mu\text{M/L}$, micromolar per liter; Σ , total; 1-HP, 1-hydroxypyrene; 2, 3-DHBA, 2,3-dihydroxybenzoic Acid; 2cx-MMHP, mono-(2-carboxymethylhexyl) phthalate; 3PBA, 3-phenoxybenzoic acid; 4F3PBA, 4-fluoro-3-phenoxybenzoic acid; 5cx-MEPP, mono-(5-carboxy-2-ethylpentyl) phthalate; 5OH-MEHP, mono-(2-ethyl-5-hydroxyhexyl) phthalate; 5oxo-MEHP, Mono-(2-ethyl-5-oxo-hexyl) phthalate; AAMA, N-acetyl-S-(2-carbamoyl-ethyl)-L-cysteine; AAs, alkylating agents; ADI, acceptable daily intake; ALARP, as low as reasonably practicable; AM, arithmetic mean; APGAR, adaptation, partnership, growth, affection, resolve; As, arsenic; AUDIT, alcohol use disorders identification Test; BAC, blood alcohol content; BAT, biological tolerance value; BDCM, bromodichloromethane; BDE 99, 2,2',4,4',5-pentabromodiphenyl ether; BE, biomonitoring equivalents; BEL, biological exposure indices; BFRs, brominated flame retardants; BMD-L, benchmark dose lower confidence limit; BoE, biomarker of exposure; BPA, bisphenol A; BPA-glu, glucuronidated metabolite of BPA; BPAD, biological pathway altering dose; BPF, bisphenol F; BPS, bisphenol S; BPP, butylbenzyl phthalate; Br₂CA, 2,2-dibromovinyl-2,2-dimethylcyclopropanecarboxylic acid; BzBP, benzylbutyl phthalate; CAL REL, California acute reference exposure levels; CC, critical concentration; Cd, cadmium; cis-Cl₂CA, cis-3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropane-1-carboxylic acid; cis-DCCA, 2,2-dichloro-2-dimethylvinyl-cyclopropane carboxylic acid; CYP, cytochrome P450; CYP1A1, cytochrome P450 1A1; CIT, citrinin; CPK, creatine phosphokinase; Cr, chromium; CRP, C-reactive protein; crea, creatinine; Cu, copper; dBA, decibel; DAP, dialkylphosphate; DBCA, 2,2-Dibromo-2-Dimethylvinyl-Cyclo-Propane Carboxylic Acid; DBCM, dibromochloromethane; DBP, di-n-butyl phthalate; DBPs, disinfection by-products; DCCA, 2,2-Dichloro-2-Dimethylvinyl-Cyclopropane Carboxylic Acid; DDE, dichlorodiphenyldichloroethylene; DDT, dichlorodiphenyltrichloroethane; DEDTP, diethyl dithiophosphate; DEHP, di-(2-ethylhexyl) phthalate; DEHT, di-(2-ethylhexyl) terephthalate; DEP, diethyl phthalate; DETP, diethyl thiophosphate; DiBP, di-iso-butyl phthalate; DINCH, diisononyl 1,2-cyclohexanedicarboxylic acid; DiNP, diisononyl phthalate; DMP, dimethyl phosphate; DMDDTP, dimethyl dithiophosphate; DMTP, dimethyl thiophosphate; DNA, deoxyribonucleic acid; DnBP, Di-n-butyl Phthalate; DON, deoxynivalenol; ECO, expired carbon-monoxide; EMF, electromagnetic field; EU's FP7, European Union's 7th Framework Programme; EWAS, environment-wide association studies; FAO, food and agriculture organization; FAS, family affluence scale; Fe, iron; FFO, food frequency questionnaires; GAMA, N-acetyl-S-(2-carbamoyl-2-hydroxyethyl)-L-cysteine; GGT, γ -glutamyl transferase; GM, geometric mean; GPAQ, global physical activity questionnaires; GWAS, genetic-wide association studies; h, hours; HBCDD, hexabromocyclododecane; HBM, human biomonitoring; HCB, hexachlorobenzene; HEALS, health and environment-wide associations based on large population surveys; Hg, mercury; ICC, intraclass correlation coefficient; IL-6, interleukin-6; IMD, index of multiple deprivation; IPAQ, international physical activity questionnaires; JEM, job-exposure-matrix; LDH, lactate dehydrogenase; LOAEL, lowest observed adverse effect level; m7Gua, 7-methylguanine; MAA, 2-methoxy acetic acid; MBP, monobutyl phthalate; MBzP, monobenzyl phthalate; MCT, measure of central tendency; MEHP, mono-(2-ethylhexyl) phthalate; MEHHP, mono-(2-ethyl-5-hydroxyhexyl) phthalate; MEOHP, mono-(2-ethyl-5-oxohexyl) phthalate; MEP, mono-ethyl phthalate; MHA, methylhippuric acid; MiNP, mono-isononyl phthalate; Mn, manganese; mg/kg/day, milligram per kilogram per day; mg/m³, milligram per cubic meter; MnBP, mono-n-butyl phthalate; MOA, mode of action; MRL, minimal risk level; MVOC, microbial volatile organic compounds; n, sample size; NDMA, N-nitrosodimethylamine; NMTCA, N-nitroso-2-methylthiazolidine-4-carboxylic acid; NNAL, 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol; NNK, 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone; No., number; NOAEL, no observed adverse effect level; NOC, N-nitroso compounds; NOx, nitrogen oxides; NPRO, N-nitrosoproline; NPs, nanoparticles; NSAR, N-nitrososarcosine; NTCA, N-nitrosothiazolidine-4-carboxylic acid; O₃, Ozone; OH-MiNP, 7OH-mono-methyloctyl phthalate; OCPs, organochlorine pesticides; OPPs, organophosphate pesticides; OTA, ochratoxin A; oxo-MiNP, 7oxo-mono-methyloctyl phthalate; P₉₀, 90th percentile; PAH, polycyclic aromatic hydrocarbon; Pb, lead; PBB, polybrominated biphenyls; PBBK, physiology-based biokinetic; PBDE, polybromodiphenyl ether; PCB, polychlorinated biphenyl; PCDD, polychlorinated dibenzo-p-dioxins; PCDF, polychlorinated dibenzofurans; PCP, pentachlorophenol; PER, perchlorethylene; PFC, perfluorinated compounds; PFOA, perfluorooctanoic acid; PFOS, perfluorooctanesulfonic acid; pg/ml, pictogram per milliliter; PGA, phenylglyoxylic acid; PK, pharmacokinetic; PM, particulate matter; POD, point of departure; POPs, persistent organic pollutants; PSS, perceived stress scale; PTWI, provisional tolerable weekly intake; PYR, pyrene; RFC, reference concentration; RfD, reference dose; RI, reference interval for clinical guidance; Rn, radon; RV₉₅, reference value; S-PMA, S-phenyl mercapturic acid; SC, stachybotrys chartarum; SD, standard deviation; Se, selenium; SED, systemic exposure dose; SES, socioeconomic status; SG, satratoxin G; SHS, second-hand smoke; STA, state-trait anxiety inventory; TBBPA, Tetrabromobisphenol A; TCAA, trichloroacetic acid; TCEQ ReV, reference value of the Texas commission on environmental quality; TCDD, tetrachlorodibenzo-p-dioxin; TDI, tolerable daily intake; THMs, trihalomethanes; THS, third-hand smoke; TLV, threshold limit values; trans-Cl₂CA, trans-2-dichlorovinyl-2,2-dimethylcyclopropane-1-carboxylic acid; trans-DCCA, 2,2-dichloro-2-dimethylvinyl-cyclopropane carboxylic acid; U/L, units per litre; UFPs, ultrafine particles; UK, United Kingdom; US, United States; UVR, ultraviolet radiation; Zn, Zinc

E-mail address: Nadine.Steckling@med.uni-muenchen.de (N. Steckling).

* Corresponding author.

¹ <http://arbmed.klinikum.uni-muenchen.de>.

<https://doi.org/10.1016/j.envres.2018.02.041>

Sciences, Faculty of Biology, Medicine and Health, University of Manchester, Manchester Academic Health Science Centre, Manchester M13 9BL, United Kingdom

^f Institute of Environmental Assessment and Water Research – Spanish Council for Scientific Research, Barcelona, Spain

^g Jožef Stefan Institute, Department of Environmental Sciences, Jamova cesta 39, 1000 Ljubljana, Slovenia

^h University of Bath, UK Centre for Tobacco and Alcohol Studies, Department for Health Bath BA2 7AY, United Kingdom

ⁱ Nofer Institute of Occupational Medicine, Department of Environmental Epidemiology, 8 Teresy Street, 91-348 Lodz, Poland

^j Universitat Rovira i Virgili, Tarragona, Spain

^k Green Infrastructure Ltd., Fallerovo setaliste 22, HR-10000 Zagreb, Croatia

^l Nederlandse Organisatie voor Toegepast Natuurwetenschappelijk Onderzoek, Zeist, The Netherlands

^m Université Pierre et Marie Curie, Paris, France

ARTICLE INFO

Keywords:

Human biomonitoring
Biomarkers of exposure
Environment-wide association studies
Reference values
Exposure limit values
Biomonitoring equivalents

ABSTRACT

Background: The European Union's 7th Framework Programme (EU's FP7) project HEALS – Health and Environment-wide Associations based on Large Population Surveys – aims a refinement of the methodology to elucidate the human exposome. Human biomonitoring (HBM) provides a valuable tool for understanding the magnitude of human exposure from all pathways and sources. However, availability of specific biomarkers of exposure (BoE) is limited.

Objectives: The objective was to summarize the availability of BoEs for a broad range of environmental stressors and exposure determinants and corresponding reference and exposure limit values and biomonitoring equivalents useful for unraveling the exposome using the framework of environment-wide association studies (EWAS).

Methods: In a face-to-face group discussion, scope, content, and structure of the HEALS deliverable “Guidelines for appropriate BoE selection for EWAS studies” were determined. An expert-driven, distributed, narrative review process involving around 30 individuals of the HEALS consortium made it possible to include extensive information targeted towards the specific characteristics of various environmental stressors and exposure determinants. From the resulting 265 page report, targeted information about BoE, corresponding reference values (e.g., 95th percentile or measures of central tendency), exposure limit values (e.g., the German HBM I and II values) and biomonitoring equivalents (BEs) were summarized and updated.

Results: 64 individual biological, chemical, physical, psychological and social environmental stressors or exposure determinants were included to fulfil the requirements of EWAS. The list of available BoEs is extensive with a number of 135; however, 12 of the stressors and exposure determinants considered do not leave any measurable specific substance in accessible body specimens. Opportunities to estimate the internal exposure stressors not (yet) detectable in human specimens were discussed.

Conclusions: Data about internal exposures are useful to decode the exposome. The paper provides extensive information for EWAS. Information included serves as a guideline – snapshot in time without any claim to comprehensiveness – to interpret HBM data and offers opportunities to collect information about the internal exposure of stressors if no specific BoE is available.

1. Introduction

The European Union's 7th Framework Programme (EU's FP7) project HEALS – Health and Environment-wide Associations based on Large Population Surveys – started in 2013 with a term of 5 years. The objective of HEALS is the refinement of an integrated methodology and the application of analytical and computational tools for elucidating human exposome through the integrated use of advanced statistical tools for environment-wide association studies (EWAS) in support of EU-wide environment and health assessments (www.heals-eu.eu).

Important determinants for the development of diseases are genetic influences and the interaction of environmental stressors (Schwartz and Collins, 2007). Described with the complementary approach of nature and nurture, the term “environment” includes everything that is not genetic (Smith et al., 1999). Consequently, the genome needs to be complemented by the exposome (Wild, 2005, 2012). While the human “genome is fixed at conception” (but changed by mutagenic influences) (Rappaport, 2011), “the exposome encompasses life-course environmental exposures [...], from the prenatal period onwards” (Wild, 2005). Based on the above, genome-wide association studies (GWAS) attempt to describe the influence of genetic factors for the development of diseases (Hirschhorn and Daly, 2005), while EWAS investigate the associations between a wide range of environmental factors and diseases (Patel et al., 2010). In this context, human biomonitoring (HBM) –procedures to determine substances or biological markers in human specimens (Angerer et al., 2007) – provides a valuable tool for understanding the magnitude of exposure from all pathways and sources. A biomarker of exposure (BoE) “may be the identification of an exogenous substance within the system, the interactive product between a

xenobiotic compound and endogenous components, or other event in the biological system related to the exposure”(NRC, 1987). BoEs include either stressors themselves (e.g. the parent compounds), or their metabolites (reaction products), identified in a variety of human specimens such as blood, urine, deciduous teeth or hair (CDC, 2005).

HEALS encompasses a more integrative approach for associating environmental exposures and disease mechanisms and outcomes. Data from the external environment, e.g., measurements of chemicals in different media (e.g. air, water, soil and food), are combined with data regarding internal exposure, e.g., measurements of chemicals in urine or blood, to build the exposome and to derive environment-wide associations between exposure and disease. Starting from HBM samples, quantification of exposure biomarkers, together with identification of markers of effect and susceptibility (mainly-omics), builds the analytical exposure biology framework for unraveling the human exposome using multi-omics technologies according to the HEALS paradigm.

To evaluate HBM data, reference and exposure limit values as well as biomonitoring equivalents are useful and receive particular attention in the HEALS project. Reference values describe the upper level of the populations' background concentration (Angerer et al., 2007; Schulz et al., 2011). The HBM Commission of the German Environment Agency defines the reference value RV₉₅ as “the 95 population percentile [...] rounded off within the 95% confidence interval” of the respective parameter in the matrix obtained from the reference population (Schulz et al., 2011). Reference values contain no information about health-related biological exposure limits (Angerer et al., 2007).

Popular health-related biological exposure limit values are the German HBM I and II values. There is no health risk assumable if the concentration of a substance in urine or blood is below the HBM I level.

A health risk cannot be excluded if the concentration of a substance in urine or blood is between HBM I and HBM II. An increased risk for adverse health effects is given if the concentration is above HBM II (Schulz et al., 2011). Additional exposure limit values are used in the literature. Mocarelli et al. (1986) defined a cut-off limit for pathological results set at “eight times the SD [standard deviation] value above the mean”. Critical concentrations (CC) define the concentration below which the probability of health effects is negligible as was it observed in children at birth (ANSES, 2013). Specific exposure limit values are also mentioned. For example, the copper concentration indicating probable depletion resulting in health effects (Burtis et al., 2012), the early morning cortisol concentration suggesting adrenal insufficiency, and cut-off points which distinguish tobacco use vs. no tobacco use (Kim, 2016) have been determined. The BAT (biological tolerance value) and BEI (biological exposure indices) values are occupational exposure limit values. BAT is the “concentration for a substance [...] in the corresponding biological material at which the health of an employee generally is not adversely affected even when the person is repeatedly exposed during long periods” (DFG, 2016). The BEI is the “level of the determinant most likely to be observed in specimens collected from a worker with an internal dose equivalent to that arising solely from inhalation exposure at the TLV [threshold limit value] concentration”. The TLV represents a safe concentration in air in occupational contexts (Morgan, 1997).

Besides reference values and exposure limit values, biomonitoring equivalents (BEs) are of importance, because they are a first screening method to evaluate potential risk from exposure to environmental stressors using HBM data. BEs are defined as the concentration of a chemical or metabolite in a biological matrix (blood, urine, human milk, etc.), consistent with defined exposure guidance values or toxicity criteria. These include reference doses (RfD) and reference concentrations (RfC), minimal risk levels (MRL) and tolerable daily intakes (TDI), which have been defined using the knowledge available regarding the toxicokinetic properties of the chemical (Boogaard et al., 2011). The application of BEs is based on the assumption that intake and excretion are at equilibrium. This ensures coherence between the guidance values for chronic exposure and the estimated BE (Angerer et al., 2011). Use of reliable physiology-based biokinetic (PBBK) models is the most convenient way to translate external exposure reference values into BEs. Details on the methodology and the specific assumptions for the derivation of BEs for each compound can be found in the references given in Table 4. In general, the main steps for deriving a BE are summarized below:

- (I) The identification of the point of departure (POD) that was used for deriving the external exposure reference value (e.g., TDI or RfD).
- (II) If the POD has been derived from an animal study (which is the most common case), then the respective uncertainty factors that account for interspecies extrapolation and, if needed, the lowest observed adverse effect level (LOAEL) to no observed adverse effect level (NOAEL) extrapolation, are used to calculate the human-equivalent POD.
- (III) By using either a simple pharmacokinetic (PK) or more sophisticated PBBK model, we estimate the expected concentration at the matrix of interest, assuming an intake equal to the human-equivalent POD. For rapidly metabolized compounds, when a urinary metabolite is identified the daily urinary excretion of the compound normalized by average urine volume and average creatinine excretion at the daily exposure rate equal to the human-equivalent POD has to be estimated. For this we have to make an assumption on the percentage of intake that is eliminated via the urinary tract. In both cases, the result of the toxicokinetic calculation helps us to derive the biological matrix-related BE_{POD}.
- (IV) Finally, to end up with a BE value that is relevant to humans, uncertainty factors related to intraspecies differences have to be

applied on the BE_{POD}. When a detailed PBTK model is available, intraspecies variability can be directly incorporated in the relevant anthropometric (i.e. bodyweight, body mass index) and biochemical (e.g. metabolic rates based on the genetic polymorphisms of the cytochrome P450 [CYP] isozymes) parameters.

For non-persistent compounds, such as phthalates and bisphenol A, BEs refer usually to levels of metabolite(s) measured in urine; for persistent compounds the biological matrix of reference is either milk (e.g. for POPs) or blood (e.g. heavy metals like Cd and Pb).

In the framework of HEALS, BoEs of a large number of environmental stressors were reviewed and used for supporting environment-wide associations. The main objective of this work was to summarize the availability of BoEs for the broad range of environmental stressors and exposure determinants of interest in HEALS (including heavy metals, persistent and non-persistent organic compounds, particulate matter and biologicals) and corresponding reference and exposure limit values and biomonitoring equivalents useful for unraveling the exposure using the EWAS framework. Additionally, environmental stressors and exposure determinants without known BoEs were discussed.

2. Methodology

This review was based on an expert panel discussion to determine scope, content, and structure of the HEALS guidelines for appropriate BoE selection for EWAS studies. An extensive list of the most important environmental stressor categories as well as selected stressors relevant to human health of the population in the EU was created based on expert opinion. An expert-driven, distributed, narrative review process involving around 30 scientists of the HEALS consortium made it possible to include extensive information targeted towards the specific characteristics of the individual stressor. A narrative/qualitative review design was preferred in contrast to a systematic one, because the intention was to give a broad comprehensive overview of the great number of topics included (Callcut and Branson, 2009; Cook et al., 1997).

The review process was organized on the basis of stressor-specific fact sheets. Every author summarized the latest information about chemical properties, effects on biological systems, exposure routes, absorption, elimination, specimens for analysis, and eventually reference and exposure limit values for at least one (mostly more than one) fact sheet(s). There was no common systematic strategy for literature searches because of the diversity of topics. However, an internal review process (see below) reduced possible researcher bias during the literature search. While most fact sheets were created for specific environmental stressors (e.g., mercury), in some cases it was necessary to summarize a group of stressors in one fact sheet (e.g., psychological occupational hazards). This was an essential, yet feasible approach in some cases, so as to represent a wide range of stressors important to determine the exposome of the EU population.

Information was obtained from comprehensive reports of international organizations (e.g., WHO's Environmental Health Criteria) and other mainstream scientific literature supplemented by the latest research results published in PubMed listed journal papers. Overall, more than 800 references were reviewed.

For quality assurance, all contributors were involved in an internal review process. Each fact sheet was reviewed by at least two project partners, while one of them was the project coordinator, co-coordinator, or leader of the HEALS HBM work package. The leading question for the review process was: “Is the quality, content, and extent of the fact sheet as well as the literature selection suitable and is the information included up to date?” The literature review process described above resulted in a dedicated technical report available for download on the HEALS website: http://www.heals-eu.eu/wp-content/uploads/2013/08/HEALS_D4.2.pdf. A concise selection of information was extracted, updated, and key conclusions are summarized in this

Table 1

Summarizing table, comprising a list of stressor categories, individual stressors and biomarkers of exposure considered and availability (✓: available, X: not available) of reference values (R), exposure limit values (E) and biomonitoring equivalents (BE).

Stressor categories		Individual stressors		Biomarker of Exposure		Availability		
No.		No.	(alphabetically by stressor category)	No.	(alphabetically by individual stressor)	R	E	BE
1	persistent organic pollutants (POPs)	1	BFRs	1	BDE-99	✓	X	✓
				2	HBCDD	X	X	✓
				3	PBDE	✓	X	X
		2	dioxins and furans	4	dioxin-like compounds	X	X	X
				5	GGT	✓	✓	X
				6	PCDD	✓	X	X
				7	PCDF	✓	X	X
				8	TCDD	X	X	X
				9	bile acids steroids	X	X	X
		3	OCPs	10	β-HCH	✓	X	X
				11	DDE	✓	X	✓
				12	DDT	✓	X	✓
				13	HCB	✓	X	✓
		4	PCBs	14	ΣPCB	✓	✓	X
				15	dioxin-like PCBs	✓	X	X
				16	indicator PCBs	✓	X	X
				17	PCB 28	✓	X	X
				18	PCB 52	✓	X	X
				19	PCB 101	✓	X	X
		5	PFC	20	PFOA	✓	✓	X
				21	PFOS	✓	✓	X
2	other organic contaminants	6	bisphenols	22	BPA	✓	✓	X
				23	BPA-glu	X	X	✓
		7	OPPs	24	DAP	✓	X	X
				25	DETTP	✓	X	X
				26	diethyl phosphate	✓	X	X
				27	DMDTP	✓	X	X
				28	DMP	✓	X	X
				29	DMTP	✓	X	X
		8	PAHs	30	1-hydroxypyrene	✓	X	X
				31	fluoranthene	✓	X	X
				32	fluorene	✓	X	X
				33	3-hydroxy-fluorene	✓	X	X
				34	1-naphthol	✓	X	X
				35	2-naphthol	✓	X	X
				36	1-hydroxy-phenanthrene	✓	X	X
				37	2-hydroxy-phenanthrene	✓	X	X
				38	3-hydroxy-phenanthrene	✓	X	X
		9	parabens	39	butyl parabens	✓	X	X
				40	ethyl parabens	✓	X	X
				41	methyl parabens	✓	X	X
				42	propyl parabens	✓	X	X
		10	phthalates	43	BzBP	X	X	✓
				44	DBP	X	X	✓
				45	DEHP	X	X	✓
				46	5cx-MEPP	X	X	✓
				47	MEHHP	X	X	✓
				48	MEHP	✓	✓	✓
				49	5oxo-MEHP	✓	✓	✓
				50	MEOHP	X	X	✓
				51	MEP	X	X	✓
				52	DiBP	X	X	X
				53	DiNP	X	X	✓
				54	oxidative metabolites	X	X	✓
				55	carboxy-MiNP	X	X	✓
				56	OH-MiNP	X	X	✓
				57	oxo-MiNP	X	X	✓
				58	DnBP	X	X	X
					MnBP	X	X	X

(continued on next page)

Table 1 (continued)

Stressor categories		Individual stressors		Biomarker of Exposure			Availability						
No.		No.	(alphabetically by stressor category)	No.	(alphabetically by individual stressor)		R	E	BE				
3	toxic and potential toxic elements	11	PYR	59	ΣPYR		✓	X	X				
				60	3PBA		✓	X	X				
				61	cyfluthrin	4F3PBA		X	X	✓			
				62		cis-Cl ₂ CA		✓	X	X			
				63		cis-DCCA		X	X	✓			
				64		DCCA		X	X	✓			
				65		trans-Cl ₂ CA		✓	X	X			
				66		trans-DCCA		X	X	✓			
				63*	cypermethrin	cis-DCCA		X	X	✓			
				66*		trans-DCCA		X	X	✓			
				67	deltamethrin	Br ₂ CA		✓	X	X			
				62*		cis-Cl ₂ CA		✓	X	X			
				68		DBCA		X	X	✓			
				65*		trans-Cl ₂ CA		✓	X	X			
				63*	permethrin	cis-DCCA		X	X	✓			
				66*		trans-DCCA		X	X	✓			
				12	As	69	As		✓	✓	X		
		70	dimethylated As				X	X	✓				
		71	inorganic As				X	X	✓				
		72	monomethylated As				X	X	✓				
		73	Cd				✓	✓	✓				
		74	Cr				✓	X	X				
		75	Cu				✓	✓	X				
		76	Fe				X	X	X				
		77	Hg				✓	✓	X				
		78	Mn				✓	X	X				
		79	Pb				✓	X	X				
		80	Se				✓	X	X				
		81	Zn				✓	✓	X				
		4	volatile organic compounds (VOCs)			22	acrylamide	82	AAMA		✓	X	X
								83	GAMA		✓	X	X
				84	benzene				✓	X	✓		
85	S-PMA				✓			X	X				
24	cyanide			86	2-Aminothiazoline-4-carboxylic acid		✓	X	X				
				25	ethylbenzene	87	ethylbenzene		✓	X	✓		
						88	PGA		✓	X	X		
						26	glycol ethers	89	MAA		✓	✓	X
27	PCP							90	PCP		✓	✓	X
				28	PER			91	PER		✓	X	X
								29	styrene	92	mandelic acid		✓
						93	N-Acetyl-S-(phenyl-2-hydroxyethyl)-L-cysteine				✓	X	X
88*	PGA						✓			X	X		
94	styrene				✓	X	✓						
30	toluene			95	hippuric acid		✓	X	X				
				96	N-Acetyl-S-(benzyl)-L-cysteine		✓	X	X				
				97	toluene		✓	X	✓				
				98	triclosan		✓	X	✓				
31	triclosan			99	2-MHA		✓	X	X				
				100	3-MHA		✓	X	X				
				101	4-MHA		✓	X	X				
				102	<i>m</i> , <i>p</i> -xylene		✓	X	X				
103	MHA			104	<i>o</i> -, <i>m</i> -, <i>p</i> -xylene		✓	X	X				
				105	<i>o</i> -xylene		✓	X	✓				
				106	xylene		✓	X	X				
				→	see substance of interest		→	→	→				
34	chemotherapy			→	see substance of interest		→	→	→				
				X	no BoE available		X	X	X				
36	tobacco smoke			107	nicotine	cotinine	✓	✓	X				
				37	bioaerosols	108	mold	SC	SG	X	X	X	
109	MVOC							X	X	X			
110	mycotoxins							X	X	X			
111	1-HP					✓	X	X					
112	NO _x					✓	X	X					
X	no BoE available					X	X	X					
113	2,3-DHBA					X	X	X					
X	no BoE available					X	X	X					
X	no BoE available					X	X	X					
8	food contamination	44	biological agents			114	mycotoxins	CIT	✓	X	X		
						115		DON	DON15GlcA	✓	X	X	
						116		OTA		✓	X	X	
						→	see substance of interest		→	→	→		
						→	see substance of interest		→	→	→		
						→	see substance of interest		→	→	→		

(continued on next page)

Table 1 (continued)

Stressor categories		Individual stressors		Biomarker of Exposure		Availability		
No.		No.	(alphabetically by stressor category)	No.	(alphabetically by individual stressor)	R	E	BE
9	water contamination	46	DBPs	117	TCAA	✓	X	X
		47	THMs	118	BDCM	✓	X	✓
				119	bromoform	✓	X	✓
				120	chloroform	✓	X	✓
				121	DBCM	✓	X	✓
10	noise	48	noise	X	no BoE available	X	X	X
11	DNA-damaging agents	49	AAs	122	m ⁷ Gua	✓	X	X
				123	nitrosamines	✓	X	X
				124	Enitrosamines NSAR	X	X	X
				125		✓	X	X
				126	NNK	✓	X	X
				127	NOC	✓	X	X
		50	EMF	X	no BoE available	X	X	X
		51	Rn	X	no BoE available	X	X	X
		52	UVR	128	thymine dimers	✓	X	X
		53	biological	→	see substance of interest	→	→	→
		54	chemical	→	see substance of interest	→	→	→
		55	mechanical	X	no BoE available	X	X	X
		56	physical	X	no BoE available	X	X	X
		57	psychological	X	no BoE available	X	X	X
		58	alcohol consumption	129	ethanol	X	X	X
		59	consumer products	X	no BoE available	X	X	X
		60	drug consumption	→	see substance of interest	→	→	→
		61	nutritional status	→	see substance of interest	→	→	→
12	occupational hazards			130	folate	✓	X	X
				131	vitamin C	✓	X	X
				132	ammonia	✓	X	X
				133	creatinine	✓	X	X
				134	lactate	✓	X	X
		62	physical activity	X	no BoE available	X	X	X
				135	cortisol	✓	✓	X
		63	SES					
		64	stress					

Abbreviations: ✓, available; X, not available; →, see substance of interest. * same BoE for more than one stressor; No., number (used to count the number of stressor categories, individual stressors, and biomarkers included in this manuscript), R, reference values; E, exposure limit values; BE, biomonitoring equivalent. Abbreviations of stressor groups and biomarkers are explained in the list of abbreviations at the end of the manuscript.

paper. The paper focuses on the availability of BoE in body fluids (blood/serum/plasma, breast milk, urine) as well as hair. Presented are reference values, exposure limit values and biomonitoring equivalents (BEs). If available, the reference value (RV₉₅) as defined by the Human Biomonitoring Commission of the German Environment Agency (Schulz et al., 2007) on the basis of a guideline from the International Union of Pure and Applied Chemistry (IUPAC) (Poulsen et al., 1997) is presented. If not available, the 95th percentile (P₉₅) was included as reference value. Otherwise, the third choice was the 90th percentile and the fourth choice was (the range of) measures of central tendency (MCT) like mean or median presented in combination with the maximum value, if available. Condensed values for a population (distinguished in children and adults) were preferred (e.g., P₉₅ for adults aged 18–69 years) instead of values separated by subgroup (e.g., P₉₅ for 18–19 years old, P₉₅ for 20–29 years old, etc.). If a condensed value is not given in the original publication, the range of youngest to oldest is presented in this paper. Values based on the general population are preferred instead of subgroups with special exposures (e.g. like smokers, people with amalgam fillings or high fish consumption). Latest values are presented. Non-creatinine-corrected values are preferred, if available. For reference values, the main – but not exclusive – focus lay on populations in the EU.

The first choice of exposure limit values was the German HBM values (HBM I and II). Otherwise, critical concentrations, cut-offs or other values are included. Some examples of occupational exposure limit values (e.g., BAT) were included. Completeness was not intended. Stressors without measurable BoE are explicitly discussed. All content was updated to at least January 2017 or later as appropriate.

3. Results

A total of 64 chemical, biological, physical, social, or psychological stressors organized in 13 broad stressor categories were selected (Table 1) to fulfil the requirements of EWAS, although the BoEs for some exposure determinants/modifiers (e.g., socioeconomic status) were not expected to be available. In total, information of 135 BoE is summarized. If available, reference values (Table 2), exposure limit values (Table 3), and biomonitoring equivalents (Table 4) are presented. From the complete list of individual stressors (Table 1), 12 were identified without a BoE. These stressors (and some summarized groups of stressors like psychological occupational hazards) are included in Table 5 to discuss opportunities other than HBM to collect information about their internal exposure.

Table 1 includes the stressor categories and stressors with available BoEs as well as – if available – an incomplete selection of corresponding reference values. Reference values were found for 104 of the 135 considered BoEs. Table 3 contains exposure limit values and Table 4 BEs by stressor, when available. Exposure limit values are available for 16 of the 130 considered BoEs. BEs are available for not more than 42 of the 130 BoEs considered.

4. Discussion

Specific BoEs are available for several environmental stressors but not for others. While chemicals and their primary metabolites may be measurable in human specimens, it is not possible at this time to identify BoEs for stressors such as electromagnetic fields or for exposure determinants/modifiers such as socioeconomic status using

Table 2
Biomarkers of exposure and reference values.

Stressor group Stressor	Biomarker of exposure	Matrix	Reference value P ₉₅ or measure of central tendency	Subgroup (years of age; n: sample size), country	Survey year	Reference
POPs	BFRs	serum	MCT (median): 2.1–15.4 ng/g lipid	Population (age not specified; n: 1667), several EU countries (Belgium, France, Greece, Norway, Spain, Sweden, United Kingdom)	1994–2004	(Gari and Grimalt, 2013)
			P ₉₅ : 34.6 ng/g lipid	Adults (18–74 years; n: 731), Catalonia/Spain	2002	(Gari and Grimalt, 2013)
	BDE-99	serum	MCT (median): 0.08–2.4 ng/g lipid	Population (age not specified; n: 1667), several EU countries (Belgium, France, Greece, Norway, Spain, Sweden, United Kingdom)	1994–2004	(Gari and Grimalt, 2013)
			P ₉₅ : 5.2 ng/g lipid	Adults (18–74 years; n: 731), Catalonia/Spain	2002	(Gari and Grimalt, 2013)
dioxins and furans	dioxin-like compounds CYP1A1	peripheral blood lymphocytes	/	/	/	(Päpke et al., 2011; Saurat et al., 2012; Van Duursen et al., 2010)
			Reference limit ([#]): 4–27 U/L	Children (6–10 years; n: about 1000), Italy	1976–1982	(Mocarelli et al., 1986)
	GGT PCDD and PCDF	serum breast milk	MCT (mean): 3.3–22.3 pg/g fat	Subgroup not specified (age not specified; sample size not specified), 27 countries (among others: Fiji [lower value], Egypt [upper value])	Survey year	(Costopoulou et al., 2006)
			MCT (mean): 6.8–37 pg/g fat	Subgroup not specified (age not specified; sample size not specified), 10 countries (among others: Greece [lower value], Finland [upper level])	Survey year	(Costopoulou et al., 2006)
OCPs	TCDD	24-h urine 24-h urine serum whole blood	WHO-TEQ	Adults (24–76 years; n: 126), Slovak Republic	2006–2007	(Chovancova et al., 2012)
			AM: 6.9–28.6 pg WHO-TEQ g ⁻¹ lipid (^{##}) (Max: 88.1 pg WHO-TEQ g ⁻¹ lipid)			
			/	/	/	(Jeanneret et al., 2014)
			/	/	/	(Jeanneret et al., 2014)
	Bile acids steroids β-HCH	serum whole blood	P ₉₅ : 190 ng/g	Adults (18–74 years; n: 386), France	2006–2007	(InVS, 2010)
			P ₉₅ : 0.1 µg/l	Children (7–14 years; n: 1063), Germany	2003–2006	(Becker et al., 2008; Schulz et al., 2009)
			MCT (median): 12–860 ng/g lipid	Adults (age not specified; n: 47–2824), several EU countries (Belgium, Czech Republic, Germany, Italy, Romania, Slovakia, Spain, Sweden, UK)	1992–2009	(Gari et al., 2014)
			P ₉₅ : 0.1–0.3 µg/l ([§])	Children (7–14 years; n: 1063), Germany	2003–2006	(Schulz et al., 2009; Schulz et al., 2011)
	ΣDDTs	breast milk	P ₉₅ : 0.3–0.9 µg/l (^{##})	Adults (18–69 years; n: 2749), Germany	1997–1999	(Schulz et al., 2011; Wilhelm et al., 2003)
			P ₉₅ : 0.07 mg/kg fat	Breast-feeding women (age not specified; sample size not specified), Germany	2004–2005	(Schulz et al., 2011)
			RV ₉₅ : 0.5 mg/kg fat	Breast-feeding women (age not specified; sample size not specified), West Germany	2003–2005	(HBM-UBA, 2008; Schulz et al., 2011)
			RV ₉₅ : 0.7 µg/l	Children (7–14 years; n: 942), West Germany	2003–2006	(Schulz et al., 2009)
DDT + DDE HCB	DDE	blood	RV ₉₅ : 1.5–11 µg/l (^{##})	Adults (18–69 years; n: 2290), West Germany	1997–1999	(Schulz et al., 2012; Wilhelm et al., 2003)
			RV ₉₅ : 1.4 µg/l	Children (7–14 years; n: 137), East Germany	2003–2006	(Schulz et al., 2009)
			RV ₉₅ : 3.0–31.0 µg/l (^{##})	Adults (18–69 years; n: 535), East Germany	1997–1999	(Schulz et al., 2012; Wilhelm et al., 2003)
			P ₉₅ : 730 ng/g lipid	Adults (18–74 years; n: 386), France	2006–2007	(InVS, 2010)
	DDT + DDE HCB	serum breast milk plasma	MCT (median): 100–2500 ng/g lipid	Adults (age not specified; n: 47–2824), several EU countries (Belgium, Czech Republic, Germany, Italy, Romania, Slovakia, Spain, Sweden, UK)	1992–2009	(Gari et al., 2014)
			/	/	/	/
			P ₉₅ : 0.06 mg/kg fat	Breast-feeding women (age not specified; sample size not specified), Germany	2004–2005	(HBM-UBA, 2008; Schulz et al., 2011)
			P ₉₅ : 0.13 µg/l	Students (age not specified; n: 116), Germany (Ulm)	2010	(UBA, 2017)
	DDT + DDE HCB	plasma	P ₉₅ : 0.14 µg/l	Students (age not specified; n: 111), Germany (Münster)	2010	(UBA, 2017)
			P ₉₅ : 0.32 µg/l	Students (age not specified; n: 113), Germany (Greifswald)	2010	(UBA, 2017)
						(continued on next page)

Table 2 (continued)

Stressor group Stressor	Biomarker of exposure	Matrix	Reference value P ₉₅ or measure of central tendency	Subgroup (years of age; n: sample size), country	Survey year	Reference
PCBs	serum		P ₉₅ : 0.18 µg/l	Students (age not specified; n: 104), Germany (Halle/Saale)	2010	(UBA, 2017)
			P ₉₅ : 73 ng/g lipid	Adults (18–74 years; n: 386), France	2006–2007	(InVS, 2010)
			MCT (median): 11–2400 ng/g lipid	Population (age not specified; n: 47–2824), several EU countries (Belgium, Czech Republic, Germany, Italy, Romania, Slovakia, Spain, Sweden, UK)	1992–2009	(Gari et al., 2014)
	whole blood		P ₉₅ : 0.1, 0.2 or 0.3 µg/l (***)	Children (7–14 years; n: 1,079), Germany	2003–2006	(Becker et al., 2008; Schulz et al., 2009; Schulz et al., 2011; Schulz et al., 2012)
			P ₉₅ : 0.4–5.8 µg/l (###)	Adults (18–69 years; n: 2824), Germany	1997–1999	(Schulz et al., 2011; Schulz et al., 2012; Wilhelm et al., 2003)
			P ₉₅ : 1 µg/l	Children (7–14 years; n: 1079), Germany	2003–2006	(Becker et al., 2008; Schulz et al., 2009)
	breast milk		RV ₉₅ : 1.1–7.8 µg/l (###)	Adults (18–69 years; n: 2816), Germany	1997–1999	(Schulz et al., 2011; Wilhelm et al., 2003)
			RV ₉₅ : 0.5 mg/kg fat	Breast-feeding women (age not specified; sample size not specified), West Germany	2003–2005	(HBM-UBA, 2008; Schulz et al., 2011)
		plasma	P ₉₅ : 0.73–0.82 µg/l (###)	Children (6–17 years; n: 601), Germany	2010–2014	(Schettgen et al., 2015)
		serum	P ₉₅ : 0.88–4.82 µg/l (###)	Adults (18–65 years; n: 2317), Germany	2010–2014	(Schettgen et al., 2015)
PFC	dioxin-like PCBs	serum	P ₉₅ : 720 ng/g lipid	Adults (18–74 years; n: 386), France	2006–2007	(InVS, 2010)
		breast milk	MCT (mean): 1.8–20.0 pg/g fat	Subgroup not specified (age not specified; sample size not specified), 27 countries (among others: Fiji [lower value], Ukraine [upper value])	Survey year not specified	(Costopoulou et al., 2006)
		serum	MCT (mean): 1.2–6.4 pg/g fat	Subgroup not specified (age not specified; sample size not specified), 10 countries (among others: Greece [lower value], New Zealand [upper value],	Survey year not specified	(Costopoulou et al., 2006)
	indicator PCBs		WHO-TEQ	Adults (24–74 years; n: 126), Slovak Republic	2006–2007	(Chovancova et al., 2012)
		breast milk	AM: 13.6–47.5 pg WHO-TEQ g ⁻¹ lipid (*) (Max: 220 pg WHO-TEQ g ⁻¹ lipid)	Subgroup not specified (age not specified; sample size not specified), 27 countries (among others: Fiji [lower value], Czech Republic [upper value])	Survey year not specified	(Costopoulou et al., 2006)
			MCT (mean): 17–502 ng/g fat	Children (7–14 years; n: 1079), Germany	2003–2006	(Becker et al., 2008; Schulz et al., 2009)
	PCB 28	whole blood	P ₉₅ : < 0.1 µg/l (*)	Children (7–14 years; n: 1079), Germany	2003–2006	(Becker et al., 2008; Schulz et al., 2009)
	PCB 52	whole blood	P ₉₅ : < 0.1 µg/l (*)	Children (7–14 years; n: 1079), Germany	2003–2006	(Becker et al., 2008; Schulz et al., 2009)
	PCB 101	whole blood	P ₉₅ : < 0.1 µg/l (*)	Children (7–14 years; n: 1079), Germany	2003–2006	(Becker et al., 2008; Schulz et al., 2009)
	PFOA	plasma	Preliminary P ₉₅ : 10 µg/l	Children (6 years; n: 80), Germany	2003–2006	(Wilhelm et al., 2009)
	serum		Preliminary P ₉₅ : 10 µg/l	Males (5–77 years; n: 342), Germany	2003–2006	(Wilhelm et al., 2009)
			Preliminary P ₉₅ : 10 µg/l	Females (5–84; n: 317), Germany	2003–2006	(Wilhelm et al., 2009)
			MCT (mean): 4–20 µg/l	Population (age not specified; sample size not specified), several European countries	Survey year not specified	(Stahl et al., 2011)
	cord blood serum		GM: 0.716 ng/ml (Max: 8.97 ng/ml)	Adults (18–65 years; n: 300) Czech Republic	2015	(Sochorova et al., 2017)
			Mean: 1.92–3.88 ng/ml ⁻¹ (###) (Max: 10.21 ng/ml)	Adults (15–89 years; n: 142), Greece	2009	(Vassiliadou et al., 2010)

(continued on next page)

Table 2 (continued)

Stressor group Stressor	Biomarker of exposure	Matrix	Reference value RV ₉₅ (otherwise P ₉₅ or measure of central tendency)	Subgroup (years of age; n: sample size), country	Survey year	Reference
other organic contaminants bisphenols	PFOS	plasma	Preliminary P ₉₅ : 10 µg/l Preliminary P ₉₅ : 25 µg/l	Children (6 years old; n: 170), Germany	2003-2006	(Wilhelm et al., 2009)
			Preliminary P ₉₅ : 15 µg/l	Males (5-77 years; n: 443), Germany	2003-2006	(Wilhelm et al., 2009)
		serum	MCT (mean): 4-55 µg/l	Females (5-84 years; n: 539), Germany Population (age not specified; sample size not specified), several European countries (e.g., Italy [lower value], Poland [upper value])	2003-2006 Survey year not specified	(Wilhelm et al., 2009) (Stahl et al., 2011)
	ΣBPA		GM: 2.29 ng/ml (Max: 51.1 ng/ml) Mean: 7.49-14.93 ng/ml (#)# (Max: 40.36 ng/ml)	Adults (18-65 years; n: 300), Czech Republic Adults (24-87 years; n: 142), Greece	2015 2009	(Sochorova et al., 2017) (Vassiliadou et al., 2010)
		cord blood serum	GM: 1.10 µg/l	Children (newborns; n: 269), Belgium	2012-2015	(Schoeters et al., 2016)
		24-h urine	Average concentration: 1-3 µg/l	Population (age not specified; sample size not specified), several cohorts from Japan, USA	Survey year not specified	(Dekant and Volkel, 2008)
		spot urine	Median: 1.51 µg/l Median: 3.78 µg/l	General adults (51 ± 12 years, n: 122), Cyprus General adults (47 ± 13 years, n: 90), Romania	2013-2014 2014-2015	(Andrianou et al., 2016) (Andrianou et al., 2016)
		24-h urine first morning urine	P ₉₅ : 7.07 µg/l P ₉₅ : 30 µg/l P ₉₅ : 15 µg/l P ₉₅ : 7 µg/l P ₉₅ : 13.1 µg/l	Students (age not specified; n: 60), Germany (Münster) Children (3-5 years; n: 137), Germany Children (6-14 years; n: 462), Germany Adults (20-29 years; n: 600), Germany Children (5-12 years, n: 653), several European countries (Belgium, Denmark, Luxembourg, Slovenia, Spain, Sweden)	2009 2003-2006 2003-2006 1995-2009 2011-2012	(UBA, 2017) (HBM-UBA, 2012) (HBM-UBA, 2012) (HBM-UBA, 2012) (Covaci et al., 2015)
			P ₉₅ : 11.1 µg/l	Mothers (age not specified; n: 639), several European countries (Belgium, Denmark, Luxembourg, Slovenia, Spain, Sweden)	2011-2012	(Covaci et al., 2015)
	OPP	first morning urine	RV ₉₅ : < 0.3 µg/l (§§)	Children (3-14 years, n: 599), Germany	2003-2006	(Becker et al., 2008; Schulz et al., 2009)
		first morning urine	RV ₉₅ : 10 µg/l	Children (3-14 years, n: 599), Germany	2003-2006	(Becker et al., 2008; Schulz et al., 2009; Schulz et al., 2011)
Diethyl phosphate		urine (not further specified)	P ₉₅ : 6.53 µg/g crea.	Adults (18-74 years; n: 392), France	2006-2007	(InVS, 2010)
		first morning urine	RV ₉₅ : 30 µg/l	Children (3-14 years, n: 599), Germany	2003-2006	(Becker et al., 2008; Schulz et al., 2009; Schulz et al., 2011)
			RV ₉₅ : 16 µg/l	General population (children and adults, age not specified; n: 1149), Germany	1998	(HBM-UBA, 2003; Heudorf et al., 2006; Schulz et al., 2011)
	DMDTP	urine (not further specified)	P ₉₅ : 15.91 µg/g crea.	Adults (18-74 years; n: 392), France	2006-2007	(InVS, 2010)
		first morning urine	RV ₉₅ : 10 µg/l	Children (3-14 years, n: 599), Germany	2003-2006	(Becker et al., 2008; Schulz et al., 2009)
		first morning urine	P ₉₅ : 75 µg/l	Children (3-14 years; n: 599), Germany	2003-2006	(Becker et al., 2008; Schulz et al., 2009; Schulz et al., 2011)
	DMP	spot urine	P ₉₅ : 135 µg/l	General population (children and adults, age not specified; n: 1149), Germany (Frankfurt/Main)	1998	(HBM-UBA, 2003; Heudorf and Angerer, 2001; Heudorf et al., 2006; Schulz et al., 2011)
	DMTP	first morning urine	RV ₉₅ : 100 µg/l	Children (3-14 years; n: 599), Germany	2003-2006	(Becker et al., 2008; Schulz et al., 2009; Schulz et al., 2011)
						(continued on next page)

Table 2 (continued)

Stressor group Stressor	Biomarker of exposure	Matrix	Reference value RV ₉₅ (otherwise P ₉₅ or measure of central tendency)	Subgroup (years of age; n: sample size), country	Survey year	Reference
PAHs	1-hydroxypyrene	spot urine	RV ₉₅ : 160 µg/l	General population (children and adults; age not specified; n: 1149), Germany	1998	(HBM-UBA, 2003; Heudorf and Angerer, 2001; Heudorf et al., 2006; Schulz et al., 2011) (Health Canada, 2013) (InVS, 2010)
		urine (not further specified)	P ₉₅ : 37 µg/l	General population (6-79 years; n: 2559), Canada Adults (18-74 years; n: 392), France	2009-2011	(Becker et al., 2006)
		morning urine	P ₉₅ : 48.74 µg/g crea.		2006-2007	
		(not specified if first morning urine)	P ₉₅ : 124 µg/l			
		first morning urine	P ₉₅ : 23.83 µg/g crea.			
	fluoranthene 3-hydroxyfluoranthene	spot urine	P ₉₅ : 210.9 µg/l	Children (2-17 years; n: 363), Germany	2001-2002	(Roca et al., 2014)
			P ₉₅ : 23.83 µg/g crea.	Children (6-11 years; n: 125), Spain (Valencia)	2010	
			P ₉₅ : 210.9 µg/l	Children (3-7 years; n: 89), Canada (Quebec)	2003	
			Median: 99.3 nmol/g crea. (Max: 1526.0 nmol/g crea.)	Children (6-7 years; n: 195), Italy (Siena)	1995	
			P ₉₅ : 62.0 µg/l	Children (6-11 years; n: 471), USA	1999-2000	
PAHs	fluorene 2-hydroxyfluorene 3-hydroxyfluorene		P ₉₅ : 69.0 µg/l	Adolescents (12-19 years; n: 664), USA	1999-2000	(Barr et al., 2004) (Barr et al., 2004) (Barr et al., 2004) (Wilhelm et al., 2008) (Wilhelm et al., 2008) (Grainger et al., 2006)
			P ₉₅ : 38.0 µg/l	Adults (20-59 years; n: 814), USA	1999-2000	
			RV ₉₅ : 0.5 µg/l	Non-smoking adults (18-69 years; n: 389), Germany	1997-1999	
			RV ₉₅ : 0.5 µg/l	Non-smoking children (3-14 years; n: 571), Germany	2003-2004	
			P ₉₅ : 730 ng/l	Population (≥ 6 years; n: 2312), USA	1999-2000	
	naphthalene	urine (not further specified)		Population (≥ 6 years; n: 2236), USA	1999-2000	(Grainger et al., 2006) (Grainger et al., 2006) (Grainger et al., 2006) (Fustinoni et al., 2010)
		urine (not further specified)	P ₉₅ : 98.8 ng/l		1999-2000	
		urine (not further specified)	P ₉₅ : 6450 ng/l	Population (≥ 6 years; n: 2315), USA	1999-2000	
		urine (not further specified)	P ₉₅ : 3390 ng/l	Population (≥ 6 years; n: 2312), USA	1999-2000	
		spot urine (sampled three times during a week)	P ₉₅ (geometric mean of three determinations): 266 ng/l	General population (19 to 75 years; n: 100), Italy (Milan)	2007-2008	
Parabens	1-naphthol	spot urine	P ₉₅ : 108.0 nmol/l	General adults (> 18 years; n: 298), UK	2006	(IEH, 2008) (Preuss et al., 2004; Wilhelm et al., 2008) (IEH, 2008) (Preuss et al., 2004; Wilhelm et al., 2008)
		morning urine	P ₉₅ : 10.7-29.9 µg/l	Non-smoking general population (2.5-51 years; n = 259), 4 cohorts from Germany	Survey year not specified	
		spot urine	P ₉₅ : 81.0 nmol/l	General adults (> 18 years; n: 298), UK	2006	
		morning urine	P ₉₅ : 6.5-17.1 µg/l	Non-smoking general population (2.5-51 years; n = 259), 4 cohorts from Germany	Survey year not specified	
		urine (not further specified)	P ₉₅ : 1070 ng/l	Population (≥ 6 years; n: 2246), USA	1999-2000	
	2-naphthol	urine (not further specified)	P ₉₅ : 828 ng/l	Population (≥ 6 years; n: 2179), USA	1999-2000	(Grainger et al., 2006) (Grainger et al., 2006) (Grainger et al., 2006)
		urine (not further specified)	P ₉₅ : 657 ng/l	Population (≥ 6 years; n: 2299), USA	1999-2000	
		spot urine	P ₉₅ : 974 µg/l	General population (≥ 6 years; n: 2548), USA	2005-2006	
		spot urine	P ₉₅ : 299 µg/l	General population (≥ 6 years; n: 2548), USA	2005-2006	
		spot urine	P ₉₅ : 19.6 µg/l	General population (≥ 6 years; n: 2548), USA	2005-2006	
Phthalates	phenanthrene 1-hydroxyphenanthrene 2-hydroxyphenanthrene 3-hydroxyphenanthrene	spot urine	P ₉₅ : 57.2 µg/l	General population (≥ 6 years; n: 2548), USA	2005-2006	(Calafat et al., 2010) (Calafat et al., 2010) (Calafat et al., 2010) (Calafat et al., 2010)
		spot urine				
		spot urine				

(continued on next page)

Table 2 (continued)

Stressor group	Biomarker of exposure	Matrix	Reference value P ₉₅ or measure of central tendency)	Subgroup (years of age; n; sample size), country	Survey year	Reference
PYR	5OH-MEHP 5oxo-MEHP DEP	spot urine spot urine urine	P ₉₅ : 146.0 nmol/l P ₉₅ : 230.0 nmol/l /	General adults (> 18 years; n: 337), UK General adults (> 18 years; n: 337), UK /	2006 2006 /	(IEH, 2008) (IEH, 2008) (Koch and Angerer, 2012)
	MEP	urine	/	/	/	(Koch and Angerer, 2012)
	DiBP	urine	/	/	/	(Koch and Angerer, 2012)
	DiNP	urine	/	/	/	(Koch and Angerer, 2012)
	DnBP	urine	/	/	/	(Koch and Angerer, 2012)
	MnBP	urine	/	/	/	(Koch and Angerer, 2012)
	ΣPYR ^(§§§)	breast milk	Mean: 4.89 ng/g ¹ lipid weight (Max: 7.79 ng/g ¹ lipid weight)	Mothers (age not specified; n: 6), Spain	2009	(Corcellas et al., 2012)
	3PBA	spot urine urine (not further specified) first morning urine	P ₉₅ : 28.3 nmol/l P ₉₅ : 3.48 µg/g crea. RV ₉₅ : 2 µg/l	General adults (> 18 years; n: 336), UK Adults (18-74 years; n: 396), France General population (children and adults; age not specified; n: 1149), Germany	2006 2006-2007 1998	(IEH, 2008) (InVS, 2010) (HBM-UBA, 2003; Heudorf et al., 2006; Schulz et al., 2011)
	cyfluthrin	first morning urine	RV ₉₅ : 2 µg/l	Children (3-14 years; n: 598), Germany	2003-2006	(Becker et al., 2008; Schulz et al., 2011)
		spot urine	P ₉₅ : 28.3 nmol/l	General adults (> 18 years; n: 336), UK	2006	(IEH, 2008)
deltamethrin	cis-Cl ₂ CA [also a biomarker for deltamethrin]	spot urine urine (not further specified) spot urine	P ₉₅ : 3.8 nmol/l P ₉₅ : 1.24 µg/g crea. P ₉₅ : 10.4 nmol/l	General adults (> 18 years; n: 336), UK Adults (18-74 years; n: 396), France General adults (> 18 years; n: 92), UK	2006 2006-2007 2006	(IEH, 2008) (InVS, 2010) (IEH, 2008)
	cis-Cl ₂ CA & trans-Cl ₂ CA [also a biomarker for deltamethrin]	spot urine	P ₉₅ : 7.7 nmol/l	General adults (> 18 years; n: 335), UK	2006	(IEH, 2008)
	trans-Cl ₂ CA [also a biomarker for deltamethrin]	urine (not further specified)	P ₉₅ : 2.64 µg/g crea.	Adults (18-74 years; n: 396), France	2006-2007	(InVS, 2010)
	Br ₂ CA	spot urine	P ₉₅ : 5.3 nmol/l	General adults (> 18 years; n: 336), UK	2006	(IEH, 2008)
		urine (not further specified)	P ₉₅ : 2.18 µg/g crea.	Adults (18-74 years; n: 396), France	2006-2007	(InVS, 2010)
		spot urine	P ₉₅ : 3.8 nmol/l	General adults (> 18 years; n: 336), UK	2006	(IEH, 2008)
	cis-Cl ₂ CA [also a biomarker for cyfluthrin]	urine (not further specified)	P ₉₅ : 1.24 µg/g crea.	Adults (18-74 years; n: 396), France	2006-2007	(InVS, 2010)
	cis-Cl ₂ CA & trans-Cl ₂ CA [also a biomarker for cyfluthrin]	spot urine	P ₉₅ : 10.4 nmol/l	General adults (> 18 years; n: 92), UK	2006	(IEH, 2008)
	trans-Cl ₂ CA [also a biomarker for cyfluthrin]	spot urine	P ₉₅ : 7.7 nmol/l	General adults (> 18 years; n: 335), UK	2006	(IEH, 2008)
		urine (not further specified)	P ₉₅ : 2.64 µg/g crea.	Adults (18-74 years; n: 396), France	2006-2007	(InVS, 2010)
toxic and potential toxic elements As	As	24-h urine	P ₉₅ : 41.9 µg/l P ₉₅ : 46.4 µg/l P ₉₅ : 57.7 µg/l P ₉₅ : 32.6 µg/l	Students (age not specified; n: 126), Germany (Münster) Students (age not specified; n: 132), Germany (Greifswald) Students (age not specified; n: 118), Germany (Halle/Saale) Students (age not specified; n: 131), Germany (Ulm)	2016 2016 2016 2016	(UBA, 2017) (UBA, 2017) (UBA, 2017) (UBA, 2017)

(continued on next page)

Table 2 (continued)

Stressor group	Biomarker of exposure	Matrix	Reference value P ₉₅ or measure of central tendency	Subgroup (years of age; n: sample size), country	Survey year	Reference
Cd	Cd	first morning urine	P ₉₅ : 18.9 µg/l	General adults (18–96 years; n: 4741), Germany	1997–1999	(Wilhelm et al., 2004)
			RV ₉₅ : 15.0 µg/l	Adults who did not eat fish 48 h before sample collection (18–69 years; n: 3924), Germany	1997–1999	(Schulz et al., 2011; Wilhelm et al., 2004)
			P ₉₅ : 14.0 µg/l	Children (3–14 years; n: 1734), Germany	2003–2006	(Becker et al., 2008)
			RV ₉₅ : 15.0 µg/l	Children who did not eat fish 48 h before sample collection (3–14 years; n: 1487), Germany	2003–2006	(Schulz et al., 2009; Schulz et al., 2011)
		urine (not further specified)	P ₉₅ : 61.3 µg/g crea.	Adults (18–74 years; n: 1515), France	2006–2007	(InVS, 2010)
Cd	Cd	24-h urine	P ₉₅ : 0.23 µg/l	Students (age not specified; n: 126), Germany (Münster)	2016	(UBA, 2017)
			P ₉₅ : 0.28 µg/l	Students (age not specified; n: 132), Germany (Greifswald)	2016	(UBA, 2017)
			P ₉₅ : 0.29 µg/l	Students (age not specified; n: 118), Germany (Halle/Saale)	2016	(UBA, 2017)
			P ₉₅ : 0.25 µg/l	Students (age not specified; n: 131), Germany (Ulm)	2016	(UBA, 2017)
		first morning urine	RV ₉₅ : 0.2 µg/l	Non-smoking children (3–14 years; n: 1667), Germany	2003–2006	(Becker et al., 2008; Schulz et al., 2009; Schulz et al., 2011)
Cr	Cr		RV ₉₅ : 0.8 µg/l	Non-smoking adults (18–69 years; n: 3128), Germany	1997–1999	(Schulz et al., 2011; Wilhelm et al., 2004)
		spot urine	P ₉₅ : 7.9 nmol/l	General adults (> 18 years; n: 362), UK	2006	(IEH, 2008)
		blood	RV ₉₅ : < 0.3 µg/l	Non-smoking children (3–14 years; n: 1498), Germany	2003–2006	(Becker et al., 2008; Schulz et al., 2009; Schulz et al., 2011)
			RV ₉₅ : 1.0 µg/l	Non-smoking adults (18–69 years; n: 3061), Germany	1997–1999	(Wilhelm et al., 2004)
		blood	Rt: 0.7–28.0 µg/l	Population (age not specified; sample size not specified), country not specified	Survey year not specified	(Burtis et al., 2012)
Cr	Cr	24-h urine	Reference value(**): < 0.2 µg/l	Population (age not specified; sample size not specified), country not specified	Survey year not specified	(Burtis et al., 2012)
		serum	Rt: 0.1–0.2 µg/l	Population (age not specified; sample size not specified), country not specified	Survey year not specified	(Burtis et al., 2012)
Cu	Cu	24-h urine	P ₅ -P ₉₅ : 3.63–13.9 µg/l	Students (age not specified; n: 126), Germany (Münster)	2016	(UBA, 2017)
			P ₅ -P ₉₅ : 4.01–14.9 µg/l	Students (age not specified; n: 132), Germany (Greifswald)	2016	(UBA, 2017)
			P ₅ -P ₉₅ : 3.22–13.4 µg/l	Students (age not specified; n: 118), Germany (Halle/Saale)	2016	(UBA, 2017)
			P ₅ -P ₉₅ : 2.72–13.4 µg/l	Students (age not specified; n: 131), Germany (Ulm)	2016	(UBA, 2017)
		plasma	P ₅ -P ₉₅ : 0.70–1.84 µg/l	Students (age not specified; n: 125), Germany (Münster)	2016	(UBA, 2017)
Cu	Cu		P ₅ -P ₉₅ : 0.75–2.12 mg/l	Students (age not specified; n: 131), Germany (Greifswald)	2016	(UBA, 2017)
			P ₅ -P ₉₅ : 0.70–1.89 mg/l	Students (age not specified; n: 118), Germany (Halle/Saale)	2016	(UBA, 2017)
			P ₅ -P ₉₅ : 0.70–1.89 mg/l	Students (age not specified; n: 132), Germany (Ulm)	2016	(UBA, 2017)
		serum	Rt: 70–140 µg/dl	Men (age not specified; sample size not specified), country not specified	Survey year not specified	(Burtis et al., 2012)
			Rt: 80–155 µg/dl	Women (age not specified; sample size not specified), country not specified	Survey year not specified	(Burtis et al., 2012)

(continued on next page)

Table 2 (continued)

Stressor group Stressor	Biomarker of exposure	Matrix	Reference value P ₉₅ or measure of central tendency	Subgroup (years of age; n: sample size), country	Survey year	Reference
Hg	Hg	24-h urine	P ₉₅ : 0.23 µg/l	Students (age not specified; n: 126), Germany (Münster)	2016	(UBA, 2017)
			P ₉₅ : 0.37 µg/l	Students (age not specified; n: 132), Germany (Greifswald)	2016	(UBA, 2017)
			P ₉₅ : 0.36 µg/l	Students (age not specified; n: 118), Germany (Halle/Saale)	2016	(UBA, 2017)
			P ₉₅ : 0.22 µg/l	Students (age not specified; n: 132), Germany (Ulm)	2016	(UBA, 2017)
			P ₉₅ : 1.9 µg/g	Mothers (< 45 years; n: 1839), Europe (Belgium, Switzerland, Cyprus, Czech Republic, Germany, Denmark, Spain, Hungary, Ireland, Luxembourg, Poland, Portugal, Romania, Sweden, Slovenia, Slovak Republic, UK)	2011–2012	(Den Hond et al., 2015)
Mn	Mn	blood	P ₉₅ : 1.3 µg/g	Children (5–11 years; n: 1836), Europe (Belgium, Switzerland, Cyprus, Czech Republic, Germany, Denmark, Spain, Hungary, Ireland, Luxembourg, Poland, Portugal, Romania, Sweden, Slovenia, Slovak Republic, UK)	2011–2012	(Den Hond et al., 2015)
			P ₉₅ : 1.8 µg/g	Adults (18–74 years; n: 365), France	2006–2007	(InVS, 2010)
			P ₉₅ : 1.2 µg/g	Children (3–17 years; n: 1364), France	2006–2007	(InVS, 2010)
			RV ₉₅ : 0.8 µg/l	Children who ate fish ≤ 3 times per month (3–14 years; n: 891), Germany	2003–2006	(Schulz et al., 2009; Schulz et al., 2011)
			RV ₉₅ : 2.0 µg/l	Adults who ate fish ≤ 3 times per month (18–69 years; n: 2310), Germany	1997–1999	(Wilhelm et al., 2004)
Pb	Pb	first morning urine	RV ₉₅ : 0.4 µg/l	Children without amalgam fillings (3–14 years; n: 1612), Germany	2003–2006	(Schulz et al., 2009; Schulz et al., 2011)
			RV ₉₅ : 1.0 µg/l	Adults without amalgam fillings (18–69 years; n: 1560), Germany	1997–1999	(Wilhelm et al., 2004)
			P ₉₅ : 15 nmol/l	General adults (> 18 years; n: 362), UK	2006	(IEH, 2008)
			RI: 5–15 µg/l	(age not specified; sample size not specified), country not specified	Survey year	(Burtis et al., 2012)
			RI: 0.5–1.3 µg/l	(age not specified; sample size not specified), country not specified	Survey year	(Burtis et al., 2012)
Pb	Pb	urine (not further specified)	RI: 0.5–9.8 µg/l	(age not specified; sample size not specified), country not specified	Survey year	(Burtis et al., 2012)
			RV ₉₅ : 35 µg/l	Children (3–14 years; n: 1560), Germany	not specified	(Schulz et al., 2009; Schulz et al., 2011)
			RV ₉₅ : 70 µg/l	Women (18–69 years; n: 2303), Germany	2003–2006	(Schulz et al., 2009; Schulz et al., 2011)
			RV ₉₅ : 90 µg/l	Men (18–69 years; n: 2342), Germany	1997–1999	(Wilhelm et al., 2004)
			P ₉₅ : 18.5 µg/l	Students (age not specified; n: 126), Germany (Münster)	1997–1999	(Schulz et al., 2011; Wilhelm et al., 2004)
Se	Se	blood	P ₉₅ : 26.4 µg/l	Students (age not specified; n: 132), Germany (Greifswald)	2016	(UBA, 2017)
			P ₉₅ : 22.5 µg/l	Students (age not specified; n: 116), Germany (Halle/Saale)	2016	(UBA, 2017)
			P ₉₅ : 21.8 µg/l	Students (age not specified; n: 130), Germany (Ulm)	2016	(UBA, 2017)
			RI: 60–120 µg/l (females); 79–130 µg/l (males)	General population (age not specified; sample size not specified), country not specified	Survey year	(Wilhelm et al., 2004)
			P ₅ -P ₉₅ : 68.2–109 µg/l	Students (age not specified; n: 125), Germany (Münster)	not specified	(UBA, 2017)
Zn	Zn	plasma	P ₅ -P ₉₅ : 69.6–111 µg/l	Students (age not specified; n: 131), Germany (Greifswald)	2016	(UBA, 2017)
			P ₅ -P ₉₅ : 64.6–109 µg/l	Students (age not specified; n: 118), Germany (Halle/Saale)	2016	(UBA, 2017)
			P ₅ -P ₉₅ : 70.8–114 µg/l	Students (age not specified; n: 132), Germany (Ulm)	2016	(UBA, 2017)
			RI: 16–71 µg/l	Children (< 2 years; sample size not specified), country not specified	Survey year	(Burtis et al., 2012)
			RI: 40–103 µg/l	Children (2–4 years; sample size not specified), country not specified	not specified	(Burtis et al., 2012)
Zn	Zn	24-h urine	RI: 55–134 µg/l	Children (4–16 years; sample size not specified), country not specified	Survey year	(Burtis et al., 2012)
			RI: 63–160 µg/l	Adults (age not specified; sample size not specified), country not specified	Survey year	(Burtis et al., 2012)
			P ₅ -P ₉₅ : 48.8–529 µg/l	Students (age not specified; n: 126), Germany (Münster, Ulm)	not specified	(Burtis et al., 2012)
					2016	(UBA, 2017)
						(continued on next page)

Table 2 (continued)

Stressor group Stressor	Biomarker of exposure	Matrix	Reference value RV ₉₅ (otherwise P ₉₅ or measure of central tendency)	Subgroup (years of age; n: sample size), country	Survey year	Reference
volatile organic compounds acrylamide	AAMA	plasma	P ₅ -P ₉₅ : 55.8-527 µg/l	Students (age not specified; n: 132), Germany (Greifswald)	2016	(UBA, 2017)
			P ₅ -P ₉₅ : 54.2-559 µg/l	Students (age not specified; n: 118), Germany (Halle/Saale)	2016	(UBA, 2017)
			P ₅ -P ₉₅ : 50.0-603 µg/l	Students (age not specified; n: 131), Germany (Ulm)	2016	(UBA, 2017)
			P ₅ -P ₉₅ : 0.56-0.94 mg/l	Students (age not specified; n: 125), Germany (Münster)	2016	(UBA, 2017)
			P ₅ -P ₉₅ : 0.55-1.07 mg/l	Students (age not specified; n: 131), Germany (Greifswald)	2016	(UBA, 2017)
	GAMA	serum	P ₅ -P ₉₅ : 0.55-0.88 mg/l	Students (age not specified; n: 118), Germany (Halle/Saale)	2016	(UBA, 2017)
			P ₅ -P ₉₅ : 0.55-0.93 mg/l	Students (age not specified; n: 132), Germany (Ulm)	2016	(UBA, 2017)
			Rt: 80-120 µg/dl	Adults (age not specified; sample size not specified), country not specified	Survey year not specified	(Burtis et al., 2012)
			P ₉₅ : 139 µg/l	Children (6-11 years; n: 394), USA	2011-2012	(CDC, 2017)
			P ₉₅ : 279 µg/l	Children (12-19 years; n: 384), USA	2011-2012	(CDC, 2017)
benzene	GAMA	urine (not further specified)	P ₉₅ : 285 µg/l	Adults (≥ 20 years; n: 1688), USA	2011-2012	(CDC, 2017)
			P ₉₅ : 50.9 µg/l	Children (6-11 years; n: 394), USA	2011-2012	(CDC, 2017)
			P ₉₅ : 68.0 µg/l	Children (12-19 years; n: 384), USA	2011-2012	(CDC, 2017)
			P ₉₅ : 64.5 µg/l	Adults (≥ 20 years; n: 1688), USA	2011-2012	(CDC, 2017)
			MCT (mean, median, or GM): 50-200 ng/l	Non-smoking general population (age not specified; sample size not specified), countries not specified (several cohorts)	Survey year not specified	(Arnold et al., 2013)
	benzene	blood	P ₉₅ : 0.120 ng/ml	Children (12-19 years; n: 912), USA	2007-2008	(CDC, 2017)
			P ₉₅ : 0.328 ng/ml	Adults (20-59 years; n: 1445), USA	2007-2008	(CDC, 2017)
			P ₉₅ : 0.213 ng/ml	Adults (≥ 60 years; n: 814), USA	2007-2008	(CDC, 2017)
			P ₉₅ : 311.5 ng/l	Non-smoking and non-occupationally exposed general population (27-78 years; n: 86), Italy	2006-2007	(Campagna et al., 2014)
			MCT (mean, median, or GM): 0.10-0.25 µg/l	Non-smoking general population (age not specified; sample size not specified), countries not specified (several cohorts)	Survey year not specified	(Arnold et al., 2013)
cyanide	S-PMA	urine (not further specified)	P ₉₅ (geometric mean of three determinations): 1598 ng/l	General population (19 to 75 years; n: 100), Italy (Milan)	2007-2008	(Fustinoni et al., 2010)
			Median: 118 ng l ⁻¹	General population (18-83 years; n: 48), Cyprus (Nicosia)	2013	(Tsangari et al., 2017)
			P ₉₅ : 38.0 nmol/l	Adults (> 18 years; n: 355), UK	2006	(IEH, 2008)
			MCT (mean, median, or GM): 0.5-9 µg/l	Non-smoking general population (age not specified; sample size not specified), countries not specified (several cohorts)	Survey year not specified	(Arnold et al., 2013)
			P ₉₅ : 911 µg/l	Children (6-11 years; n: 394), USA	2011-2012	(CDC, 2017)
	2-Aminothiazoline-4-carboxylic acid	blood	P ₉₅ : 583 µg/l	Children (12-19 years; n: 384), USA	2011-2012	(CDC, 2017)
			P ₉₅ : 483 µg/l	Adults (≥ 20 years; n: 1688), USA	2011-2012	(CDC, 2017)
			P ₉₅ : 0.068 ng/ml	Children (12-19 years; n: 448), USA	2007-2008	(CDC, 2017)
			P ₉₅ : 0.131 ng/ml	Adults (20-59 years; n: 1473), USA	2007-2008	(CDC, 2017)
			P ₉₅ : 0.100 ng/ml	Adults (≥ 60 years; n: 829), USA	2007-2008	(CDC, 2017)
ethylbenzene	ethylbenzene	spot urine	P ₉₅ : 289 ng/l ⁻¹	Primary school children (age not specified; n: 151) Italy (Treviglio)	1995	(Minoia et al., 1996)
			P ₉₅ : 75 ng/l ⁻¹	Primary school children (age not specified; n: 107), Italy (Poggibonsi)	1995	(Minoia et al., 1996)
			P ₉₅ : 98 ng/l ⁻¹	Primary school children (age not specified; n: 139), Italy (Valenza)	1995	(Minoia et al., 1996)
			P ₉₅ (geometric mean of three determinations): 130 ng/l	General population (19 to 75 years; n: 100), Italy (Milan)	2007-2008	(Fustinoni et al., 2010)
	ethylbenzene	spot urine (sampled three times during a week)	Median: 9.2 ng l ⁻¹	General population (18-83 years; n: 48), Cyprus (Nicosia)	2013	(Tsangari et al., 2017)

(continued on next page)

Table 2 (continued)

Stressor group	Biomarker of exposure	Matrix	Reference value P ₉₅ or measure of central tendency	Subgroup (years of age; n: sample size), country	Survey year	Reference
glycol ethers	PGA [also a biomarker for styrene]	urine (not further specified)	P ₉₅ : 508 µg/l P ₉₅ : 662 µg/l P ₉₅ : 732 µg/l	Children (6–11 years; n: 394), USA Children (12–19 years; n: 384), USA Adults (≥ 20 years; n: 2466), USA	2011–2012 2011–2012 2011–2012	(CDC, 2017) (CDC, 2017) (CDC, 2017)
		24-h urine	P ₉₅ : 0.3 mg/l	General population (19–52 years; n: 44), Germany (Bavaria)	2007–2008	(Fromme et al., 2013; HBM-UBA, 2014)
		Blood	P ₉₅ : 2.2 µg/l P ₉₅ : 1.13 µg/l P ₉₅ : 1.15 µg/l P ₉₅ : 1.14 µg/l RV ₉₅ : 5 µg/l Mean: < 5 µg/l	Students (age not specified; n: 116), Germany (Ulm) Students (age not specified; n: 111), Germany (Münster) Students (age not specified; n: 128), Germany (Greifswald) Students (age not specified; n: 104), Germany (Halle/Saale) Adults (18–69 years; n: 691; living in homes without wood preservatives), Germany Not exposed population (age not specified; sample size not specified), Germany	2010 2010 2010 2010 1997–1999 Survey year not specified	(UBA, 2017) (UBA, 2017) (UBA, 2017) (UBA, 2017) (Schulz and Butte, 2007; Schulz et al., 2011) (Scholz 2001a)
	PCP	plasma		Population (age not specified; sample size not specified), country not specified	Survey year not specified	(Scholz 2001a)
		serum	P ₉₅ : 25 µg/l RV ₉₅ : 12 µg/l	Adults (41–65 years; n: 251), Germany	1995–1996	(HBM-UBA, 1999; Schulz et al., 2011)
		first morning urine	RV ₉₅ : 2.0 µg/l (*)	Children (3–14 years; n: 599), Germany	2003–2006	(Becker et al., 2008; Schulz et al., 2011)
	PER	urine (not further specified)	P ₉₅ : < 10 µg/l	Population (age not specified; sample size not specified), country not specified	Survey year not specified	(Scholz 2001a)
		24-h urine	P ₉₅ : 0.30 µg/l P ₉₅ : 0.18 µg/l P ₉₅ : 0.16 µg/l P ₉₅ : 0.17 µg/l	Students (age not specified; n: 116), Germany (Ulm) Students (age not specified; n: 112), Germany (Münster) Students (age not specified; n: 128), Germany (Greifswald) Students (age not specified; n: 105), Germany (Halle/Saale)	2010 2010 2010 2010	(UBA, 2017) (UBA, 2017) (UBA, 2017) (UBA, 2017)
		Blood	P ₉₅ : < 1 µg/l	Population (age not specified; sample size not specified), country not specified	Survey year not specified	(Gratza and Kevekordes, 2001)
		urine (not further specified)	P ₉₅ : 384 µg/l P ₉₅ : 421 µg/l P ₉₅ : 638 µg/l P ₉₅ : 3.21 µg/l P ₉₅ : 3.50 µg/l P ₉₅ : 3.26 µg/l P ₉₅ : 508 µg/l	Children (6–11 years; n: 394), USA Children (12–19 years; n: 384), USA Adults (≥ 20 years; n: 1688), USA Children (6–11 years; n: 394), USA Children (12–19 years; n: 384), USA Adults (≥ 20 years; n: 1688), USA Children (6–11 years; n: 394), USA	2011–2012 2011–2012 2011–2012 2011–2012 2011–2012 2011–2012 2011–2012	(CDC, 2017) (CDC, 2017) (CDC, 2017) (CDC, 2017) (CDC, 2017) (CDC, 2017) (CDC, 2017)
toluene	hippuric acid	urine (not further specified)	P ₉₅ : 0.200 ng/ml P ₉₅ : 512 ng/l	Population (≥ 12 years; n: 950), USA Not occupationally exposed hospital staff and blood donors (20–58 years; n = 81), country not specified (author team from Italy and China)	2001–2008 Survey year not specified	(CDC, 2017) (Brugnone et al., 1993)
		spot urine	P ₉₅ : 0.36 g/g crea.	Non-occupational exposed population (18–60 years; n: 115), Brazil	Survey year not specified	(Siqueira and Paiva, 2002)
		urine (not further specified)	P ₉₅ : 29.7 µg/l P ₉₅ : 36.5 µg/l P ₉₅ : 38.7 µg/l P ₉₅ : 0.318 ng/ml P ₉₅ : 0.839 ng/ml P ₉₅ : 0.610 ng/ml Reference value: < 1 µg/l (**)	Children (6–11 years; n: 394), USA Children (12–19 years; n: 384), USA Adults (≥ 20 years; n: 1688), USA Children (12–19 years; n: 439) USA Adults (20–59 years; n: 1483) USA Adults (≥ 60 years; n: 809) USA Non-smoker (age not specified; sample size not specified), country not specified	2011–2012 2011–2012 2011–2012 2007–2008 2007–2008 2007–2008 Survey year not specified	(CDC, 2017) (CDC, 2017) (CDC, 2017) (CDC, 2017) (CDC, 2017) (CDC, 2017) (Scholz, 2001b)
	N-Acetyl-S-(benzyl)-L-cysteine	blood		Children (6–11 years; n: 394), USA	2007–2008	(Minoia et al., 1996)
		spot urine	P ₉₅ : 481–1361 ng/l ¹ , depending on the city	Primary school children (age not specified; n: 107–147, depending on the city), Italy (Poggibonsi, Treviglio, Valenza)	1995	(continued on next page)
		urine (not further specified)				
	styrene	urine (not further specified)				
		urine (not further specified)				
		urine (not further specified)				
	PGA [also a biomarker for ethylbenzene]	urine (not further specified)				
		urine (not further specified)				
		urine (not further specified)				

Table 2 (continued)

Stressor group Stressor	Biomarker of exposure	Matrix	Reference value RV ₉₅ (otherwise P ₉₅ or measure of central tendency)	Subgroup (years of age; n: sample size), country	Survey year	Reference
pharmaceuticals antibiotics	the substance of interest	spot urine (sampled three times during a week)	P ₉₅ (geometric mean of three determinations): 618 ng/l	General population (19 to 75 years; n: 100), Italy (Milan)	2007-2008	(Fustinoni et al., 2010)
		First morning urine	Median: 124 ng l ⁻¹	General population (18-83 years; n: 48), Cyprus (Nicosia)	2013	(Tsangari et al., 2017)
		urine (not further specified)	P ₉₅ : 124 µg/l	Children (6-11 years; n: 409), USA	2013-2014	(CDC, 2017)
			P ₉₅ : 224 µg/l	Children (12-19 years; n: 462), USA	2013-2014	(CDC, 2017)
			P ₉₅ : 420 µg/l	Adults (≥ 20 years; n: 1815), USA	2013-2014	(CDC, 2017)
		urine (not further specified)	P ₉₅ : 122 µg/l	Children (6-11 years; n: 394), USA	2011-2012	(CDC, 2017)
			P ₉₅ : 173 µg/l	Children (12-19 years; n: 384), USA	2011-2012	(CDC, 2017)
			P ₉₅ : 232 µg/l	Adults (≥ 20 years; n: 1688), USA	2011-2012	(CDC, 2017)
		urine (not further specified)	P ₉₅ : 889 µg/l	Children (6-11 years; n: 394), USA	2011-2012	(CDC, 2017)
			P ₉₅ : 1680 µg/l	Children (12-19 years; n: 384), USA	2011-2012	(CDC, 2017)
chemotherapy tobacco smoke	the substance of interest		P ₉₅ : 1740 µg/l	Adults (≥ 20 years; n: 1688), USA	2011-2012	(CDC, 2017)
			P ₉₅ : 0.216 ng/ml	Children (12-19 years; n: 447), USA	2007-2008	(CDC, 2017)
			P ₉₅ : 0.379 ng/ml	Adults (20-59 years; n: 1520), USA	2007-2008	(CDC, 2017)
			P ₉₅ : 0.308 ng/ml	Adults (≥ 60 years; n: 854), USA	2007-2008	(CDC, 2017)
		spot urine (sampled three times during a week)	P ₉₅ (geometric mean of three determinations): 178 ng/l	General population (19 to 75 years; n: 100), Italy (Milan)	2007-2008	(Fustinoni et al., 2010)
		first morning urine	Median: 29 ng l ⁻¹	General population (18-83 years; n: 48), Cyprus (Nicosia)	2013	(Tsangari et al., 2017)
		spot urine	P ₉₅ : 440.0 µmol/l	Adults (> 18 years; n: 360), UK	2006	(IEH, 2008)
			P ₉₅ : 94.7 mg/g crea.	Adults (> 18 years; n: 360), UK	2006	(IEH, 2008)
		spot urine	P ₉₅ : 230-909 ng/l ⁻¹ , depending on the city	Primary school children (age not specified; n: 96-144, depending on the city), Italy (Poggibonsi, Treviglio, Valenza)	1995	(Minoia et al., 1996)
		Blood	P ₉₅ : 0.066 ng/ml	Children (12-19 years; n: 457), USA	2007-2008	(CDC, 2017)
smoking	the substance of interest		P ₉₅ : 0.086 ng/ml	Adults (20-59 years; n: 1524), USA	2007-2008	(CDC, 2017)
			P ₉₅ : 0.091 ng/ml	Adults (≥ 60 years; n: 854), USA	2007-2008	(CDC, 2017)
		spot urine (sampled three times during a week)	P ₉₅ (geometric mean of three determinations): 64 ng/l	General population (19 to 75 years; n: 100), Italy (Milan)	2007-2008	(Fustinoni et al., 2010)
		first morning urine	Median: 28 ng l ⁻¹	General population (18-83 years; n: 48), Cyprus (Nicosia)	2013	(Tsangari et al., 2017)
		blood	Reference value: < 1 µg/l (**)	Non-smoker (age not specified; sample size not specified), country not specified	Survey year not specified	(Scholz, 2001c)
			/	/	/	/
		blood	/	/	/	/
		urine	/	/	/	/
		urine	/	/	/	/
		spot urine	P ₉₅ : 3230 µg/l	General adults (> 18 years; n: 356), UK	2006	(IEH, 2008)
tobacco smoke	the substance of interest		P ₉₅ : 233.73 µg/g crea.	Ex-smoker (> 18 years; n: 129), UK	2006	(IEH, 2008)
			P ₉₅ : 43.45 µg/g crea.	Never smoker (> 18 years; n: 175), UK	2006	(IEH, 2008)
			P ₉₅ : 7243.47 µg/g crea.	Current smoker (> 18 years; n: 46), UK	2006	(IEH, 2008)
			P ₉₅ : 9162.68 µg/g crea.	Second-hand smoke exposed who shares home with smoker (> 18 years; n: 40), UK	2006	(IEH, 2008)
		Nicotine				
		Cotinine				

(continued on next page)

Table 2 (continued)

Stressor group Stressor	Biomarker of exposure	Matrix	Reference value P ₉₅ or measure of central tendency)	Subgroup (years of age; n: sample size), country	Survey year	Reference
air pollution bioaerosols	Mold	blood	P ₉₅ : 2454.59 µg/g crea.	Second-hand smoke exposed who not shares home with smoker (> 18 years; n: 313), UK	2006	(IEH, 2008)
	SC		/	/	/	(Beirao and Araujo, 2013)
	1-HP	serum spot urine	/	/	/	(Yike et al., 2006)
	NO _x	plasma	P ₇₅ : 0.74 pmol/ml (Max.: 1.40 pmol/ml) Mean: 47.5 µM/l	African American students (12–14 years; n: 24), USA (Harlem/New York City) Healthy subjects (20–69 years; n: 738), country not specified (author team from Japan)	1997 Survey year not specified	(Northridge et al., 1999) (Kawakatsu et al., 2002)
	O ₃	plasma	/	/	/	(Liu et al., 1999)
	2,3-DHBA	morning urine (not specified if first morning urine)	Mean: 31.4 pg/ml (Max.: 392.8 pg/ml)	Children (3–12 years; n: 155), Belgian	2013–2014	(Heyndrickx et al., 2015)
	Mycotoxins	24-h urine	Mean: 56.7 pg/ml (Max.: 1398.0 pg/ml) Mean: 11.89 ng/ml (Max.: 67.36 ng/ml)	Adults (19–65 years; n: 239), Belgian General population 3–85 years; n: 50) Italy	2013–2014 2011	(Heyndrickx et al., 2015) (Solfrizzo et al., 2014)
	CIT	morning urine (not specified if first morning urine)	Mean: 58.4 ng/ml (Max.: 343.0 ng/ml) Mean: 53.8 ng/ml (Max.: 460.8 ng/ml)	Children (3–12 years; n: 155), Belgian Adults (19–65 years; n: 239), Belgian	2013–2014 2013–2014	(Heyndrickx et al., 2015) (Heyndrickx et al., 2015)
	DON	morning urine (not specified if first morning urine)	Mean: 79.5 pg/ml (Max.: 3683.0 pg/ml) Mean: 27.8 pg/ml (Max.: 368.1 pg/ml)	Children (3–12 years; n: 155), Belgian Adults (19–65 years; n: 239), Belgian	2013–2014 2013–2014	(Heyndrickx et al., 2015) (Heyndrickx et al., 2015)
	OTA	24-h urine	Mean: 0.144 ng/ml (Max.: 2.129 ng/ml)	General population 3–85 years; n: 52) Italy	2011	(Solfrizzo et al., 2014)
Water contamination DBPs THMs	TCAA	spot urine	P ₉₅ : 49.6 nmol/l	General adults (> 18 years; n: 330), UK	2006	(IEH, 2008)
	BDCM	blood first morning urine	P ₉₅ : 9.5 pg/ml AM: 131 ng g ⁻¹ (summer), 61 ng g ⁻¹ (winter)	General population (≥20 years; n: 1322), USA General population (18–87 years; n: 310) Cyprus (Nicosia)	2003–2004 2012–2013	(LaKind et al., 2010) (Andrianou et al., 2014)
	bromoform	blood first morning urine	P ₉₅ : 7.2 pg/ml AM: 32 ng g ⁻¹ (summer), 147 ng g ⁻¹ (winter)	General population (≥20 years; n: 1310), USA General population (18–87 years; n: 310) Cyprus (Nicosia)	2003–2004 2012–2013	(LaKind et al., 2010) (Andrianou et al., 2014)
	chloroform	blood first morning urine	P ₉₅ : 50.0 pg/ml AM: 608 ng g ⁻¹ (summer), 243 ng g ⁻¹ (winter)	General population (≥20 years; n: 1238), USA General population (18–87 years; n: 310) Cyprus (Nicosia)	2003–2004 2012–2013	(LaKind et al., 2010) (Andrianou et al., 2014)
	DBCM	blood first morning urine	P ₉₅ : 77 ng g ⁻¹ (summer), 119 ng g ⁻¹ (winter)	General population (≥20 years; n: 1333), USA General population (18–87 years; n: 310) Cyprus (Nicosia)	2003–2004 2012–2013	(LaKind et al., 2010) (Andrianou et al., 2014)
	m ⁷ Gua	spot urine	P ₉₅ : 105 mol/mol crea.	Population-based matched control group without lung cancer (50–64 years; n: 261), Denmark	1993–1997	(Loft et al., 2007)
	Σnitrosamines (°)	24-h urine	Mean: 57.33 nmol/l (Max.: 178.4 nmol/l)	Control group without urinary diversion (age not specified; n: 20), Germany	1989	(Tricker et al., 1989)
	NNAL	spot urine	P ₉₅ : 11.7 pg/ml	Non-tobacco users in general population (≥6 years; n: 4831); USA	2011–2012	(Wei et al., 2016)
	NNK	hair	Mean: 1.1 pg/mg	Non-smokers not exposed at home (age not specified; n: 24) Spain (Barcelona)	Survey year not specified	(Perez-Ortuno et al., 2016)
	NOC	12-h overnight urine	Mean 1.12 µmol/l (Max: 3.8 µmol/l)	Non-smoking healthy adults (age not specified; n: 12), France (Lyon)	Survey year not specified	(Pignatelli et al., 1989)

(continued on next page)

Table 2 (continued)

Stressor group Stressor	Biomarker of exposure	Matrix	Reference value P ₉₅ or measure of central tendency)	Subgroup (years of age; n: sample size), country	Survey year	Reference
UVR	thymine dimers	morning urine	Mean: 189 fmol/mol crea. (Max.: 519 fmol/mol crea.)	Adult lifeguards/farm workers (18-54 years; n: 22), Sweden	2006	(Lijendahl et al., 2013)
occupational Hazards biological chemical cultural factors	see substance of interest					
	see substance of interest					
	see substance of interest					
drug consumption	see substance of interest (two examples below)					
	folate					
nutritional status	vitamin C / ascorbate	serum	P ₉₅ : 28.5 ng/ml	General population (≥ 1 years; n: 16411); USA	2003-2006	(CDC, 2012)
		serum	Adequacy level: ≥ 10 nmol/L	Adults	Survey year not specified	(EFSA, 2014)
		red cells	Adequacy level: ≥ 340 nmol/L	Adults	Survey year not specified	(EFSA, 2014)
physical activity						

Table 3
Exposure limit values.

Stressor group Stressor	Biomarker	Matrix	Exposure limit values (BAT, BEL, HBM, CC, etc.)	Subgroup (years of age), country	Reference
POPs dioxins and furans PCBs	GGT ΣPCBs	serum plasma	Cut-off limit (*): 80 U/L CC: 700 ng/g serum lipid	Children (6–10 years), Italy Children (< 3 years), women in childbearing age, pregnant women, breastfeeding women, USA Adults (excluded: women in childbearing age, pregnant women, breastfeeding women), USA	(Mocarelli et al., 1986) (Aylward et al., 2013) (Aylward et al., 2013)
PFC	PFOA PFOS	plasma	HBM I: 2 ng/ml HBM I: 5 ng/ml	Adults (age not specified), Germany Adults (age not specified), Germany	(HBM-UBA, 2009) (HBM-UBA, 2009)
other organic contaminants BPA	ΣBPA	urine (not further specified) (**)	HBM I: 1.5 ng/l HBM I: 2.5 ng/l	Children (age not specified), Germany Adults (age not specified), Germany	(HBM-UBA, 2012) (HBM-UBA, 2012)
phthalates	Σoxo -and 5OH-MEHP	urine (not further specified) (**)	HBM I: 500 µg/l HBM I: 300 µg/l HBM I: 750 µg/l	Children (6–13 years), Germany Women in child-bearing age, Germany Adult men and women of the general population (≥ 14 years) except women in child-bearing age, Germany	(Schulz et al., 2011) (Schulz et al., 2011) (Schulz et al., 2011)
toxic and potential toxic elements As Cd	As Cd	/	ALARP HBM I: 1 µg/g crea. HBM II: 4 µg/g crea. HBM I: 0.5 µg/g crea HBM II: 2 µg/g crea.	Adults (age not specified), Germany Adults (age not specified), Germany Children (age not specified), Germany Children (age not specified), Germany Adults (age not specified)	(EFSA, 2009) (Schulz et al., 2011) (Schulz et al., 2011) (Schulz et al., 2011) (Burtis et al., 2012)
Cu	Cu	plasma	Concentration indicates probable depletion: < 50 µg/dl Concentration indicates probable depletion: < 30 µg/dl	Infants (age not specified)	(Burtis et al., 2012)
Hg	Hg	blood	HBM I: 5 µg/l HBM I: 15 µg/l HBM I: 7 µg/l	Children and adults (age not specified), Germany Children and adults (age not specified), Germany Children and adults (age not specified), Germany	(Schulz et al., 2011) (Schulz et al., 2011) (Schulz et al., 2011)
Pb Zn	Pb Zn	urine (not further specified) (**) Blood Serum	HBM II: 25 µg/l suspended (***) Concentration suggests likely deficiency: < 30 µg/dl	Children and adults (age not specified), Germany Adults (age not specified)	(Schulz et al., 2011) (Burtis et al., 2012)
Volatile organic compounds glycol ethers	MAA	urine (not further specified) (**)	HBM I: 0.4 mg MAA/g crea. HBM II: 1.6 mg MAA/g crea.	General population (age not specified), Germany General population (age not specified), Germany	(HBM-UBA 2014) (HBM-UBA 2014)
PCP	PCP	urine (not further specified) (**)	HBM I: 25 µg/l HBM I: 20 µg/g crea. HBM II 40 µg/l HBM I: 30 µg/g crea. HBM I: 40 µg/l HBM II: 70 µg/l	General population (age not specified), Germany General population (age not specified), Germany General population (age not specified), Germany General population (age not specified), Germany General population (age not specified), Germany General population (age not specified), Germany	(Schulz et al., 2007, 2011) (Schulz et al., 2007, 2011) (Schulz et al., 2007, 2011) (Schulz et al., 2007, 2011) (Schulz et al., 2007, 2011) (Schulz et al., 2007, 2011)
smoking active tobacco smoke	nicotine	serum	Cut-off points to distinguish tobacco use vs. no tobacco use: 3–20 ng/ml Cut-off points to distinguish smokers from nonsmokers: 3 ng/ml Cut-off points to distinguish tobacco use vs. no tobacco use: 31.5–330 ng/ml	Range of 14 studies (≥ 4 years, depending on study), Germany, India, Italy, Norway, Spain, USA Overall population (≥ 12 years and older), USA	(Kim, 2016) (Benowitz et al., 2009) (Kim, 2016)
second-hand smoke (SHS)	nicotine	urine (sample collection differ depending on the study) first morning urine	Cut-off points to distinguish SHS exposed from non-exposed: 3.2 ng/g crea.;	Range of 5 studies (≥ 18 years, depending on study), several countries (e.g., Poland, USA) Children (5–11 years), Poland	(Lupsa et al., 2015)

(continued on next page)

Table 3 (continued)

Stressor group Stressor	Biomarker	Matrix	Exposure limit values (BAT, BEI, HBM, CC, etc.)	Subgroup (years of age), country	Reference
Occupational Hazards chemical occupational hazards VOCs	2-Butoxy-ethanol	spot urine	Cut-off points to distinguish SHS exposed from non-exposed: 2.2 ng/g crea.; Cut-off points to distinguish SHS exposed from non-exposed 2.5 ng/g crea.	Children (5–11 years), Portugal Children (5–11 years), Romania	
	2-Ethoxy-ethanol	spot urine	Cut-off points to distinguish SHS exposed from non-exposed: 1.2 ng/g crea.	Non-smoking mothers (27–45 years), Poland	
	benzene	spot urine	Cut-off points to distinguish SHS exposed from non-exposed: 2.1 ng/g crea.	Non-smoking mothers (26–45 years), Portugal	
	styrene	spot urine	Cut-off points to distinguish SHS exposed from non-exposed: 1.5 ng/g crea.	Non-smoking mothers (25–45 years), Romania	
	toluene	spot urine	Cut-off points to distinguish SHS exposed from non-exposed: 1.5 ng/g crea.	Non-smoking mothers (25–45 years), Romania	
cultural factors stress	Butoxyacetic acid	spot urine	BAT: 100 mg/l	Employee (age not specified), Germany	(DFG, 2016)
	Ethoxyacetic acid	spot urine	BAT: 200 mg/l (after hydrolysis) BAT: 50 mg/l	Employee (age not specified), Germany Employee (age not specified), Germany	(DFG, 2016) (DFG, 2016)
	S-PMA	spot urine	BEI: 25 µg/g crea.	Employee (age not specified), USA	(ACGIH cited in (Arnold et al. 2013)
	Styrene	spot urine	BAT: 600 mg/g crea.	Employee (age not specified), Germany	(DFG, 2016)
	Toluene	blood	BEI: 0.05 mg/l BAT: 600 µg/l	Employee (age not specified), USA Employee (age not specified), Germany	(ACGIH cited in (ATSDR 2000) (DFG, 2016)
cultural factors stress	Hippuric acid	urine (not further specified)	BEI: 1.6 g/g crea.	Employee (age not specified), USA	(ACGIH cited in (ATSDR 2000)
	o-xylene	blood	BAT: 1.5 mg/l	Employee (age not specified), Germany	(DFG, 2016)
	MHA (all isomers)	spot urine	BAT: 2000 mg/l	Employee (age not specified), Germany	(DFG, 2016)
	cortisol	serum	Early morning concentrations below 140 nmol/L suggests adrenal insufficiency	General population (age not specified), 12 countries (e.g., UK)	(Kazlauskaitė et al., 2008) cited in (El-Farhan et al., 2017)

Abbreviations: ALARP, as low as is reasonably practicable; CC, critical concentration (ANSES, 2013; Aylward et al., 2013); OCPs, organochlorine pesticides; PCP, pentachlorophenol.

Abbreviations of stressor groups and biomarkers are explained in the list of abbreviations at the end of the manuscript.

(*) The cut-off limit is defined as “eight times the SD [standard deviation] value above the mean” (Mocarelli et al., 1986).

(**) In the corresponding publications, it is not further specified into first morning urine, spot urine, or 24-h urine. However, the commission “Human Biomonitoring” of the German Environment Agency published a statement with the opinion, that 24-h urine is favorable but less feasible (HBM-UBA, 2007).

(***) In 2009 the HBM Commission of the German Environment Agency suspended the HBM values for lead in blood of children and adults, because several findings consistently show no threshold levels especially for developmental toxicity in children. The HBM Commission concluded that establishing an effect threshold for blood lead levels would be arbitrary and therefore not justified (Schulz et al., 2011).

Table 4
Biomonitoring equivalent (BE) values for selected stressors (based on WHO, 2015 and supplemented).

Stressor group	Biomarker	Matrix	BE value	Subgroup	Intake-based reference value publishing institute, type of value, (value and unit)	Reference
POPs	BFRs	blood	520 ng/g lipid	/	US EPA, RfD (0.1 µg/kg/day)	(Krishnan et al., 2011)
			190,000 ng/g lipid	/	EU Draft, BMD-L (2 mg/kg/day)	(Aylward and Hays, 2011)
OCs	DDT + DDE	breast milk	190,000 ng/g lipid	/	EU Draft, BMD-L (2 mg/kg/day)	(Aylward and Hays, 2011)
		serum	5,000 ng/g lipid	/	US EPA, RfD; RIVM, TDI; ATSDR, Intermediate oral MRL (0.0005 mg/kg/day)	(Kirman et al., 2011)
		serum	47 ng/g lipid	/	ATSDR, MRL (5×10^{-4} mg/kg/day)	(Aylward et al., 2013)
		24-h urine	2,000 µg/l	/	EFSA, TDI (50 µg/kg/day)	(Krishnan et al., 2010a)
other organic contaminants	bisphenols phthalates	24-h urine	12 µg/l	/	EFSA, TDI (500 µg/kg/day)	(Aylward et al., 2009a)
		24-h urine	0.2 µg/l	/	EFSA, TDI (10 µg/kg/day)	(Aylward et al., 2009a)
		24-h urine	660 µg/l	/	EFSA, TDI (50 µg/kg/day)	(Aylward et al., 2009b)
		24-h urine	1000 µg/l	/	EFSA, TDI (50 µg/kg/day)	(Aylward et al., 2009b)
		24-h urine	1100 µg/l	/	EFSA, TDI (50 µg/kg/day)	(Aylward et al., 2009b)
		24-h urine	18 µg/l	/	USEPA, RfD (800 µg/kg/day)	(Aylward et al., 2009a)
		24-h urine	15 µg/l	children (6–11 years)	EFSA, TDI (150 µg/kg/day)	(Hays et al., 2011)
		24-h urine	10.7 µg/l	adolescents (11–16 years)	EFSA, TDI (150 µg/kg/day)	(Hays et al., 2011)
		24-h urine	12.7 µg/l	men (> 16 years)	EFSA, TDI (150 µg/kg/day)	(Hays et al., 2011)
		24-h urine	10.6 µg/l	women (> 16 years)	EFSA, TDI (150 µg/kg/day)	(Hays et al., 2011)
PYR	cyfluthrin 4F3PBA	24-h urine	0.7 µg/l	children (6–11 years)	EFSA, TDI (150 µg/kg/day)	(Hays et al., 2011)
		24-h urine	0.5 µg/l	adolescents (11–16 years)	EFSA, TDI (150 µg/kg/day)	(Hays et al., 2011)
		24-h urine	0.6 µg/l	men (> 16 years)	EFSA, TDI (150 µg/kg/day)	(Hays et al., 2011)
		24-h urine	0.5 µg/l	women (> 16 years)	EFSA, TDI (150 µg/kg/day)	(Hays et al., 2011)
Toxic and potential toxic elements	inorganic As Cd VOCs	24-h urine	400 µg/l	/	FAO/WHO, ADI (10 µg/kg/day)	(Hays et al., 2009)
		24-h urine	240 µg/l	/	US EPA, chronic RfD (0.024 mg/kg/day)	(Aylward et al., 2013; Hays et al., 2009)
		plasma	20 µg/l	adults	US EPA, RfD (0.01 mg/kg/day)	(Aylward et al., 2011)
		24-h urine	2 µg/l	infants and children	US EPA, RfD (0.001 mg/kg/day)	(Aylward et al., 2011)
Toxic and potential toxic elements	inorganic As Cd VOCs	24-h urine	50 µg/l	adults	US EPA, RfD (0.01 mg/kg/day)	(Aylward et al., 2011)
		24-h urine	7 µg/l	Children (≥ 6 years)	US EPA, RfD, (0.001 mg/kg/day)	(Aylward et al., 2011)
		24-h urine	6.4 µg/l	/	ATSDR, chronic MRL (0.3 µg/kg/day)	(Hays et al., 2010)
		24-h urine	1.2 µg/l	/	FAO/WHO, PTWI (10 µg/kg/day)	(Hays et al., 2008)

(continued on next page)

Table 4 (continued)

Stressor group	Biomarker	Matrix	BE value	Subgroup	Intake-based reference value publishing institute, type of value, (value and unit)	Reference
benzene	benzene	blood	0.15 µg/l	/	US EPA, Chronic RfC (30 µg/m³)	(Hays et al., 2012)
		urine*	0.16 µg/l	/		(Hays et al., 2012)
		blood	1.29 µg/l	/	TCEQ, chronic ReV (280 µg/m³)	(Hays et al., 2012)
		urine*	1.42 µg/l	/		(Hays et al., 2012)
		blood	0.29 µg/l	/	California, CAL REL (60 µg/m³)	(Hays et al., 2012)
ethylbenzene	ethylbenzene	urine*	0.33 µg/l	/		(Hays et al., 2012)
		blood	0.04 µg/l	/	ATSDR, chronic inhalation MRL (10 µg/m³)	(Hays et al., 2012)
		urine*	0.05 µg/l	/		(Hays et al., 2012)
		blood	1 µg/l	/	ATSDR, MRL (0.25 mg/m³)	(Hays et al., 2012)
				/		(Aylward et al., 2010; Aylward et al., 2013)
styrene	styrene	blood	3 µg/l	/	US EPA, RfC (1 mg/m³)	(Aylward et al., 2010; Aylward et al., 2013)
toluene	toluene	blood	50 µg/l	/	US EPA, chronic RfC (128 mg/m³)	(Aylward et al., 2008)
			40 µg/l	/	Health Canada, chronic inhalation TDI (150 mg/m³)	(Aylward et al., 2008)
			3 µg/l	/	WHO, lowest level of chronic occupational toluene exposure unequivocally associated with neurobehavioral functional decrement (332 mg/m³)	(Aylward et al., 2008)
			3 µg/l	/	ATSDR, chronic inhalation MRL (132 mg/m³)	(Aylward et al., 2008)
			30 µg/l	/	ATSDR, acute MRL (150 mg/m³)	(Aylward et al., 2008)
triclosan xylene	Σtriclosan (free plus conjugates) o-xylene	24-h urine	2,600 µg/l	/	EC, SED (0.12 mg/kg/day)	(Krishnan et al., 2010b)
		whole	0.3 µg/l	/	US EPA, RfC (0.1 mg/m³)	(Aylward et al., 2010; Aylward et al., 2013)
		blood		/		
water contamination						
THMs						
bromoform BDCM chloroform DBCM	bromoform BDCM chloroform DBCM	blood	130 pg/ml	/	US EPA, RfD(0.03 mg/kg/day)	(Aylward et al., 2013)
		blood	80 pg/ml	/	US EPA, RfD(0.02 mg/kg/day)	(Aylward et al., 2013)
		blood	230 pg/ml	/	US EPA, RfD (0.01 mg/kg/day)	(Aylward et al., 2013)
		blood	20 pg/ml	/	US EPA, RfD (0.003 mg/kg/day)	(Aylward et al., 2013)

Abbreviations: µg/kg/day, microgram per kilogram per day; µg/l, microgram per liter; BE, biomonitoring equivalents; BMD-L, benchmark dose lower confidence limit; CAL REL, California Acute reference exposure levels; mg/kg/day, milligram per kilogram per day; mg/m³, milligram per cubic meter; pg/ml, pictogram per milliliter; MRL, minimal risk level; pictogram per milliliter; PTWI, provisional tolerable weekly intake; RfC, reference concentrations; RfD, reference doses; SED, systemic exposure dose; TCEQ Rev, reference value of the Texas Commission on Environmental Quality.

* derived by using the urine to blood benzene relationship (Hays et al., 2012).

Abbreviations of stressor groups and biomarkers are explained in the list of abbreviations at the end of the manuscript.

Table 5

Opportunities to collect information about the internal exposure of stressors if no specific biomarker of exposure (BoE) is available.

Stressor group	Stressor	Opportunities
smoking	smoking	Besides measuring cotinine in human specimens (see Table 2) questionnaires are useful to determine internal exposure to smoking. Also, a measurement of expired carbon-monoxide (ECO) is an useful information to determine the internal exposure of smoking (Krautter et al., 2015).
air pollution	PM _{2.5} , PM ₁₀	Exposure is measurable in terms of mass or number and composition of PM (like specific chemicals, e.g. metals, PAHs) (Karanasiou et al., 2014; Kolosnjaj-Tabi et al., 2015), however, no specific biomarker is currently available.
	NO _x NPs	Products of NO _x can be measured in body fluids (Halatek et al., 2005). Markers of NPs exposure can range from measurements of specific NPs components, their metabolites, their reaction with cellular macromolecules such as DNA or protein or other effects on cellular processes taken in various bio specimens.
	ozone	Besides 2,3-DHBA (see Table 1), biomarkers of oxidative stress (e.g., 8-iso-PGF, 8-OHdG) in blood, urine or other fluids can be useful to identify human ozone exposure; however, the marker are not specific to ozone exposure (Chen et al., 2007; Kadiiska et al., 2013; Ren et al., 2011).
	UFPs	
noise	noise	There are no specific markers for noise exposure in the case of non-auditory effects. However, reactions at the organism level can be assessed in terms of effect (immune system, cardiac response). Questionnaires and measurements of the noise level are useful to determine noise exposure.
DNA-damaging agents EMF		Some non-specific biomarkers are discussed (e.g., hormones), however, they cannot be used as effective markers of exposure. Further research is urgently needed.
radon		Radon progeny is measurable in blood, hair, and urine, however, not specific. Radon progeny is also a BoE for radium and uranium. Because radon progeny have short half-lives, the time at which the biological sample is taken relevant to time of exposure may be important (Nazaroff and Nero, 1988).
Occupational Hazards	biological	Besides analyzing the biological agent in body fluids, an occupational exposure can be surveyed by requesting the job history (Nowak, 2010). A Job-Exposure-Matrix (JEM) can be used to draw conclusions about internal exposure (Pearce and Douwes, 2008).
	chemical	Besides using BoE for exposure assessment, a JEM is a possibility to identify internal exposure (Pearce and Douwes, 2008).
	mechanical	A JEM can be used to draw conclusions about internal exposure (Pearce and Douwes, 2008).
	physical	Physical stressors can be measured in specific units (e.g., hertz for the frequency of vibration (Levy et al., 2011); decibel (dB(A)) for the noise level (Nowak, 2010)). Personal dosimetry can measure ionizing radiation (Liljendahl et al., 2013). High or low temperatures as acute risk factors can be identified by changing body temperatures (Nowak, 2010). A JEM can be used to draw conclusions about internal exposure (Pearce and Douwes, 2008).
	psychological	Several instruments (questionnaires and observational instruments) are available to measure psychosocial factors in the work environment (Tabanelli et al., 2008). Job stress surveys or specific scales are developed (Levy et al., 2011). Medical history, interviews and employee surveys may give indications to possible psychological exposures during work (Nowak, 2010). Specific psychological exposures can be identified by using JEMs (Pearce and Douwes, 2008).
cultural factors	SES	

(continued on next page)

Table 5 (continued)

Stressor group	Stressor	Opportunities
		Censuses, surveys and national data registers can provide data on SES of population, using a variety of measures at an individual (e.g. personal education or occupation, (Galobardes et al., 2006a; Galobardes et al., 2006b) household (e.g. Family Affluence Scale (FAS) (Boyce et al., 2006)) or neighbourhood scale (e.g. English Index of Multiple Deprivation, IMD (Smith et al., 2015)). The relative importance of measures of socioeconomic status varies with the location of the study so an understanding of local relevant socioeconomic status measures is paramount. In other international studies the availability of similar SES measures may be critical. Furthermore different manifestations of socioeconomic status may be relevant for different health outcomes but other studies have found the strongest relationships between housing tenure and heart disease (Woodward et al., 1992). For example elsewhere in HEALS we are using car access as a measure of socioeconomic status because it is central to time activity and exposure to air pollution. Such sources often include measures of medical conditions, mental health and wellbeing including self-report, health service records and a few surveys such as national health surveys may include biological samples (Brummett et al., 2013) Note that socioeconomic status itself may not be intrinsically linked to health. Instead it may be a marker for other exposures such as tobacco smoke or poor diet (Giesinger et al., 2014) or poor housing (Gibson et al., 2011). Furthermore the length of exposure to low SES or low SES in childhood may be more important than current SES for some diseases (Giesinger et al., 2014).
	alcohol consumption	Acute alcohol consumption can be measured in blood (BAC: blood alcohol content), urine etc. (see Table 2). The AUDIT (Alcohol Use Disorders Identification Test) questionnaire can be used to collect data about alcohol consumption (Babor et al., 2001).
	drug consumption	Besides blood or urine analyses of the substance of interest, questionnaires (such as CRAFFT or DAST – 10 from US National Institute on Drug Abuse) are useful to collect data about the internal exposure of drugs.
	nutritional status	Anthropometric, clinical, biochemical (according nutrient to be evaluated: water-soluble vitamin, fat-soluble vitamins and nutrients, trace elements, Isoflavones and Lignans; Hepatic proteins, Hormones, Nitrogen in urine) and dietary evaluation (Food Frequency Questionnaires (FFQ) such as EPIC-Norfolk and Food4Me) methods (Blössner and de Onis, 2005; CDC, 2012; Wasantwisut and Neufeld, 2012).
	physical activity	Biomarkers of physical activity and exercise are the following: Cortisol and testosterone for chronic stress and fatigue; lactate, c(CPK), creatinine, ammonia, lactate dehydrogenase (LDH), uric acid and urea are markers of overtraining; C-reactive protein (CRP), interleukin – 6 (IL – 6) and leukocytes are markers of inflammation associated to physical activity (Palacios et al. 2015). The most used biomarkers to muscle fatigue are cortisol, lactate and IL – 6 and moreover ammonia, leukocytes and oxidative stress parameters are being increasingly used (Palacios et al., 2015). Reactions of organism are measurable like increased inflammation biomarkers (Margeli et al., 2005). Global Physical Activity and International Physical Activity Questionnaires (GPAQ and IPAQ) (Craig et al. 2003; Bull et al. 2009).
	consumer products	Analytical determination of endocrine disruptors and chemicals of concern contained in consumer products in biological samples are reported in several publications (Faniband et al., 2014). Use frequency and life style questionnaires. Due the complexity to analyze all the activities, consumer products, and chemicals containing in them, and the different routes of exposures the use of biomarkers of exposure to the chemicals contained in consumer products seems to be a more reasonable way to assess the exposure to this confounder (WHO, 2006).
	stress	

(continued on next page)

Table 5 (continued)

Stressor group	Opportunities
Stressor	
	<p>Stress enhances cortisol in blood what can be measured (Kingston et al., 2012). Although a broad range of instruments is available to assess psychological stress, there is no measure that is appropriate for all the aspects of stress (e.g. occupational stress, anxiety, depression, daily hassles, life events, socio-environmental stressors) and for all populations (children, adolescents, adults, pregnant and postpartum women). The exact stress measure that one may choose depends on the question that is being posed (Nast et al., 2013). Questionnaires/scales are usually validated and their psychometric value is proven but the core challenge is the choice of a proper tool (Kingston et al., 2012). Exemplary instruments: Perceived Stress Scale (PSS) (Cohen et al., 1983), State-trait Anxiety Inventory (STA) (Spielberger et al., 1983), Social Readjustment Rating Scale (Holmes and Rahe, 1967), APGAR (Adaptation, Partnership, Growth, Affection, Resolve) Family Scale (Smilkstein, 1993; Smilkstein et al., 1982).</p>

Abbreviations: APGAR, adaptation, partnership, growth, affection, resolve; AUDIT, alcohol use disorders identification Test; BAC, blood alcohol content; CRP, C-reactive protein; dB(A), decibel; DNA, deoxyribonucleic acid; ECO, expired carbon-monoxide; EMF, electromagnetic fields; FAS, family affluence scale; FFQ, food frequency questionnaires; GPAQ, global physical activity questionnaires; IL-6, interleukin-6; IMD, index of multiple deprivation; IPAQ, international physical activity questionnaires; JEM, job-exposure-matrix; LDH, lactate dehydrogenase; PSS, perceived stress scale; STA, state-trait anxiety inventory.

Abbreviations of stressor groups are explained in the list of abbreviations at the end of the manuscript.

biomonitoring. Although possible ways of representing the aggregate exposure of some stressors without specific BoEs were found (see Table 5), lack of specificity introduces uncertainties in using these to unravel the exposome. As the characteristics of environmental stressors may be very diverse, HBM measurements need to be complemented by tools and technologies that would allow effective HBM data assimilation (Sarigiannis et al., 2014) to accurately relate HBM values to actual human exposure to potential health stressors. This includes an array of technologies, employing environmental monitoring or food item analysis for chemical residuals, or ancillary exposure information retrieved from questionnaires or exposure related databases.

Currently exposure limit values used in chemical safety regulations are derived for the most part on toxicological (i.e. hazard-based) considerations using animal models and extrapolated to human exposure limit values with corrections using assessment factors that pertain to intra-species differences and inter-species variability. Given the cost and the burden to derive such acceptable limit values, they tend to be identified only for a limited number of chemicals and for an even more limited number of primary metabolites. In addition, the lack of harmonization among the various cohort and human biomonitoring studies results in a paucity of widely accepted exposure limit values based on HBM data. Most of these studies are designed to answer specific questions of limited scope, which are mostly related to the quantification of exposure levels among the study participants.

In order to derive exposure limit values, exposure characterization and quantification have to be associated with health observations. In addition, the methods used for interpretation of exposure-to-health associations, including both the statistical methods employed and proper consideration of potential modifiers (genetics, dietary, socio-economic conditions), are hardly consistent among the studies performed thus far. Several of these issues are addressed in HEALS, and they are to be addressed in the European Human Biomonitoring Initiative (HBM4EU project) (Ganzleben et al., 2017). Nonetheless, it should be noted that BEs do not address shortcomings in the derivation of current regulatory guidelines. They simply provide estimates of the urinary concentrations corresponding to the regulatory exposure limit as per the respective safety regulation.

Thus, beyond the difficulties in deriving BEs for given exposure reference values, a major problem is the lack of properly defined exposure-based limit and/or reference values. Widespread use of PBBK models will facilitate the derivation of BEs and will support the

derivation of more robust associations between external exposures and biomonitoring data. Moreover, this will also allow the use of rapidly produced and inexpensive *in vitro* reference values, such as the ones derived by US EPA's Toxicity Forecaster (ToxCast) (US EPA, 2016). In this case, for calculating a BE, the starting point will be the biological pathway altering dose (BPAD) instead of an animal based POD. BPAD is analogous to current risk assessment metrics in that it combines dose-response data with analysis of uncertainty and population variability so as to derive exposure limits (Judson et al., 2010, 2011). The analogy is closest when perturbation of a pathway is a key event in the mode of action (MOA) leading to a specified adverse outcome. An application of this method (use of BPAD for deriving human BE) has been showcased by Sarigiannis et al. (2016) for bisphenol A. Methods for deriving BEs that are based on *in vitro* systems will allow the faster screening of newly produced chemicals, considering the rapidity, the lower costs and the lack of ethical concerns which occur when animal studies are used to derive PODs.

For EWAS, it is essential to consider a large range of diverse environmental stressors to enable the most complete decoding of the exposome. Relying on only one monitoring method (in this work we refer to biomonitoring) is insufficient. Although analysis of human bio-samples for identifying the BoE levels is a good starting point, further elucidation of the individual exposome requires the use of additional molecular analysis such as transcriptomics, metabolomics or adductomics according to the HEALS paradigm; in turn, this requires additional computational tools that have to be used to interpret the biomonitoring and multi-omics results in the frame of a more integrative approach. This is actually one of the key aspects investigated in HEALS.

4.1. Limitations and strengths

Despite the amount of information collected in this narrative review, this work has limitations. Information was collected in an expert-driven, distributed, narrative review process which might involve individual researcher decisions. The internal review process reduced this potential researcher bias. The list of stressors included is not exhaustive but evaluated based on the joint opinion of the participating partners as a list of important stressors for the population in the EU. Completeness of the list of stressors is impossible because of the countless number of stressors available and the constant production and release of new

chemicals. This is the case of some substitutes, such as other bisphenols for BPA (e.g. BPF, BPS) (Chen et al., 2016) or non-phthalate plasticizers like DINCH (diisononyl cyclohexane-1,2-dicarboxylate) or DEHT (di(2-ethylhexyl) terephthalate) (Fromme et al., 2016; Larsson et al., 2017). There are also many other BoE for POPs, including brominated flame retardants such as TBBPA (tetrabromobisphenol A) (Lu et al., 2017) or PBB (polybrominated biphenyls) (Ploteau et al., 2016). Other examples are organic compounds like glyphosate-based herbicides (Conrad et al., 2017), carbamates (Haines et al., 2017), or micro- as well as macro-nutrients that have not been included. While vitamin C and folate are included in our work as examples, these and further micronutrients (e.g. iodine or other trace elements, proteins, water or fat soluble vitamins) with health-relevance (e.g., deficiencies) have been discussed previously in another review (Combs et al., 2013).

The lists of reference values, exposure limit values and biomonitoring equivalents were not intended to be complete; rather, examples are listed to provide an inside in the interpretation of data. Presented are rather condensed information and stratifications by age, gender, or other subgroups were not reported. HBM itself contains limitations such as the use of diverse methods for analyses. Also, the derivation of reference and exposure limit values is based on expert decisions usually on the basis of a consensus process.

This paper's scope lies in the availability of BoEs and does not include further technical information. For example, it must be kept in mind that the half-life of BoEs is an essential piece of information for the practical use of BoEs. For example, as the biological half-life of nicotine is ~2 h (h), cotinine, the major metabolite of nicotine, with a half-life of 17 h, is a more suitable BoE (Benowitz, 1996). For biomarkers with a half-life of less than 2 h, biomonitoring is not feasible. When the half-life between 2 and 10 h, a sample collected at the end of the day reflects the exposure over the day, while with half-lives of 10–100 h, the optimal sampling time is at the end of the week, and the results reflect exposure during the preceding few days (HSE, 1992). The half-life of the marker of choice is a key parameter to be taken into account to achieve representative spot sampling results.

In addition the intraclass correlation coefficient (ICC) is important if spot-measurements are intended to estimate long-term exposures. The ICC is a value between 0 and 1 and if the value is close to 0 then repeated measurements of the BoE from the same individual would result in any test result, whereas if the ICC is close to 1 repeated measurements would be very similar (Pleil and Sobus, 2013).

Also, information about the representativeness of the samples is not included in this paper. For example, the reference values for 3PBA, diethyl phosphate, DMP, DMTP, PCP, PFOA, and PFOS are based on not representative samples of the population in Germany, as underlined by the authors (HBM-UBA, 2003; Heudorf et al., 2006; Schulz et al., 2011). In contrast, the data from the French National Survey on Nutrition and Health (InVS, 2010) and the German Environmental Survey on Children (Schulz et al., 2012) as well as the reference values for PBDEs (Gari and Grimalt, 2013) and β -HCH (Gari et al., 2014) were derived based on representative samples.

If no other reference value (e.g., RV_{95}) was available, information was included on measures of central tendency (MCT). In several cases, the presented MCTs represent arithmetic mean, geometric mean, median, or a mixture of them. It needs to be mentioned, that the mean (arithmetic mean, average) should be avoided in HBM studies, since the distribution of the values do not follow a normal distribution. Thus, mean values do not represent the central tendency.

Strengths of this work are the broad inclusion of diverse environmental stressors, the extensive list of BoEs and corresponding reference values, exposure limit values and biomonitoring equivalents as well as the inclusion of possibilities to measure the internal exposure of stressors without specific BoE.

5. Conclusions

Given the diversity of environmental stressors that need to be examined to unravel the exposome, current-day human biomonitoring is suitable for determining the internal exposome of several stressors (e.g., metals, PCBs, VOCs) but not for many others (e.g., NO_x , PM, physical activity). Most chemical and biological stressors are measurable in human specimens whereas exposure to the majority of physical, social and psychological stressors needs to be assessed using methods complementary to HBM. The joint and harmonized application of methods and tools to unravel the exposome represents the main task of the HEALS project.

Acknowledgments

This work has received funding from the European Union's Seventh Programme for research, technological development and demonstration under grant agreement No. 603946 (Health and Environment-wide Associations based on Large population Surveys, HEALS). All HEALS partners involved in the discussion of deliverable 4.2 during the annual HEALS meeting in Edinburgh, UK, on September 16, 2014, are gratefully acknowledged. Special thanks go to our colleagues outside WP4 and outside HEALS who reviewed chapters of their expertise: Thank you, Tomislav Bituh, Danijela Štimac, and Matthias Weigl.

Declaration of interest

All authors received funding from the European Union's Seventh Programme for research, technological development and demonstration under grant agreement No. 603946 (HEALS).

Appendices

The complete HEALS report can be downloaded from the HEALS website, http://www.heals-eu.eu/wp-content/uploads/2013/08/HEALS_D4.2.pdf.

References

- Andrianou, X.D., et al., 2014. Spatial and seasonal variability of urinary trihalomethanes concentrations in urban settings. *Environ. Res.* 135, 289–295.
- Andrianou, X.D., et al., 2016. Human Exposures to Bisphenol A, Bisphenol F and Chlorinated Bisphenol A Derivatives and Thyroid Function. *PLoS One* 11, e0155237.
- Angerer, J., et al., 2011. Human biomonitoring assessment values: approaches and data requirements. *Int. J. Hyg. Environ. Health* 214, 348–360.
- Angerer, J., et al., 2007. Human biomonitoring: state of the art. *Int. J. Hyg. Environ. Health* 210, 201–228.
- ANSES, 2013. What are the critical blood concentration levels for PCBs? Agence nationale de sécurité sanitaire Alimentation Environnement Travail (ANSES), Maisons-Alfort.
- Aprea, C., et al., 2000. Biologic monitoring of exposure to organophosphorus pesticides in 195 Italian children. *Environ. Health Perspect.* 108, 521–525.
- Arnold, S.M., et al., 2013. The use of biomonitoring data in exposure and human health risk assessment: benzene case study. *Crit. Rev. Toxicol.* 43, 119–153.
- Benowitz, N.L., 1996. Cotinine as a biomarker of environmental tobacco smoke exposure. *Epidemiol. Rev.* 18, 188–204.
- Aylward, L.L., et al., 2009a. Derivation of Biomonitoring Equivalents for di-n-butyl phthalate (DBP), benzylbutyl phthalate (BzBP), and diethyl phthalate (DEP). *Regul. Toxicol. Pharmacol.* 55, 259–267.
- Aylward, L.L., et al., 2009b. Derivation of Biomonitoring Equivalents for di(2-ethylhexyl) phthalate (CAS No. 117-81-7). *Regul. Toxicol. Pharmacol.* 55, 249–258.
- Aylward, L.L., et al., 2013. Evaluation of biomonitoring data from the CDC National Exposure Report in a risk assessment context: perspectives across chemicals. *Environ. Health Perspect.* 121, 287–294.
- Barr, D.B., et al., 2014. Concentrations of dialkyl phosphate metabolites of organophosphorus pesticides in the U.S. population. *Environ. Health Perspect.* 122, 189–200.
- Becker, K., et al., 2008. German Environmental Survey for Children 2003/06 – GerES IV – Human Biomonitoring Levels of selected substances in blood and urine of children in Germany. Research Report 202 62 219. UBA-FB 001026 WaBoLu-Hefte. Federal Environment Agency (Umweltbundesamt, UBA), Dessau-Roßlau, Berlin, pp. 1–85.
- Beirao, F., Araujo, R., 2013. State of the art diagnostic of mold diseases: a practical guide for clinicians. *Eur. J. Clin. Microbiol. Infect. Dis.* 32, 3–9.
- Boogaard, P.J., et al., 2011. Human biomonitoring as a pragmatic tool to support health risk management of chemicals-examples under the EU REACH programme. *Regul. Toxicol. Pharmacol.* 59, 125–132.

- Brugnone, F., et al., 1993. Blood styrene concentrations in a "normal" population and in exposed workers 16 hours after the end of the workshift. *Int. Arch. Occup. Environ. Health* 65, 125–130.
- Burtis, C., et al., 2012. Tietz Textbook of Clinical Chemistry and Molecular Diagnostics, 5th ed. Saunders, Missouri.
- Calafat, A.M., 2010. Urinary concentrations of four parabens in the U.S. population: NHANES 2005–2006. *Environ. Health Perspect.* 118, 679–685.
- Callcut, R.A., Branson, R.D., 2009. How to read a review paper. *Respir. Care* 54, 1379–1385.
- CDC, 2005. Third National Report on Human Exposure to Environmental Chemicals. Centers for Disease Control and Prevention (CDC), Atlanta, Georgia.
- CDC, 2017. Fourth National Report on Human Exposure to Environmental Chemicals. Updated Tables, January 2017, Volume One. Centers for Disease Control and Prevention (CDC), Atlanta, Georgia.
- Chen, D., et al., 2016. Bisphenol Analogues Other Than BPA: environmental Occurrence, Human Exposure, and Toxicity-A Review. *Environ. Sci. Technol.* 50, 5438–5453.
- Chovanova, J., et al., 2012. PCDD/PCDF, dl-PCB and PBDE serum levels of Slovak general population. *Chemosphere* 88, 1383–1389.
- Combs Jr., G.F., et al., 2013. Biomarkers in nutrition: new frontiers in research and application. *Ann. N. Y. Acad. Sci.* 1278, 1–10.
- Conrad, A., et al., 2017. Glyphosate in German adults - Time trend (2001 to 2015) of human exposure to a widely used herbicide. *Int. J. Hyg. Environ. Health* 220, 8–16.
- Cook, D.J., et al., 1997. Systematic reviews: synthesis of best evidence for clinical decisions. *Ann. Intern. Med.* 126, 376–380.
- Costopoulou, D., et al., 2006. Levels of dioxins, furans and PCBs in human serum and milk of people living in Greece. *Chemosphere* 65, 1462–1469.
- DFG, 2016. List of MAK and BAT Values 2016: maximum Concentrations and Biological Tolerance Values at the Workplace (Permanent Senate Commission for the Investigation of Health Hazards of Chemical Compounds in the Work Area. Vol. Report 52). Deutsche Forschungsgemeinschaft (DFG), Weinheim.
- Covaci, A., et al., 2015. Urinary BPA measurements in children and mothers from six European member states: Overall results and determinants of exposure. *Environ. Res.* 141, 77–85.
- Dekant, W., Volkel, W., 2008. Human exposure to bisphenol A by biomonitoring: methods, results and assessment of environmental exposures. *Toxicol. Appl. Pharmacol.* 228, 114–134.
- Den Hond, E., et al., 2015. First steps toward harmonized human biomonitoring in Europe: demonstration project to perform human biomonitoring on a European scale. *Environ. Health Perspect.* 123, 255–263.
- EFSA, 2014. Scientific Opinion on Dietary Reference Values for folate. *EFSA J.* 12, 3893.
- El-Farhan, N., et al., 2017. Measuring cortisol in serum, urine and saliva – are our assays good enough? *Ann. Clin. Biochem.* 54, 308–322.
- Fromme, H., et al., 2016. Non-phthalate plasticizers in German daycare centers and human biomonitoring of DINCH metabolites in children attending the centers (LUPE 3). *Int. J. Hyg. Environ. Health* 219, 33–39.
- Fustinoni, S., et al., 2010. Urinary BTEX, MTBE and naphthalene as biomarkers to gain environmental exposure profiles of the general population. *Sci. Total Environ.* 408, 2840–2849.
- Ganzleben, C., et al., 2017. Human biomonitoring as a tool to support chemicals regulation in the European Union. *Int. J. Hyg. Environ. Health* 220, 94–97.
- Gari, M., Grimalt, J.O., 2013. Inverse age-dependent accumulation of decabromodiphenyl ether and other PBDEs in serum from a general adult population. *Environ. Int.* 54, 119–127.
- Gari, M., et al., 2014. Impacts of atmospheric chlor-alkali factory emissions in surrounding populations. *Environ. Int.* 65, 1–8.
- Haines, D.A., et al., 2017. An overview of human biomonitoring of environmental chemicals in the Canadian Health Measures Survey: 2007–2019. *Int. J. Hyg. Environ. Health* 220, 13–28.
- Grainger, J., et al., 2006. Reference range levels of polycyclic aromatic hydrocarbons in the US population by measurement of urinary monohydroxy metabolites. *Environ. Res.* 100, 394–423.
- Gratz, T., Kevekordes, S., 2001. [Perchloroethylene (PER)] Perchloroethylen (PER). In: Böse-O'Reilly, S. (Ed.), Leitfaden Umweltmedizin. Urban & Fischer. München, Jena, pp. 373–375.
- Hays, S.M., et al., 2009. Derivation of Biomonitoring Equivalents for cyfluthrin. *Regul. Toxicol. Pharmacol.* 55, 268–275.
- Hays, S.M., et al., 2012. Biomonitoring Equivalents for benzene. *Regul. Toxicol. Pharmacol.* 62, 62–73.
- Hirschhorn, J.N., Daly, M.J., 2005. Genome-wide association studies for common diseases and complex traits. *Nat. Rev. Genet.* 6, 95–108.
- HBM-UBA, 2003. Innere Belastung der Allgemeinbevölkerung in Deutschland mit Organophosphaten und Referenzwerte für die Organophosphatmetabolite DMP, DMTP und DEP im Urin. Stellungnahme der Kommission „Human-Biomonitoring“ des Umweltbundesamtes. Bundesgesundheitsblatt – Gesundheitsforschung – Gesundheitsschutz. 46, 1107–1111.
- HSE, 1992. The Workplace (Health, Safety and Welfare) Regulations 1992. Health and Safety Executive (HSE), London.
- Health Canada, 2013. Second Report on Human Biomonitoring of Environmental Chemicals in Canada. Results of the Canadian Health Measures Survey Cycle 2 (2009–2011).
- Heudorf, U., et al., 2006. Reference values for metabolites of pyrethroid and organophosphorous insecticides in urine for human biomonitoring in environmental medicine. *Int. J. Hyg. Environ. Health* 209, 293–299.
- Heyndrickx, E., et al., 2015. Human biomonitoring of multiple mycotoxins in the Belgian population: Results of the BIOMYCO study. *Environ. Int.* 84, 82–89.
- Judson, R.S., et al., 2010. In vitro screening of environmental chemicals for targeted testing prioritization: the ToxCast project. *Environ. Health Perspect.* 118, 485–492.
- IEH, 2008. Background Incidence of Key Biomarkers of Chemical Exposure within the General UK Population. Institute of Environment and Health (IEH), Cranfield.
- InVS, 2010. Exposure of the French population to environmental pollutants French Institute for Public Health Surveillance (InVS), Saint-Maurice Cedex.
- Jeanneret, F., et al., 2014. Human urinary biomarkers of dioxin exposure: analysis by metabolomics and biologically driven data dimensionality reduction. *Toxicol. Lett.* 230, 234–243.
- Judson, R.S., et al., 2011. Estimating toxicity-related biological pathway altering doses for high-throughput chemical risk assessment. *Chem. Res. Toxicol.* 24, 451–462.
- Kazlauskaitė, R., et al., 2008. Corticotropin tests for hypothalamic-pituitary-adrenal insufficiency: a metaanalysis. *J. Clin. Endocrinol. Metab.* 93, 4245–4253.
- Kim, S., 2016. Overview of Cotinine Cutoff Values for Smoking Status Classification. *Int. J. Environ. Res. Public Health* 13.
- Koch, H., Angerer, J., 2012. Phthalates: Biomarkers and Human Biomonitoring. In: Knudsen, L., Merlo, D. (Eds.), Biomarkers and Human Biomonitoring, Volume 1: Ongoing Programs and Exposures. Royal Society of Chemistry, Cambridge, pp. 179–233.
- Krishnan, K., et al., 2010a. Biomonitoring Equivalents for bisphenol A (BPA). *Regul. Toxicol. Pharmacol.* 58, 18–24.
- Krishnan, K., et al., 2010b. Biomonitoring Equivalents for triclosan. *Regul. Toxicol. Pharmacol.* 58, 10–17.
- Krishnan, K., et al., 2011. Biomonitoring equivalents for 2,2',4,4',5-pentabromodiphenylether (PBDE-99). *Regul. Toxicol. Pharmacol.* 60, 165–171.
- LaKind, J.S., et al., 2010. Public health interpretation of trihalomethane blood levels in the United States: NHANES 1999–2004. *J. Expo. Sci. Environ. Epidemiol.* 20, 255–262.
- Larsson, K., et al., 2017. Phthalates, non-phthalate plasticizers and bisphenols in Swedish preschool dust in relation to children's exposure. *Environ. Int.* 102, 114–124.
- Liljendahl, T.S., et al., 2013. Urinary levels of thymine dimer as a biomarker of exposure to ultraviolet radiation in humans during outdoor activities in the summer. *Mutagenesis* 28, 249–256.
- Lu, D., et al., 2017. Multi-analyte method development for analysis of brominated flame retardants (BFRs) and PBDE metabolites in human serum. *Anal. Bioanal. Chem.* 409, 5307–5317.
- Lupsa, I.R., et al., 2015. Urinary cotinine levels and environmental tobacco smoke in mothers and children of Romania, Portugal and Poland within the European human biomonitoring pilot study. *Environ. Res.* 141, 106–117.
- Minoia, C., et al., 1996. Environmental and urinary reference values as markers of exposure to hydrocarbons in urban areas. *Sci. Total Environ.* 192, 163–182.
- Mocarelli, P., et al., 1986. Clinical laboratory manifestations of exposure to dioxin in children. A six-year study of the effects of an environmental disaster near Seveso, Italy. *Jama* 256, 2687–2695.
- Morgan, M.S., 1997. The biological exposure indices: a key component in protecting workers from toxic chemicals. *Environ. Health Perspect.* 105 (Suppl 1), 105–115.
- Northridge, M.E., et al., 1999. Diesel exhaust exposure among adolescents in Harlem: a community-driven study. *Am. J. Public Health* 89, 998–1002.
- NRC, 1987. Biological markers in environmental health research. Committee on Biological Markers of the National Research Council. *Environ. Health Perspect.* 74, 3–9.
- Päpke, O., et al., 2011. Chapter 3C: Biomarkers of Exposure: Polychlorinated Dibenzo-p-dioxins and Polychlorinated Dibenzofuranes. In: Knudsen, L., Merlo, D.F. (Eds.), Biomarkers and Human Biomonitoring: Volume 1.
- Patel, C.J., et al., 2010. An Environment-Wide Association Study (EWAS) on type 2 diabetes mellitus. *PLoS One* 5, e10746.
- Pignatelli, B., 1989. Group-selective determination of total N-nitroso compounds in nitrate-containing human urine samples. *Analyst* 114, 1103–1108.
- Pleil, J.D., Sobus, J.R., 2013. Estimating lifetime risk from spot biomarker data and intraclass correlation coefficients (ICC). *J. Toxicol. Environ. Health A* 76, 747–766.
- Ploteau, S., et al., 2016. Distribution of persistent organic pollutants in serum, omental, and parietal adipose tissue of French women with deep infiltrating endometriosis and circulating versus stored ratio as new marker of exposure. *Environ. Int.* 97, 125–136.
- Poulsen, O.M., et al., 1997. A supplement to the approved IFCC recommendation on the theory of reference values. *Pure Appl. Chem.* 69, 1601–1611.
- Preuss, R., et al., 2004. Pilot study on the naphthalene exposure of German adults and children by means of urinary 1- and 2-naphthol levels. *Int. J. Hyg. Environ. Health* 207, 441–445.
- Rappaport, S.M., 2011. Implications of the exposome for exposure science. *J. Expo. Sci. Environ. Epidemiol.* 21, 5–9.
- Roca, M., et al., 2014. Biomonitoring exposure assessment to contemporary pesticides in a school children population of Spain. *Environ. Res.* 131, 77–85.
- Sarigiannis, D., et al., 2014. Integra: From global scale contamination to tissue dose. Proceedings - 7th International Congress on Environmental Modelling and Software: Bold Visions for Environmental Modeling, iEMSS 2014, Vol. 2, pp. 1001–1008.
- Sarigiannis, D.A., et al., 2016. Integrated exposure and risk characterization of bisphenol-A in Europe. *Food Chem. Toxicol.* 98, 134–147.
- Saurat, J.H., et al., 2012. The cutaneous lesions of dioxin exposure: lessons from the poisoning of Victor Yushchenko. *Toxicol. Sci.* 125, 310–317.
- Schulz, C., et al., 2007. The German human biomonitoring commission. *Int. J. Hyg. Environ. Health* 210, 373–382.
- Schettgen, T., et al., 2015. Current data on the background burden to the persistent organochlorine pollutants HCB, p,p'-DDE as well as PCB 138, PCB 153 and PCB 180 in plasma of the general population in Germany. *Int. J. Hyg. Environ. Health* 218, 380–385.
- Schulz, C., et al., 2009. Revised and new reference values for environmental pollutants in urine or blood of children in Germany derived from the German environmental

- survey on children 2003-2006 (GerES IV). *Int. J. Hyg. Environ. Health* 212, 637–647.
- Schulz, C., et al., 2011. Update of the reference and HBM values derived by the German Human Biomonitoring Commission. *Int. J. Hyg. Environ. Health* 215, 26–35.
- Schoeters, G., et al., 2016. Three cycles of human biomonitoring in Flanders - Time trends observed in the Flemish Environment and Health Study. *Int. J. Hyg. Environ. Health*.
- Scholz, H., 2001a. [Pentachlorophenol (PCP)] Pentachlorophenol (PCP). In: Böse-O'Reilly, S. (Ed.), *Leitfaden Umweltmedizin. Urban & Fischer, München, Jena*, pp. 369–373.
- Scholz, H., 2001b. [Toluol (C7H8)] Toluene (C7H8). In: Böse-O'Reilly, S. (Ed.), *Leitfaden Umweltmedizin. Urban & Fischer, München, Jena*, pp. 398–400.
- Scholz, H., 2001c. [Xylol] Xylene. In: Böse-O'Reilly, S. (Ed.), *Leitfaden Umweltmedizin. Urban & Fischer, München, Jena*, pp. 401–403.
- Schulz, C., et al., 2012. Reprint of "Update of the reference and HBM values derived by the German Human Biomonitoring Commission". *Int. J. Hyg. Environ. Health* 215, 150–158.
- Schwartz, D., Collins, F., 2007. Medicine. *Environmental biology and human disease. Science* 316, 695–696.
- Siqueira, M.E., Paiva, M.J., 2002. Hippuric acid in urine: reference values. *Rev. Saude Publica* 36, 723–727.
- Smith, K.R., et al., 1999. How much global ill health is attributable to environmental factors? *Epidemiology* 10, 573–584.
- Sochorova, L., et al., 2017. Perfluorinated alkylated substances and brominated flame retardants in serum of the Czech adult population. *Int. J. Hyg. Environ. Health* 220, 235–243.
- Solfrizzo, M., et al., 2014. Assessment of multi-mycotoxin exposure in southern Italy by urinary multi-biomarker determination. *Toxins (Basel)* 6, 523–538.
- Stahl, T., et al., 2011. Toxicology of perfluorinated compounds. *Environ. Sci. Eur.* 23, 1–52.
- Tsangari, X., et al., 2017. Spatial characteristics of urinary BTEX concentrations in the general population. *Chemosphere* 173, 261–266.
- UBA, 2017. [Environmental specimen bank] Umweltprobenbank des Bundes [Internet] German Environment Agency (Umweltbundesamt, UBA), Dessau-Roßlau, Berlin.
- US EPA, 2016. Toxicity Forecasting. Advancing the Next Generation of Chemical Evaluation. United States Environmental Protection Agency (US EPA), Washington D.C.
- Valcke, M., et al., 2006. Biological monitoring of exposure to organophosphate pesticides in children living in peri-urban areas of the Province of Quebec. Canada. *Int. Arch. Occup. Environ. Health* 79, 568–577.
- Van Duursen, M., 2010. CYP1A1 induction in Lymphocytes from mice and humans: A biomarker of Dioxin exposure? *Organohalogen Compd.* 72, 1038–1041.
- Wild, C.P., 2005. Complementing the genome with an "exposome": the outstanding challenge of environmental exposure measurement in molecular epidemiology. *Cancer Epidemiol. Biomark. Prev.* 14, 1847–1850.
- Vassiliadou, I., et al., 2010. Levels of perfluorooctanesulfonate (PFOS) and perfluorooctanoate (PFOA) in blood samples from different groups of adults living in Greece. *Chemosphere* 80, 1199–1206.
- Wild, C.P., 2012. The exposome: from concept to utility. *Int. J. Epidemiol.* 41, 24–32.
- Wei, B., et al., 2016. Assessing exposure to tobacco-specific carcinogen NNK using its urinary metabolite NNAL measured in US population: 2011-2012. *J. Expo. Sci. Environ. Epidemiol.* 26, 249–256.
- Wilhelm, M., et al., 2003. Revised and new reference values for some persistent organic pollutants (POPs) in blood for human biomonitoring in environmental medicine. *Int. J. Hyg. Environ. Health* 206, 223–229.
- Yike, I., et al., 2006. Mycotoxin adducts on human serum albumin: biomarkers of exposure to *Stachybotrys chartarum*. *Environ. Health Perspect.* 114, 1221–1226.
- Zografos, G.N., et al., 2010. Primary pigmented nodular adrenocortical disease presenting with a unilateral adrenocortical nodule treated with bilateral laparoscopic adrenalectomy: a case report. *J. Med. Case Rep.* 4, 230.