

**Royal  
Geographical  
Society**

with IBG

Advancing geography  
and geographical learning

**WILEY**

---

The Permanence of Ocean Basins

Author(s): Hugh Robert Mill

Source: *The Geographical Journal*, Vol. 1, No. 3 (Mar., 1893), pp. 230-234

Published by: geographicalj

Stable URL: <http://www.jstor.org/stable/1773821>

Accessed: 28-06-2016 09:21 UTC

---

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at

<http://about.jstor.org/terms>

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).



*The Royal Geographical Society (with the Institute of British Geographers), Wiley* are collaborating with JSTOR to digitize, preserve and extend access to *The Geographical Journal*

range to the south, crossing numerous spurs. On the following day he crossed the watershed at an elevation of 10,000 feet, and reached the fertile district of Imbo, where he saw the broad plain of the Rufizi lying beneath him. Villages were numerous. The oil-palm and grey parrots were again seen, and many of the natives were dressed in cotton, those of northern Urundi and Ruanda having contented themselves with skins and bark-cloth. On September 25th, the traveller reached one of the stations of Rumaliza, in Uzige, at the northern end of Tanganyika. The Warundi in this district, fish, till the soil, and export palm-oil, but the Arabs principally deal in slaves.

Dr. Baumann then travelled in a south-easterly direction to Tabora, where he arrived on November 7th. Having climbed the steep mountains which shut in the lake on the east, he came upon a fine pastoral region, sloping to the eastward, and occupied by warlike Watusi. Here he crossed the Muvarazi and the Luviroza, tributaries of the Ruvuvu and consequently the most southerly head-streams of the Nile basin. Stony ranges of mountains form the watershed between the Ruvuvu and the Mlagarazi. The last village of the Warundi, built on the branches of trees, stands on the latter. Uhha, which lies beyond, has open forests of great extent in which grow acacias and a tree resembling our beech. The inhabitants are kinsmen of the Warundi. They are governed by petty chiefs and live in small villages. All their cattle were stolen a few years ago by the Masai. Crossing the Mlagarazi once more and passing through Muhambwa and Mulunga, both districts of Uhha, Dr. Baumann arrived in Kirambo, which is inhabited by Wasumbwa, a division of the Wanyamwezi. He then crossed the Igombe River, passed through Urambo, now ruled over by Tugamoto, a son of the famous Mirambo, visited Mr. and Mrs. Shaw at Kilimani Urambo, and ultimately reached Tabora. During the whole of this expedition Dr. Baumann only lost five men, one of whom was killed by the Watusi. On the other hand there were "numerous" births among the wives of the elephant hunters who accompanied him.

---

## THE PERMANENCE OF OCEAN BASINS.

By HUGH ROBERT MILL, D.Sc., F.R.S.E.

AN interesting discussion has recently taken place in the pages of *Natural Science* upon the question of the relative permanence of oceanic and continental areas. Dr. A. Russel Wallace comments on the remarkable convergence of independent lines of research which led Professor Dana, Mr. Darwin, Sir Archibald Geikie, Dr. John Murray, the Rev. O. Fisher, and himself to the same conclusion, that continents and oceans have not changed places in geological times. He adduces

three new arguments in favour of these views:—(1) The enormous disproportion between the mean height of the land and the mean depth of the ocean, which would render it very difficult for new land to reach the surface till long after the total submergence of the sinking continent; (2) the wonderful uniformity of level over by far the greater part of the ocean floor, which indicates that it is not subject to the same disturbing agencies which throughout all geological time have been creating irregularities in the land surface—irregularities which would be far greater than they are were they not continually counteracted by the lowering and equalising effects of subaerial denudation; (3) the remarkable parallelism and completeness of the series of geological formations in all the best-known continents and larger continental islands, indicating that none of them has risen from the ocean floor during any era of known geological history—a conclusion enforced by the absence from any of them of that general deposit of oceanic ooze at some definite horizon, which would be at once the result and proof of any such tremendous episode in their past history.

Mr. Jukes-Browne combats the theory of the permanence of original oceanic and continental areas by pointing out that, as a matter of fact, deposits which must have been formed in very deep water have been detected in elevated land. He denies the force of the second argument quoted above on account of our comparative ignorance of the details of sub-oceanic geography, and because long-continued sedimentation must have tended to reduce the irregularities of the ocean floor, which would in any case be less than those of the continents, and, finally, because Mr. Fisher's theory of the sub-oceanic outpouring of lava might account for the uniformity of the ocean-bed. Mr. Jukes-Browne also shows how it is possible that the volume of the ocean was formerly very much less than it is now, and that in consequence of the release of water from chemical union with hot rock-material by volcanic explosions the volume has been steadily increasing. Hence, in palæozoic times, he argues that the land area may actually have been in excess of the water area, and the continental plateaux might have been much less elevated above the ocean floor than they are now—that, in fact, the elevations and depressions might not have been of the character of continent and ocean-basin at all.

Mr. Jukes-Browne's remark as to long-continued sedimentation obliterating the irregularities of the ocean floor requires proof. Dr. Murray has shown that the shallower the sea is the more rapidly does sedimentation far from land proceed, sedimentation tending to exaggerate and not to obliterate, initial differences of level.

Dr. W. T. Blanford, whose name had been mentioned in the articles as an opponent of the theory of the permanence of oceans and continents, explained that he objected to strong statements on either side, and urged great caution in forming opinions from such incomplete data as we


possess. He objected, as did Mr. Jukes-Browne, to assuming that no part of the sea-bed at a depth exceeding 1000 fathoms has ever been dry land; but Dr. Wallace explained that he took the 1000-fathom line as only an approximate boundary, and even in the first edition of 'Island Life' admitted certain exceptions. Hence his advance to 1500 or even 2000 fathoms in some cases did not affect the consistency of his views.

Professor James Geikie in his address to Section E of the British Association last year (*Proceedings* for 1892, p. 638), touched upon the interchange of elevation on the continental margin, and Dr. Blanford quotes his statement as one fairly representative of modern views on the subject. He says, "The continental plateau and the oceanic hollows have never changed places, although from time to time portions of the latter have been ridged up and added to the margins of the former, while ever and anon marginal portions of the plateau have sunk down to very considerable depths." In a paper on the "Evolution of Climate," published in the *Scottish Geographical Magazine*, vol. 6 (1890), p. 57, Professor Geikie gives a series of maps showing the approximate distribution of land and water at different geological periods, a re-statement with new data of Dana's earlier work. He understands the evolution of continents to be a transition from insular groups of land to a solid continental form, the island groups appearing upon the gradually emerging continental plateau. The diversity in opinion of students of this department of physical geography seems to be due rather to individual differences of confidence in drawing conclusions, than to any serious conflict of belief as to the bearing of ascertained facts.

The actual dividing line to be assumed between the oceanic and continental areas is obviously to a large extent arbitrary. The great ocean basins are comparatively flat-floored, the continental areas bounded by the 100-fathom line are, in a general way, roof-shaped, partly flat, partly composed of steep irregular slopes. Between these two great and definite regions there is everywhere a wall or zone of transition equally definite, forming a relatively abrupt descent. Along the slope of the entire development of this zone the difference of vertical height for a given horizontal displacement at right angles to the contour lines is greater than in any other part of the Earth, a few mountain ranges excepted. The line separating the two areas must lie somewhere on this slope. The 2000-fathom line lies near the bottom of the slope close to the ocean floor, the 1000-fathom line lies nearer the top than the middle of the transition zone.

Calculating from Dr. Murray's data, I showed in 1890 (*Proceedings*, Roy. Soc. Edin., vol. 17 (1890), p. 185), that the contour line of 1700 fathoms divided the surface of the Earth almost exactly into two equal parts, one a nearly continuous area of elevation, the other an equally continuous area of depression. This line I termed mean-sphere level, because it had the remarkable property of repre-

senting the intersection of a geoidal shell, the depression of the crust below which was equal in volume to the protuberance of the continents above it. The exact position of the line of mean-sphere level it is, of course, impossible to determine until the volumes of oceans and continents below and above sea-level have been accurately ascertained, and until sea-level is itself corrected for gravitational distortion, but the position cannot be very far removed from that assigned to it. The accompanying sketch-map on Lambert's equal-area projection serves to show graphically the two great areas of elevation and depression separated by the line of mean-sphere level. The area of depression may be termed the abysmal area, a name which Dr. Murray originally applied to the region lying at a depth greater than 1000 fathoms; the elevated half includes the dry land or continental area proper, and the transitional area or flanks of the continental plateau, at present covered with water. If any compressive stress were limited to one-half of the



area of the Earth's surface while the crust was plastic and undistorted, the surface being everywhere at mean-sphere level, a mass would be elevated over the free half to such a height as would, by its weight, balance the force of the pressure causing the depression, and if the density of the crust were uniform the depressed and elevated volumes would be the same. It seems probable that the series of changes which have evolved continents and oceanic hollows may be related to the line of mean-sphere level, the position of which corresponds with that which Dr. Wallace has been led to adopt as his boundary between oceanic and continental areas. By the adoption of this physical constant some of the objections to the theory of permanence are overcome, especially that which charges those to whom the evidence seems more complete than it does to others with a change of ground when they slightly modify an approximate and largely arbitrary limit.

The main fact, conceded by all who have studied the subject, is, that there is such a thing as the evolution of continents, the heights and hollows of the Earth's crust having become greater with the lapse of time. And all, also, concede that the present ocean basins represent regions where subsidence has predominated over elevation, while the continental area is that in which elevation has been more active than depression. Thus the actual level of the sea is an accident depending on the volume of its water and the inequalities of the crust, equal variations in which, on the hypothesis of constant volume in the ocean, may have led to very different emergence or submergence of the border areas according to the angle of the slope. At present the coast-line lies nearly mid-way on the flattest expanse of the continental margin, so that a given increase or decrease in the volume of the ocean would cover or lay bare the largest possible area of land.

---

### THE RECLAMATION OF THE ZUIDERZEE.

By Professor P. H. SCHOUTE, of Groningen.

THE question of the reclamation of land now covered by the Zuiderzee is no new one, but it is only recently that a thoroughly practicable scheme has been matured for the accomplishment of this great engineering feat. In 1886 a Committee was appointed to consider and report upon the question under the chairmanship of Mr. A. Buma, who had frequently urged this matter upon the attention of the Government. The result of the deliberations of this Committee has been published in the form of eight memoirs, in which the whole question is thoroughly discussed in all its bearings. The first memoir deals with general principles, discussing the best position and direction for the main dyke which shall shut off communication with the sea, and sketching a general plan of the engineering works to be effected. The commercial and strategical aspects of the proposed changes are also fully considered. The three succeeding memoirs are more technical in their treatment of the engineering problems. The means of protection against floods and the maintenance of navigable channels after the seclusion of the sea are entered into in some detail. In the fifth the best form of construction for the main dyke, with its sluices and other works, is minutely discussed, and the whole question of land reclamation by different methods is gone into. The sixth memoir is confined to a delineation of the present condition of the Zuiderzee, its depth, salinity, and the volume and velocity of the rivers which enter it. The seventh memoir has to do with the geological aspects of the scheme, discussing the nature of the sea-bed; and in the last the new plans are compared with former proposals for the same purpose.

As the subject is one of considerable geographical importance, on