

the phenomena have been sufficiently pronounced, the sun is during the day encircled by a more or less distinct *colored halo or corona*. At this place the assumed supra-cirrus volcanic dust seems not to have been sufficiently dense to have developed the colored rings; and there was observed nothing more than a *whitish glare* extending over the sky from 20° to 25° from the centre of the sun. But the Rev. S. E. Bishop writes me from Honolulu, that this chromatic circle around the sun has been constantly observed in all of the Hawaiian Islands for several months. It has likewise been observed in England as a frequent accompaniment of a conspicuous manifestation of the upperglows of sunset and sunrise.

It is an interesting question, whether this more or less distinct colored zone encircling the sun is a true *ice-crystal halo*, or a *diffraction corona*. Its want of sharp definition, and the absence of the regular succession of prismatic tints due to refractive dispersion, would seem to point to diffraction as the true cause of the chromatic phenomena. On the other hand, the large size of the colored circle, having a radius of from 20° to 30° , would seem to connect it with the well-known ice-crystal halo of about 22° radius.

While I am disposed to regard this chromatic feature of the phenomena as mainly due to the diffractive action on light of the impalpable dust-particles suspended in the lofty supra-cirri regions of the atmosphere, yet it is by no means improbable that ice may be associated with the phenomena: for it appears from the experiments of M. Coulier, and more particularly from those of Mr. John Aitken, communicated to the Royal Society of Edinburgh, Dec. 20, 1880 (*Nature*, vol. xxiii, pp. 195-197; also vol. xxiii, p. 384), that the presence of dust-particles in the air is essential to the formation of fogs and clouds; that, when aqueous vapor condenses in the atmosphere, it always does so on some solid nucleus; and that the dust-particles in the air form these nuclei. Now, it is evident that the presence of these attenuated dust-particles in the supra-cirri regions of the atmosphere would produce condensation of the rarefied aqueous vapor at these lofty altitudes. But inasmuch as this region must, even within the tropics, be far above the plane of perpetual congelation, the condensed vapor must necessarily assume the form of aggregations of ice around these nuclei: hence the diffractive coronas may be associated with imperfectly developed ice-crystal halos. JOHN LECONTE.

Berkeley, Cal., Jan. 25, 1884.

Inheritance of physical injuries.

Well-authenticated instances of the inheritance of a physical injury are so rare, that I wish to put upon record one which has recently fallen under my observation. A gentleman, when a boy about seven years of age, had the second toe of the right foot deformed by wearing a tight boot. The first and third toes were crowded together, forcing the second one under and backwards, and causing a curvature of the second joint, which, in time, became permanent. The joint, being somewhat elevated above those of the other toes, received the pressure of the shoe, and always after was more or less troublesome in consequence. The gentleman was twice married. By his first wife he had six children, the second of which was a daughter; the rest, sons. The daughter inherited the crooked toe; but the feet of all the sons were normal. The deformity appeared, however, in the son of one of these, — the brother next younger than the sister, — affecting the same foot and toe as on the grandfather. By his second wife the gentleman had

only one child, a son, who also inherited the peculiarity; but in this instance it was the second toe of the *left* foot, instead of the right, that was affected.

Knowing that much doubt still exists whether the results of a slight physical injury, like the one I have described, are ever transmitted, I have taken pains to examine carefully all the evidence under my observation; and I feel assured of its correctness. All four having the deformed toes are now living, and all agree upon the facts. The gentleman is positive that his feet were normal until he was about seven years old, and says he remembers very distinctly wearing the boots which caused the deformity. An examination of the foot does not show any congenital peculiarity which might have been transmitted. The toe, when restored to its correct position, appeared normal in every way. No peculiarity of this kind has ever appeared in any other of the gentleman's relatives. I can see no way, then, of avoiding the conclusion that the injury, or rather its results, have been transmitted to two generations.

The case presents some features which render it especially interesting. The peculiarity's appearance in the children of both wives seems to eliminate altogether the element of the mother's influence in producing it. The recurrence of the variation in the grandchild, the father being normal, indicates how powerful was the tendency to perpetuate this slight deviation from nature's standard. In the other cases which I have studied personally, if a variation did not appear in a child, that child's children were free from it also. I should be glad to know if any one of your readers has observed this tendency toward reverting to the ancestral type under similar circumstances. IRVING P. BISHOP.

Perry, N.Y., Jan. 28, 1884.

Pumice from Krakatoa.

Capt. A. W. Newell, of the bark Amy Turner of Boston, has brought in some pumice which was washed aboard his vessel, Sept. 17, 1883, in latitude $7^\circ 25'$ south, longitude $103^\circ 21'$ east, about a hundred and sixty-five miles south-west from Krakatoa, Sunda Straits. It covered the sea in windrows, and was observed as fine ashes as far distant as thirteen hundred and fifty miles from its source.

A piece about seven inches by five, which came to my notice, is of a reddish-gray color, and very much inflated: it carries porphyritic crystals of plagioclase felspar, in many cases surrounded by dark-brown glass, forming small black spots in the gray mass, which might at first sight be mistaken for augite or hypersthene. There is, besides, dark-green augite and brown hypersthene, which is strongly pleochroic, and resembles closely that found in the lavas from the volcanoes of northern California and the Cascade Range (Notes on the volcanoes of northern California, Oregon, and Washington Territory, *Amer. Journ. Sci.*, September, 1883).

The percentage of silica for this pumice was found to be 62.53, and is almost identical with that of the hypersthene-bearing pumice from Mount Shasta, which is 62. It is undoubtedly the pumice of a hypersthene andesite, and is especially interesting because of its similarity to rocks found on the western coast of North America. The observations of Rénard on the ashes that fell in Batavia soon after the eruption of Krakatoa (*Nature*, Dec. 6, 1883) show the same component minerals, and have doubtless been made on similar material.

Jos. P. IDDIGS.

U. S. geological survey, New York,
Jan. 30, 1884.