

This article was downloaded by: [North Carolina State University]
On: 13 March 2015, At: 06:55
Publisher: Taylor & Francis
Informa Ltd Registered in England and Wales Registered Number:
1072954 Registered office: Mortimer House, 37-41 Mortimer Street,
London W1T 3JH, UK



Transactions of the Botanical Society of Edinburgh

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/tped18>

IV. Notice of Plants collected in the Neighbourhood of Elie, Fife

Mr J. W. Brown

Published online: 01 Dec 2010.

To cite this article: Mr J. W. Brown (1863) IV. Notice of Plants collected in the Neighbourhood of Elie, Fife , Transactions of the Botanical Society of Edinburgh, 7:1-4, 430-430, DOI: [10.1080/03746606309467879](https://doi.org/10.1080/03746606309467879)

To link to this article: <http://dx.doi.org/10.1080/03746606309467879>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

after which one, or usually two leaves, are developed, and then a little rosette of undeveloped leaves, beset with scales, forming a pseudo-hibernaculum. The young plant up to this period is generally attached to the generative leaf, so that its nourishment is entirely derived from the parent. The vital activity of the recipient plant now in a great measure ceases, the generative leaf undergoes decay, and leaves it a free and independent organism, which, ere spring returns, may have been transported to some distant nidus by the floods, which generally sweep their habitats during winter. As illustrating susceptibility to mechanical irritation, he gave the following experiments on *Drosera rotundifolia* (round-leaved sundew), performed in a temperature of 65° Fahr. :—"Selecting a vigorous plant, I carried on a gentle irritation of the hairs for a short time; their collapsing soon became evident, and in half an hour they were all curved in upon the surface of the laminæ. Again, placing an insect upon the surface of another, the hairs had begun to collapse in twelve minutes, and in twenty minutes nearly all those near the base of the leaf had their glands applied to the insect, while those on the apical part had undergone little or no change. This had been occasioned by the position of the insect, which had been accidentally placed on a line with the petiole near the base of the lamina. Thus it would appear that the *Droseras* do not possess the communicative powers of either their ally the *Dionæa*, the lobes of which collapse on the touching of a single hair, or of the *Mimosa*, which exhibits the same rapid communicability to the other pinna of the leaf when the equilibrium of one is disturbed." The author also detailed some experiments in support of Dr Nitsche's statement, "that their susceptibility to irritation is invariably proportionate to the activity of their secretions, and dependent on the process of assimilation." None of the author's experiments with chemical stimuli elicited any susceptibility to irritation, though he seemed to think that chloroform had an anæsthetic influence. In one of his experiments he placed two portions of the leaf of *Drosera binata* under a bell-glass exposed to the vapour of chloroform. In four minutes no perceptible change having taken place in either, he took one of them out, and found that the hairs were completely anæsthetised. In treating of the cause and functional import of these movements, he was inclined to suppose that in regard to the former no merely physical hypothesis was sufficient to account for the phenomenon, but that it was due entirely to the vital force; and in regard to the latter, he thought Mr Knight's view, that decomposing animal matter might be necessary for certain of the functional requirements of the plant, very plausible, on considering how peculiarly adapted the leaves of *Drosera* and *Dionæa* were for the purpose of catching insects. —

IV. *Notice of Plants collected in the Neighbourhood of Elie, Fife.* By
Mr J. W. BROWN.

Mr Brown gave a list of the rarer plants met with in the district, and noticed several which had not before been observed in that quarter. The list included the following: *Diplostaxis tenuifolia*, *Koniga maritima*, *Senebiera Coronopus*, *Sisymbrium Sophia*, *Lavatera arborea*, *Sagina maritima*, *Geranium pyrenaicum*, *G. pusillum*, *Genista tinctoria*, *Lysimachia vulgaris*, *Mertensia maritima*, *Lycium barbarum*, *Satureia Limonium*, *Beta maritima*, *Rumex aquaticus*. He accounted for the appearance of *Statice Limonium*, *Senebiera Coronopus*, *Lycium barbarum*, &c., by their introduction with ballast.