

Far Eastern Entomologist

Дальневосточный энтомолог

Journal published by Far East Branch
of the Russian Entomological Society
and Laboratory of Entomology,
Institute of Biology and Soil Science,
Vladivostok

Number 307: 1-5

ISSN 1026-051X

March 2016

http://urn.nsd.ru/urn.nsf/obj/19CAA934-11E8-4557-9FB0-803D270E0747

A NEW SPECIES OF *CHYSOTUS* MEIGEN, 1824 (DIPTERA, DOLICHOPODIDAE) FROM CAUCASUS

O. P. Negrobov¹⁾, S. Naglis²⁾, O. O. Maslova³⁾

1) Voronezh State University, Universitetskaya pl. 1, Voronezh, 394006, Russia. *
Corresponding author, E-mail: negrobov@list.ru

2) Institute of Evolutionary Biology and Environmental Studies, University of Zurich,

Winterthurerstrasse 190, Zurich CH-8057, Switzerland. E-mail: s.naglis@bluewin.ch

3) Voronezh State Pedagogical University, Lenina 86, Voronezh 394043, Russia.

Chrysotus chandleri Negrobov, Naglis et Maslova **sp. n.** is described from North Caucasus. The new species is similar to *Ch. blepharosceles* Kowarz, 1874, but differs by the color of the halters, by the lacking of long hairs on the fore tibia, and by the shape of the aedeagus.

KEY WORDS: Diptera, Dolichopodidae, *Chrysotus*, new species, Caucasus, Russia.

О. П. Негров*, С. Наглис, О. О. Маслова. Новый вид *Chrysotus* Meigen, 1824 (Diptera, Dolichopodidae) с Кавказа // Дальневосточный энтомолог. 2016. N 307. C. 1-5.

С Северного Кавказа описан *Chrysotus chandleri* Negrobov, Naglis et Maslova **sp. n.** Новый вид сходен с *Ch. blepharosceles* Kowarz, 1874, но отличается от последнего окраской жужжальцев, отсутствием длинных щетинок на передних голенях и формой эдеагуса.

*Корреспондирующий автор, Воронежский государственный университет, Университетская пл. 1, Воронеж 394006, Россия. E-mail: negrobov@list.ru

INTRODUCTION

The genus *Chrysotus* Meigen, 1824 comprises more than 335 species. The species of this genus are similar in external morphology and have a relatively small number of diagnostic characters, but most species are well distinguished by the structure of the male genitalia, particularly the structure of the aedeagus.

The Palaearctic species of *Chrysotus* were revised by Negrobov and his co-workers in a series of publications (Negrobov, 1980; Negrobov & Maslova, 1995; Negrobov *et al.*, 2000, 2003). Later *Ch. bifurcatus* Wang et Yang, 2009, *Ch. brevicercus* Wang et Yang, 2009, *Ch. luoyangensis* Wang et Yang, 2009, *Ch. sinuolatus* Wang et Yang, 2009, *Ch. xinjiangensis* Wang et Yang, 2009, *Ch. beijingensis* Wang et Yang, 2006, *Ch. serratus* Wang et Yang, 2006, *Ch. thornpenis* Liu, Wang et Yang, 2015, *Ch. shanxiensis* Liu, Wang et Yang, 2015, and *Ch. shanxiensis* Liu, Wang et Yang, 2015 were described from China (Wang & Yang, 2006, 2008, 2009; Wei & Yang, 2007; Liu *et al.*, 2015). Five species were described from Russia, namely *Ch. chukotkensis* Grichanov, 2012 from Chukotka, *Ch. brooksi* Negrobov, Barkalov et Selivanova, 2013 from Primorskii krai, *Ch. cilitibia* Maslova et Negrobov, 2015 from Sakhalin and Kuril Islands, *Ch. komovi* Negrobov, Barkalov et Selivanova, 2014 and *Ch. tricaudatus* Negrobov, Barkalov et Selivanova, 2014 from Taimyr (Grichanov, 2012; Negrobov *et al.*, 2013, 2014; Maslova & Negrobov, 2015). Naglis (2010) described *Ch. dischmaensis* Naglis, 2010 from Switzerland. *Ch. tagoi* Negrobov, Maslova et Fursov, 2015 and *Ch. saigusai* Negrobov, Fursov, Kumazawa et Tago, 2015 are described from Japan (Negrobov *et al.*, 2015). Thus, 82 species of *Chrysotus* are recorded from the Palaearctic Region. Negrobov *et al.* (2000) published the last key to males of the Palaearctic species. A key to males of the Chinese species was published by Yang *et al.* (2011).

Up to now 13 species of *Chrysotus* are known from Caucasus: *Ch. collini* Parent, 1923 from North Caucasus (Grichanov *et al.*, 2006), *Ch. laesus* (Wiedemann, 1817), *Ch. angulicornis* Kowarz, 1874, *Ch. pennatus* Lichtwardt, 1902 from Krasnodar Territory (Negrobov, 1965; Maslova *et al.*, 2007), *Ch. cilipes* Meigen, 1824, *Ch. cupreus* Macquart, 1824, *Ch. defensus* Negrobov et Maslova, 2000, *Ch. femoratus* Zetterstedt, 1843, *Ch. gramineus* (Fallen, 1823), *Ch. neglectus* (Wiedemann, 1817), *Ch. peculiariter* Negrobov et Maslova, 2000, *Ch. pulchellus* Kowarz, 1874, and *Ch. suavis* Loew, 1857 from the Caucasian State Biosphere Reserve (Negrobov, 1967; Negrobov *et al.*, 2000; Grichanov, 2007). One new species from Caucasus is described below. Holotype and paratype of the new species are deposited in the Zoological Institute of the Russian Academy of Sciences, St. Petersburg (ZIN).

DESCRIPTION OF THE NEW SPECIES


Chrysotus chandleri Negrobov, Naglis et Maslova, sp. n.

Figs 1–4

MATERIAL. Holotype – ♂, Russia: “Mountain Dzhuga, Kuban oblast, Moykopskoe otdelenie” (=Krasnodar region: Tuapse district, Mt. Dzhubga), 17.VI 1911, coll. Volnuhin (ZIN). Paratype: 1 ♂, same data as holotype (ZIN).

DESCRIPTION. Male. Frons metallic green, with gray pollen. Face dark, with brown pollen, its width is less the distance between ocellar setae. Antennae black, postpedicel large, transversely oval. Palps brown. Legs black with black setae. Mid tibia with 2 long anterodorsal bristles. Hind tibia with long setae on the outer side. Wings no darkened. Halters dark. Fore tibia without long hairs.

Head. Frons metallic green, with gray pollen. Face dark, with brown pollen, its width is less the distance between ocellar setae, its width in the middle is about twice less than the length of the postpedicel (0.2: 0.4). Antennae black, postpedicel large, transversely oval, with short hairs. Width of postpedicel longer than its length (0.9: 0.4). Stylus subapical, with short hairs, much longer than postpedicel (0.4: 2.6). Palps brown, with black hairs. Lower postoculars white.



Figs. 1–7. *Chrysotus* spp. 1–4 – *C. chandleri*, sp. n.: 1 – aedeagus, lateral view, 2 – antenna, 3 – hypopygium, lateral view, 4 – aedeagus, dorsal view; 5–7 – *C. blepharoscelis* Kowarz; 5 – hypopygium, lateral view, 6 – apical part of aedeagus, lateral view, 7 – aedeagus, dorsal view.

Thorax. Shining green, weakly grey pollinose; pleura grey pollinose. Mesonotum shiny green with violet shade. 6 pairs of strong dorsocentral bristles. Proepisternum with black setae. Acrostichals biseriate, well developed. Scutellum with a pair of strong medial scutellars. Legs black with black setae. Fore coxae with black setae. Fore femur with short anteroventral hairs. Fore tibia with 2 strong anterodorsal setae, its length is shorter than the diameter of the tibia. Fore tarsus without short hairs. Mid femur with few short anteroventral bristles. Mid tibia with 2 long anterodorsal bristles. Mid tarsus without short hairs. Hind femur with long anteroventral bristles, its length is more than half the width of the femur. Hind tibia with 2 long anterodorsal and 2 posterodorsal bristles, and with long setae on the outer side which are not longer than the diameter of the tibia, and with a number of short black bristles on the ventral side. Hind tarsus with short hairs. Podomeres (from tibia to fifth tarsomere) length ratio: fore leg: 3.3: 1.7: 0.8: 0.6: 0.4: 0.3, mid leg 5.0: 2.0: 0.9: 0.6: 0.5: 0.4, hind leg 4.5: 1.4: 1.0: 0.7: 0.5 : 0.4.

Wings. Membrane no darkened; veins dark; R_{4+5} and M_{1+2} parallel at the apex; costal section between R_{2+3} and R_{4+5} about twice as long as the costal section between R_{4+5} and M_{1+2} (1.9: 0.9); distal section of M_{1+2} longer than basal section (5.7: 3.5); ratio of cross-vein dm-cu to distal part of CuA_1 0.9: 2.8; lower calypter with black cilia; halter black.

Abdomen. Metallic green, grey pollinose, covered with black hairs; epandrium nearly circular; surstyli narrow, curved ventrally at apex, with setae in apical part. Aedeagus without appendages. Cerci oval with fine black bristles.

Female unknown.

MEASUREMENTS. Body length 1.8–2.2 mm, wing length 1.9 mm.

ETYMOLOGY. The new species is named after the English dipterologist Peter Chandler.

COMPARISON. According to the key to the Palaearctic species of the genus *Chrysotus* (Negrobov *et al.*, 2000), the new species is keyed to *Ch. blepharosceles*, but differs by the following characters:

- 1(2) Halters dark. Fore tibia without long hairs. Hypopygium – Fig. 3. Aedeagus at the apex without lateral processes (Figs 1, 4) *Chrysotus chandleri* sp. n.
- 2(1) Halters yellow. Fore tibia with long dorsal and ventral hairs, longest hairs longer than the diameter of the tibia. Hypopygium – Fig. 5. Aedeagus at the apex with 2 lateral processes (Figs 6, 7) *Ch. blepharosceles* Kowarz, 1874

ACKNOWLEDGEMENTS

This work was supported by the grant of the Russian Foundation for Basic Research No 14-04-00264.

REFERENCES

- Grichanov, I.Ya., Kustov, S.Yu. & Volfov, B.I. 2006. A checklist of Dolichopodidae (Diptera) of Krasnodar Territory and Adygea (Russia). *International Journal of Dipterological Research*, 17(1): 35–55.

- Grichanov, I.Ya. 2007. New records of Dolichopodidae (Diptera) from the Middle East. *International Journal of Dipterological Research*, 18(3): 141–153.
- Grichanov, I.Ya. 2012. A new peculiar species of *Chrysotus* from the Far East of Russia (Dolichopodidae, Diptera). *Amurian Zoological Journal*, 4(4): 333–335.
- Liu, R.S., Wang, M.Q. & Yang, D. 2015. *Chrysotus* Meigen (Diptera: Dolichopodidae) from Shanxi, China, with descriptions of two new species. *Zoological Systematics*, 40(1): 86–92.
- Maslova, O.O. & Negrobov, O.P. 2015. A new species of *Chrysotus* Meigen, 1824 (Dolichopodidae: Diptera) from Sakhalin and the Kuril Islands. *Caucasian Entomological Bulletin*, 11(1): 201–203. [In Russian].
- Naglis, S. 2010. A new species of *Chrysotus* (Diptera, Dolichopodidae) from Switzerland. *Mittheilungen Schweizer entomologischen Gesellschaft*, 83: 199–205.
- Negrobov, O.P. 1965. New and little-known species of Dolichopodidae (Diptera) of USSR fauna. *Entomologicheskoe Obozrenie*, 44(2): 438–446. [In Russian].
- Negrobov, O.P. 1967. Fauna and vertical distribution of Dolichopodidae (Diptera) of Western Caucasus. *Zoologicheskii Zhurnal*, 46(7): 1074–1079. [In Russian].
- Negrobov, O.P. 1980. A revision of Palaearctic species of the genus *Chrysotus* Mg. (Diptera, Dolichopodidae). I. *Ch. ciliipes* Mg. and *Ch. laesus* Wied. species groups. *Entomologicheskoe Obozrenie*, 59(2): 415–420. [In Russian].
- Negrobov, O.P., Barkalov, A.V. & Selivanova, O.V. 2014. Two new *Chrysotus* Meigen (Diptera, Dolichopodidae) from Siberia, with a key to the Siberian species. *Zootaxa*, 3815(3): 409–416. DOI:10.11646/zootaxa.3964.3.8
- Negrobov, O.P. & Maslova, O.O. 1995. Revision of Palaearctic species of the genus *Chrysotus* Mg. (Diptera, Dolichopodidae), II. *Entomologicheskoe Obozrenie*, 74(2): 456–466. [In Russian].
- Negrobov, O.P., Maslova O.O. & Fursov, V.N. 2015. New data on the genus *Chrysotus* Meigen, 1824 (Diptera: Dolichopodidae) from Japan. *Far East Entomologist*. 293: 10–15.
- Negrobov, O.P., Selivanova, O.V. & Grichanov, I.Ya. 2013. Redescription of male and first record of *Chrysotus pilitibia* Negrobov et Maslova from Mongolia (Dolichopodidae, Diptera). *Cesa News*, 89: 3–6.
- Negrobov O.P., Selivanova, O.V. & Maslova, O.O. 2013. A new species of *Chrysotus* Meigen (Diptera, Dolichopodidae) from Primorye. *Dipterists Digest*, 20: 187–190.
- Negrobov, O.P., Tsurikov, M.N. & Maslova, O.O. 2000. Revision of the Palaearctic species of the genus *Chrysotus* Mg. (Diptera, Dolichopodidae), III. *Entomologicheskoe Obozrenie*.79(1): 227–238. [In Russian].
- Negrobov, O.P., Tsurikov, M.N. & Maslova, O.O. 2003. A revision of the Palaearctic species of the genus *Chrysotus* Mg. (Diptera, Dolichopodidae), IV. *Entomologicheskoe Obozrenie*, 82(1): 223–228. [In Russian].
- Wang, M. & Yang, D. 2006. Species of *Chrysotus* Meigen from Beijing (Diptera, Dolichopodidae). *Deutsche Entomologische Zeitschrift*, 53(2): 249–255.
- Wang, M.Q. & Yang, D. 2008. New Species of *Chrysotus* from China (Diptera: Dolichopodidae). In: Shen, X.C. et al. (Eds.). *Classification and Distribution of Insects in China*. China Agricultural Science & Technology Press, Beijing. P. 23–32. [In Chinese].
- Wang, M. & Yang, D. 2009. Species of *Chrysotus* Meigen in Palaearctic China (Diptera: Dolichopodidae). *Entomologica Fennica*, 19: 232–240.
- Wei, L.M. & Yang, Z.H. 2007. Dolichopodidae. In: Li, Z.Z, Yang, M.F. & Jin, D.C. (Eds.). *Insects from Leigongshan Landscape*. Guizhou Science & Technology Press, Guiyang. P. 561–587. [In Chinese].
- Yang, D., Zhang, L., Wang, M. & Zhu, Y. 2011. Dolichopodidae. *Fauna Sinica, Insecta*, Vol. 53. Science Press, Beijing. 1912 pp. [In Chinese, with English summary].