

**17th International *Symposium on*
Aquatic Plants - Back to the Future of
Aquatic Plants and the Way Forward**
Lisbon 2025

Organising Committee

Francisca C. Aguiar (Chair) & Sofia F. Santos

Local Organisation Mid-Symposium Excursions

Ana D. Caperta, Portugal

Ana Mendes, Portugal

Patrícia Rodríguez-González, Portugal

Carla Pinto Cruz, Portugal

Scientific Committee

Teresa Ferreira, Portugal (Chair)

Elisabeth M. Gross, France

Francisca C. Aguiar, Portugal

Daniel Larkin, USA

Deborah Hofstra, New Zealand

Eva Papastergiadou, Greece

Iris Stiers, Belgium

Jan Vermaat, Norway

Jonas Schoelynck, Belgium

Julie Coetzee, South Africa

Kevin Wood, UK

Krzysztof Szoszkiewicz, Poland

Maria Dolores Bejarano, Spain

Rossano Bolpagni, Italy

Sabine Hilt, Germany

Sándor Szabó, Hungary

Sidinei Magela Thomaz, Brazil

Susanne Claudia Schneider, Norway

Takashi Asaeda, Japan

Tenna Riis, Denmark

Wei Li, China

Citation suggestions

This book: Aguiar, Francisca C. & Santos, Sofia F. (eds.). 2025. *Book of Abstracts 17th International Symposium on Aquatic Plants: Back to the Future of Aquatic Plants and the Way Forward*, 15-19 September 2025, Lisbon, 199 pp.

An Abstract from this book

Neiva, J., Frade, D., Serrão, E. 2025. Endangered aquatic halophytes in a cultural landscape: importance and habitat partition in man-made habitats. In: Aguiar, Francisca C. & Santos, Sofia F. (eds.). *Book of Abstracts 17th International Symposium on Aquatic Plants: Back to the Future of Aquatic Plants and the Way Forward*, 15-19 September 2025, Lisbon. p. 146.

Cover

Ranunculus peltatus Schrank (pond water-crowfoot), a freshwater hydrophyte species observed in lentic waters of Portugal, South-Western Europe; Francisca C. Aguiar.

Sponsors

MUSEU
DA ÁGUA

EPAL
Grupo Águas de Portugal



Logistics



Layout



Aleksandra Marković, Susanne C. Schneider, Danijela Vidaković, Miloš Ćirić, Vladimir Petrović, Željka Milovanović, Andreas Ballot

University of Belgrade, Institute of Chemistry, Technology and Metallurgy, National Institute of the Republic of Serbia, Njegoševa 12, 11000 Belgrade, Serbia

Norwegian Institute for Water Research – NIVA, Økernveien 94, 0579 Oslo, Norway

Developing eDNA-based methods for exploring charophyte algae diversity in lakes - Oral presentation

Over the last two decades, eDNA metabarcoding has rapidly evolved into a cost-effective and time-efficient methodology, widely used in areas such as biodiversity assessment, biomonitoring, invasive species control, and the discovery of rare, cryptic and unknown species. This has boosted a great number of new studies and the development of new methods. Still, eDNA studies on aquatic plants lag behind those on animals, as it is much more difficult to find suitable eDNA markers, but a considerable number have recently emerged. Yet, an important group of aquatic plants, the charophytes, remains neglected. Charophyte algae play a vital role in aquatic ecosystems and contribute to a better ecological status of the ecosystems. Due to their sensitivity to water quality, they have long been used for ecological status assessment of lakes. However, their diversity is often underestimated as they have great morphological plasticity and sometimes short life cycles. Within the BIOLAWEB project, we investigated charophyte algae diversity in lakes using both eDNA metabarcoding and a traditional morphological approach. Samples were collected in two saline and two freshwater lakes in Serbia, in three seasons. *rbcL* gene marker was used for metabarcoding and Oxford Nanopore technology for sequencing. Bioinformatic analyses were conducted using the cloud-based metagenomic platform CZID, using NCBI as a database. The aligned sequences were checked manually and the consensus sequences were

then aligned with confirmed sequences from the Boldv4 database. The morphological approach revealed the presence of charophytes in three out of four lakes. Two lakes had only one charophyte species, both belonging to the genus *Chara*, while the third lake had six species belonging to three genera, *Chara*, *Nitella* and *Nitellopsis*. The eDNA metabarcoding approach detected the presence of charophytes in all four lakes, all belonging to the genus *Chara*. However, due to the high similarity of the selected *rbcL* barcode, the *Chara* sequences could only be assigned to two broad charophyte species-complexes (*Chara hispida* and *Chara vulgaris* complex). This study is the first time eDNA metabarcoding has been successfully used to detect charophytes in lakes. To provide species-level identification and identify representatives of all genera, further fine-tuning of the method is required, particularly in generating primers that better discriminate charophyte diversity.

Acknowledgements: This project has received funding from the European Union's Horizon Europe research and innovation programme under grant agreement No 101079234 and from the Ministry of Science, Technological Development and Innovation of the Republic of Serbia (451-03-66/2024-03/200026).

