

This article was downloaded by: [Purdue University]  
On: 17 January 2015, At: 08:36  
Publisher: Taylor & Francis  
Informa Ltd Registered in England and Wales  
Registered Number: 1072954 Registered office:  
Mortimer House, 37-41 Mortimer Street, London W1T  
3JH, UK



## Annals and Magazine of Natural History: Series 1

Publication details,  
including instructions for  
authors and subscription  
information:

[http://  
www.tandfonline.com/loi/  
tnah07](http://www.tandfonline.com/loi/tnah07)

### XXVI.—Observations on the family Helicidæ, and description of a new genus

Dr. L. Pfeiffer<sup>a</sup>

<sup>a</sup> Cassel

Published online: 15 Mar  
2010.

To cite this article: Dr. L. Pfeiffer (1840) XXVI.—Observations on the family Helicidæ, and description of a new genus , Annals and Magazine of Natural History: Series 1, 5:31, 238-242, DOI: [10.1080/00222934009496817](https://doi.org/10.1080/00222934009496817)

To link to this article: <http://dx.doi.org/10.1080/00222934009496817>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

ally foreign to the ovule, and introduced into it from without.

#### EXPLANATION OF PLATE IV.

*Fig. 1.* Part of the ovule and conducting tissue of *Phytolacca decandra*. *a.* Conducting tissue. *b.* Pollen tube. *c.* Embryo. *d.* Embryo-sac. *e.* Nucleus. *f.* Secundine. *g.* Primine. (Schleiden.)

*Fig. 2.* The extremity of the pollen tube (embryo) indenting the embryo-sac. *a.* Pollen tube. *b.* Embryo. *c.* Embryo-sac. (Schleiden.)

*Fig. 3.* The inferior part of an ovule of *Carduus nutans*, after impregnation. *a.* Pollen tube. *b.* Embryo. *c.* Embryo-sac. *d.* Nucleus. *e.* Teguments. (Schleiden.)

*Fig. 4.* Section of the ovarium of *Zea Mays* at an early period of its development. *a.* Primine. *b.* Secundine. *c.* Nucleus. *d.* The little cavity in which the primary utricle is afterwards formed. (Mirbel and Spach.)

*Fig. 5.* The same at a more advanced period. *a.* The primary utricle. (Mirbel and Spach.)

*Fig. 6.* The primary utricle, detached from the ovule, filled with the globulo-cellular cambium. (Mirbel and Spach.)

*Fig. 7.* The embryo detached. *a.* Cotyledon. *b.* The first leaf of the plumule. *c.* The second leaf of the plumule. (Mirbel and Spach.)

*Fig. 8.* The embryo at a more advanced period. *a.* The first leaf of the plumule. *b.* Radicle. *c.* The suspensor. (Mirbel and Spach.)

#### XXVI.—*Observations on the Family Helicidæ, and description of a new Genus.* By Dr. L. PFEIFFER of Cassel\*.

THE most difficult question concerning the limits of genera among the land mollusca has of late been frequently treated of, and with widely different results. If, on the one hand, Férussac went much too far, in comprising nearly all air-breathing mollusca with four tentacula in his genus *Helix*, still, on the other hand, the attempts at a division of this large group have not yet succeeded in a satisfactory manner. Draparnaud's genera, however, form a good basis, to which I am inclined with slight deviation to return. Lamarck evidently relied too much on individual peculiarities of the shell, because he was not acquainted with a sufficient number of species in which the transitions of the forms may be distinctly

\* From Wiegmann's Archiv, Part I. 1840.

demonstrated. This is especially the case with the genus *Achatina*, which had necessarily, in as far as its character was merely founded on the truncated columella, to be again united with *Bulimus*, as the animals, both as to their exterior as well as their anatomical characters, are perfectly similar, and all the intermediate forms may be followed, from the truncate columella of *Achat. virginea* to the beautiful round aperture of *Bul. hæmastomus*. But those species are excluded from this reunion which Montfort comprised in his genus *Polyphemus*, as these are not merely distinguished by the peculiar form of the columella, but likewise by the bilobate snout of the animal.

I follow therefore in general the well sustained view of Deshayes, who reunites the genera *Achatina* and *Bulimus* of Lamarck, and shows on anatomical grounds (Lam. viii. p. 14. 2de édit.) that they cannot be comprised under *Helix*; but I cannot at all agree in the opinion expressed by the same ingenious naturalist, that *Clausilia* and *Pupa* must likewise be combined. I find, rather, no reason for separating Draparnaud's genus *Pupa* from *Bulimus*. In the European species described by Draparnaud, there were, it is true, some constant characteristics evident which appeared to justify this separation; but with how many species have we since then become acquainted, which have shown the earlier generic characters to be insufficient, and have therefore been sometimes ascribed to the one, sometimes to the other of these genera! In fact, I know at present not a single distinctive character between the two. The animals are perfectly identical; the form is in both cylindrical or ovate; and the oral edge disconnected. What therefore is left? The form of the aperture, or its folds and teeth? The form of the columella? For all these characteristics the most distinct forms of transition are afforded by Férussac's genus *Partula* or Swainson's *Achatinella*. Are the teeth and folds of the aperture to constitute the chief character,—which for instance Menke seems to adopt, since he refers to the *Pupæ*, the long-known *Bulimus Pupa*? But how many toothed species do we not now refer without scruple to *Bulimus*, following the analogy with *Helix*, while at the same time we do not class, for instance, *Pupa obtusa* among them! In the ex-

ceedingly difficult group of the large extra-European forms; the teeth of the aperture and the folds of the columella are a very uncertain character; sometimes they are present, sometimes missing: *Pupa sulcata* is perfectly toothless. On the other hand, the great affinity of this group with *Bulimus labiosus*\*, Br. is evident, and this can just as little be separated from *Bulimus faba*, Desh. (*Partula australis*, Fér.) as the latter from *Bul. agotis*, Mke. (*Auricula Sileni*, Lam.) *Bulimus citrinus* and the other true species of *Bulimus* (according to Draparnaud!). Indeed the most certain proof of a genus having been falsely established is when several of its species are thrown from one to another; or generally when we are uncertain to which any species that occurs is to be carried.

From all these reasons I consider that the genus *Pupa*, Dr., must be wholly discarded, and its species arranged with *Bulimus*. But the genus *Clausilia*, according to its old Draparnaudian characteristic, remains firm. Did there exist no other distinctive character, the clausium—a part evidently analogous to the operculum of several molluscos genera—would alone suffice to establish the genus; and if we define this with Draparnaud's short words thus: "*Testa fusiformis; peristoma continuum oblongum; clausilium!*" we have a well-defined whole. It is true, a portion of the species included by Draparnaud in this genus, to suit which Lamarck (ed. Desh. viii. p. 195.) says, Ce nom fut d'abord significatif! must then be excluded; viz. 1<sup>o</sup>, all those which have no continuous peristoma; and 2<sup>o</sup>, those which have a circular continuous peristoma but possess no clausium. Of the former I will merely mention the *Clausilia exesa*, Spix (Desh. No. 39.), and Turton's *Balea fragilis*, which indeed is reckoned by Draparnaud, Nilsson, and Lamarck as a *Pupa*; but by Studer, C. Pfeiffer (vol. iii.), and Menke as a *Clausilia*. Both must range, together with the genus *Pupa*, under *Bulimus*. To the second section belong Lamarck's

\* Desh. No. 130. This beautiful species, adopted by Deshayes only from Müller's excellent description, is in my possession. It is represented with perfect truth in Chemnitz (ix. p. 1234.), but the figure in Gualtieri (T. 4. R.) cited with a query has no relation to it, but belongs to the species which Blainville (Malacol. tab. 39. fig. 5 a.) has figured as *Pupa Mumia*.

and Deshayes' species, Nos. 2, 9, 40, 41, 42, 43. But since all these, on account of the regularly continuous peristoma, can be classed with none of the other genera of the family *Helicidae*, and on account of the structure of the animal can far less be referred anywhere else, I conceive that they form of themselves a good genus, and propose for this the name *Cylindrella*,—a name which in the first place points to the form of all the species known, modified by the termination already in use in molluscou genera.

The characteristic of this new genus would be as follows :—

*CYLINDRELLA*, *L. Pfr.*—Animal heliciforme. Testa subcylindracea, imperforata, multispirata, sæpe truncata. Peristoma continuum, suborbiculare. Operculum vel clausium nullum.

All hitherto known species are inhabitants of the West Indian Islands, and I myself have found in Cuba four evidently distinct species, of which two have already been described and figured by Férussac (*Helix Cochlodina perplicata* and *subula*), the two others are perhaps new. The latter have been preliminarily described by me in Wiegmann's Archiv (1839, p.353.) under the name *Clausilia elegans* and *crispula*. It appears remarkable to me that all the Cuban *Cyclostomata* with which I am acquainted are always truncate, i. e. cast off the apex at a certain age, and reclose the open place. Almost all known species are dextral, and we should be justified in adopting this as a generic character if Chemnitz's *Turbo elongatus* from Jamaica (*Clausilia Chemnitziana*, Desh.) was not sinistrously whorled, according to the figure and clear description. (Chemn. ix. fig. 956.) In other respects this species is so nearly allied to my *Cyl. elegans* in its habit, that we may admit with certainty that it likewise has no clausium, and belongs to the new genus, of which the following are the hitherto known species :—

1. *Cylindrella gracilicollis* (*Clausilia truncatula*, Lam. 2.)
2. ——— *collaris* (Claus. *collaris*, Lam. 9.)
3. ——— *antiperversa* (Claus. *antiperversa*, Desh. 40.)
4. ——— *subula* (Claus. *subula*, Desh. 41.)
5. ——— *perplicata* (Claus. *perplicata*, Desh. 42.)
6. ——— *Chemnitziana* (Claus. *Chemnitziana*, Desh. 43.)

7. *Cylindrella elegans* } Clausilia, L. Pfeiffer in Wiegmann's  
 8. ————— *crispula* } Archiv, Part I. 1839, p. 353.  
 9. —————? *torticollis* (Claus. *torticollis*, Lam. 1.)

The first five of these species are figured by Férussac on the 163rd plate, and are referred by him to the subgenus *Cochlodina*. Rossmässler calls them, ('Iconographie', Part II. p. 13.) "long-necked *Pupæ*," under which Sowerby ('Genera of Shells,') also classes some species belonging here. Whether *Clausilia torticollis*, Lam. from Candia, must likewise be referred to *Clausilia* I will not venture to determine, as I am not acquainted with the species, nor is the figure at present at my disposal, and the description is slight, especially with reference to the aperture. Yet much speaks in favour of this being its true position.

From what has been above stated it appears to me advantageous to divide the family of the *Helicidæ* into the following genera:—

1. *Vitrina*.
2. *Helicophanta*.
3. *Succinea*.
4. *Helix* (with *Carocolla* and *Anostoma*, Lam.)
5. *Bulimus* (with *Achatina*, Lam., *Pupa*, Drap., *Partula*, Fér., and *Megaspira*, Lea.)
6. *Vertigo*.
7. *Cylindrella*.
8. *Clausilium*.
9. *Polyphemus*, Mont.

The character which all have in common with the *Limacidæ*, and by which they are distinguished from the following orders, are the retractile tentacles provided at the apex with eyes; and the separate genera appear to me only in this way capable of being sufficiently established according to correct principles. Perhaps, however, some changes must be made in accordance with the structure of the animals, as I am chiefly led to suppose from some observations made on living specimens of *Bulimus hæmastomus*.