



Annals and Magazine of Natural History: Series 2

Publication details, including instructions
for authors and subscription information:
<http://www.tandfonline.com/loi/tnah08>

XXXII.—Anatomical description of a species of asteroid polypes, probably forming the type of a new genus of Alcyonidæ

John Denis Macdonald

Published online: 26 Oct 2009.

To cite this article: John Denis Macdonald (1857) XXXII.—Anatomical description of a species of asteroid polypes, probably forming the type of a new genus of Alcyonidæ , *Annals and Magazine of Natural History: Series 2*, 19:113, 391-391, DOI: [10.1080/00222935708693952](https://doi.org/10.1080/00222935708693952)

To link to this article: <http://dx.doi.org/10.1080/00222935708693952>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses,

actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

Both *Bowerbankia* and *Fesicularia* agree in the uniserial and unilateral distribution of the polypes, but in the present instance the cells are arranged in linear and bilateral clusters.

XXXII.—*Anatomical Description of a Species of Asteroid Polypes, probably forming the type of a new genus of Alcyonidæ.*

By JOHN DENIS MACDONALD, Assistant Surgeon R.N.*

ON leaving the Conway Reef (lat. $21^{\circ} 44' 48''$ S., long. $174^{\circ} 37' 45''$ E.), July 4, 1855, a very beautiful branched asteroid Zoophyte, belonging to the Alcyonidæ, was brought up from a depth of between 30 and 40 fathoms, on the buoy-rope of the anchor.

The polypidom, from a trunk of about one inch and a half in diameter, branched off, with much irregularity, but generally in a dichotomous manner, into very minute subdivisions.

The investing membrane was strengthened by the close deposition of elongated, fusiform, and minutely tuberculated spicula of a deep crimson-lake tint, which impart their colour to the whole mass.

The internal substance was chiefly composed of longitudinal muscular septa, radiating from the central axis (which contained no denser material), frequently communicating with one another laterally, and being fixed into the internal surface of the integument in vertical lines. These muscular septa were invested on each side with a layer of finely reticulated vessels; both sets being connected by numerous transverse trunks passing through the intervening muscular tissue, and the spaces between the septa were filled with a transparent glairy fluid.

The polyp-cells were exposed and solitary, resting on the internal surface of short branchlets strengthened by large dorsal spicula, one of which, much larger than the rest, extended considerably beyond the polyp-cells, tapering gently to a needle-like point. These latter spicula are covered with tubercles, and in every respect but in size similar to those of the general integument.

The small spicula on the internal or ventral surface of the branchlets diverge from one another in the peripheral direction, while those on the dorsal border are disposed longitudinally.

The mouth of each cup-like polyp-cell was surrounded with about eight projecting spicula, whose fixed extremities were curved upwards and inwards, festoon-fashion, while numerous smaller ones were so disposed as to fill up the open spaces posteriorly, and thus strengthen the body of the cell.

Although I have not been able to count the number of the oral tentacula satisfactorily, from their proportional size I can readily believe that there were about eight in this species, as in most if not all other asteroid polypes. They were broad and flat, tapering to a blunt point, like those of *Sarcoedictyon* (Forbes), to which genus I have no doubt this Zoophyte is nearly allied, though the habit of the polypidom is so very different.

* From the Proceedings of the Royal Society, March 12, 1857.