

Article

The Potential Exploitation of the Mediterranean Invasive Alga *Caulerpa cylindracea*: Can the Invasion Be Transformed into a Gain?

Loredana Stabili ^{1,2,*}, Simonetta Frascchetti ^{1,3}, Maria Immacolata Acquaviva ², Rosa Anna Cavallo ², Sandra Angelica De Pascali ¹, Francesco Paolo Fanizzi ¹, Carmela Gerardi ⁴, Marcella Narracci ² and Lucia Rizzo ^{1,3}

- ¹ Dipartimento di Scienze e Tecnologie Biologiche ed Ambientali, Università del Salento, Via Prov.le Lecce Monteroni, 73100 Lecce, Italy; simona.frascchetti@unisalento.it (S.F.); sandra.depascali@unisalento.it (S.A.D.P.); fp.fanizzi@unisalento.it (F.P.F.); lucia.rizzo@unisalento.it (L.R.)
- ² Istituto per l'Ambiente Marino Costiero, Unità Operativa di Supporto di Taranto, CNR, Via Roma 3, 74123 Taranto, Italy; maria.acquaviva@iamc.cnr.it (M.I.A.); rosanna.cavallo@iamc.cnr.it (R.A.C.); marcella.narracci@iamc.cnr.it (M.N.)
- ³ CoNISMa—Consorzio Nazionale Interuniversitario per le Scienze del Mare, Piazzale Flaminio 9, 00196 Rome, Italy
- ⁴ Istituto di Scienze delle Produzioni Alimentari, Unità Operativa di Supporto di Lecce, via Prov.le Lecce-Monteroni, 73100 Lecce, Italy; carmela.gerardi@ispa.cnr.it
- * Correspondence: loredana.stabili@iamc.cnr.it; Tel.: +39-0832-298971

Academic Editor: Sylvia Urban

Received: 29 July 2016; Accepted: 9 November 2016; Published: 15 November 2016

Abstract: Recently, there is a growing interest towards the development of strategies for invasive seaweed control and exploitation as source of secondary metabolites. Here, we investigated the potential of exploitation in biotechnology and recycling options in eradication programs of the lipidic extract of the Mediterranean invasive seaweed *Caulerpa cylindracea* (Chlorophyta). The chemical characterization was carried out by means of multinuclear and multidimensional NMR spectroscopy. The fatty acid profile of *C. cylindracea* assessed the presence of several types of molecules known for antioxidant activity such as carotenoids, chlorophylls, pheophytins, and sterols. The NMR spectroscopy showed also the characteristic signals of saturated, unsaturated, and free fatty acids as well as other metabolites including the biopolymer polyhydroxybutyrate. The lipidic extract exerted an antioxidant activity corresponding to 552.14 ± 69.13 mmol Trolox equivalent/g (ORAC) and to 70.3 ± 2.67 mmol Trolox equivalent/g (TEAC). The extract showed an antibacterial activity against several *Vibrio* species, suggesting its potential use in the control of diseases in mariculture. Our results show that *C. cylindracea*, representing a critical hazard in coastal areas, could be transformed into a gain supporting specific management actions to reduce the effects of human pressures.

Keywords: seaweed; *C. cylindracea*; NMR spectroscopy; lipidic extract; antioxidant activity; antimicrobial activity; fatty acids

1. Introduction

Marine organisms are rich sources of structurally novel and biologically active metabolites, exhibiting numerous interesting biological effects and thus constituting valuable opportunities for drug discovery. Recently, several studies on the growing problem of non-indigenous species have proven that the knowledge of marine bioactive compounds may indicate the capacity of non-indigenous species to invade new systems [1–9]. In a recent review, Mollo et al. [10] showed that the commercial use of the natural products obtainable from marine bioinvaders might also be considered as an effective option for reducing their impact on marine ecosystems.

In coastal habitats, macroalgae are abundant and represent a source of bioactive metabolites exhibiting numerous promising and remarkable biological activities capable of influencing the abundance, distribution, and survival of marine organisms [11,12]. Seaweeds are known for their richness in polysaccharides, minerals, and certain vitamins [13]. Marine algae have been also recognized as potential source of antioxidants [14,15] and traditionally consumed as a readily available whole food, especially among coastal communities [16–18]. Fatty acids and enzymatic and non-enzymatic antioxidant properties of have been investigated in *Caulerpa* species [19–22]. Macroalgae also contain bioactive substances like polysaccharides, proteins, lipids, and polyphenols, with antibacterial, antiviral, and antifungal properties [23].

Seaweeds have been prescribed for several diseases in different Asian traditional medical systems [18]. In recent years, pathogenic bacteria resistant to multiple drugs have become a worldwide emergency. The discovery of new antibacterial compounds, as suitable substitutes to conventional antibiotics, might be a possible solution to this problem. Seaweeds could represent a potential source of new antimicrobial compounds [24]. Ballesteros and Uriz [25] have screened several marine macrophytes from the Central Mediterranean to evaluate the production of antibacterial, antifungal, compounds, founding, among the Chlorophyta, with the maximum level of activity found in the order Bryopsidales. In particular, species belonging to the genus *Caulerpa* show interesting antibacterial activity compared to other groups of algae [26]. A study conducted on the aqueous extract of *Caulerpa racemosa* collected in India (Gulf of Mannar) showed a pronounced antibacterial activity against different pathogenic organisms [27]. Moreover, the methanol extract of *C. racemosa* was found to exhibit significant activity against the test pathogens *Staphylococcus aureus*, *Escherichia coli*, and *Enterobacter aerogenes* [28].

In the present study, we focused on the Mediterranean non indigenous seaweed *Caulerpa cilindracea* (Sonder) [29], previously known as *C. racemosa* var. *cylindracea* (Sonder) Verlaque, Huisman and Boudouresque [30]. The vector of introduction in the Mediterranean Sea is unknown. It was initially hypothesized to be a Lessepsian immigrant, or a hybrid between *C. racemosa* var. *turbinata* and an unknown tropical variety [31] until molecular analyses identified a possible source population around the Australian-Pacific area [30–32]. It can invade different habitats, grow rapidly, and spread though fragmentation and propagation [33,34], leading to profound structural and functional alterations of native benthic assemblages and fish metabolism [9,35,36]. In the Mediterranean, the production of secondary metabolites by the alga changes seasonally and the highest levels have been observed during the period of vegetative algal growth (summer and autumn) [37]. Moreover Blažina et al. [20] observed that in *C. racemosa* from sheltered sites generally small variations in total lipids were found.

Here the chemical characterization of *C. cylindracea* lipidic extract was carried out by means of the advanced analytical technique of multinuclear and multidimensional NMR spectroscopy. In addition, the presence of secondary metabolites in the seaweed extract including bioactive compounds with antimicrobial and antioxidant activities was assessed. Since eradication has been recently suggested as a powerful action to protect areas highly impacted by invasive species, the secondary metabolites were investigated with the perspective of using *C. cylindracea* in eradication programs to support biotechnology and recycling options.

2. Results

2.1. NMR Spectroscopy

The 1D ^1H (Figure 1) spectrum in spectrum in CDCl_3 of the algal lipid fractions shows the characteristic signals of fatty acids (FAs), such as saturated (SAFAs) and unsaturated (UFAs) fatty acids, as well as caulerpin and other metabolites. At low frequencies the singlet at 0.66 ppm, which in the ^1H - ^{13}C HSQC shows cross peak correlation with the carbon at 11.8 ppm, was attributed to a methyl group of sterols.




Figure 1. ^1H NMR in CDCl_3 of *C. cylindracea* lipidic extract.

The signals at 0.85 ppm were assigned to the methyl groups of all FAs except ω_3 , which give a triplet at 0.95 ppm. The large signal at 1.21 and 1.50 ppm was attributed to alkyl chain and methylene group in β position to $\text{C}=\text{O}$ of all fatty acids. The large signals between 1.94 and 2.12 ppm and the signals at 5.35 ppm were attributed to methylene groups in α position with respect to vinyl groups and vinyl groups of all UFAs, respectively. The methylene groups in α to $\text{C}=\text{O}$ of all FAs resonate at 2.29 ppm. The bisallylic protons, characteristic of fatty acids with two or more double bonds, give signals at 2.78 ppm.

In addition, the characteristic spin system of poly- β -hydroxybutyrate (PHB) was identified in extract by 2D ^1H COSY spectra (two signals at 2.45 and 2.58 ppm, attributed to the methylene group, coupled with the methyl group at 1.26 ppm and the methine at 5.23 ppm). By 1D ^1H and 2D ^1H COSY spectra of all fractions different pattern signals for esterified glycerols were observed. The coupling system connecting the multiplet at 5.25 ppm and the doublet of doublet at 4.27 and 4.12 ppm was assigned to the CH *sn2* and the two CH_2 *sn1* and *sn3* of triacylglycerol. However, by 1D ^{13}C -NMR a higher intensity of signals at 179 ppm, attributed to carbonyl groups ($\text{C}=\text{O}$) of free fatty acids (FFAs) with respect to the signals in the range 173–174 ppm assigned to carbonyl groups of esterified fatty acids (EFAs) was observed.

In the 1D ^1H NMR spectrum at high frequencies the aromatic signals of caulerpin were observed. The doublets at 7.41 and 7.28 ppm and the multiplets at 7.16 and 7.07 ppm were assigned to the bis-indolic ring of caulerpin, whereas the singlets at 9.22, 8.04, and 3.88 ppm were assigned to NH of bis-indole, $=\text{CH}-$ of central ring and ester methyl groups, respectively. The signals in the high frequencies region between 11.2 and 8.5 ppm were attributed to methine bridge protons of chlorophyll and pheophytins. These signals were confirmed by the presence at negative value of chemical shift (-1.5 to -2.00 ppm) of peaks corresponding to the N-H protons of the tetrapyrrole ring. The signals in the range 6.00 to 6.70 ppm were assigned to conjugated double bonds of carotenoids. Isomers that are all *trans* are prevalent in fresh matter and the quantities of *cis* isomers increase after thermal processing.

2.2. Antioxidant Activity

The antioxidant activity of the lipidic extract from *C. cylindracea* assayed by TEAC and ORAC assays is reported in Table 1. Antioxidant capacity of the seaweed extract measured by the ORAC assay was eight times higher than the activity measured by the TEAC assay.

Table 1. Antioxidant activity of *C. cylindracea* lipidic extract assayed by TEAC and ORAC assays.

ORAC value	552.14 ± 69.13 μmol Trolox Equivalent/g extract
TEAC value	70.03 ± 2.67 μmol Trolox Equivalent/g extract

Data are the mean \pm SD ($n = 3$).

2.3. Antimicrobial Activity

The results of antimicrobial activity of *Caulerpa cylindracea* lipidic extract towards the utilized microbial strains are shown in Table 2. Yeasts were not affected in their growth by *C. cylindracea* lipidic extract. Moreover, the extract did not show antibacterial activity against *Enterococcus* sp., *Escherichia coli*, *Staphylococcus* sp., and *Streptococcus* sp. By contrast, the degree of inhibition produced by the lipidic extract on some *Vibrio* species was quantified. In particular, using discs with 100 μ L of algal extract, *V. fischeri*, *V. inusitatus*, and *V. litoralis* were the most inhibited (diameter of growth inhibition = 0.9 cm). A lower percentage of inhibition was measured on *V. aestuarianus* (0.85 cm), *V. mediterranei* (0.8 cm), and *V. vulnificus* (0.8 cm).

Table 2. Antimicrobial activity of *C. cylindracea* lipidic extract.

Microbial Strain	Diameter of Growth Inhibition (cm)						
	10 μ L	20 μ L	30 μ L	40 μ L	60 μ L	80 μ L	100 μ L
<i>Candida albicans</i>	0	0	0	0	0	0	0
<i>Candida glabrata</i>	0	0	0	0	0	0	0
<i>Enterococcus</i> sp.	0	0	0	0	0	0	0
<i>Escherichia coli</i>	0	0	0	0	0	0	0
<i>Staphylococcus</i> sp.	0	0	0	0	0	0	0
<i>Streptococcus</i> sp.	0	0	0	0	0	0	0
<i>Vibrio aestuarianus</i>	0.7	0.7	0.7	0.7	0.7	0.8	0.85
<i>Vibrio campbelli</i>	0	0	0	0	0	0	0
<i>Vibrio carchariae</i>	0	0	0	0	0	0	0
<i>Vibrio diazotrophicus</i>	0	0	0	0	0	0	0
<i>Vibrio fischeri</i>	0.8	0.8	0.8	0.8	0.8	0.8	0.9
<i>Vibrio fluvialis</i>	0	0	0	0	0	0	0
<i>Vibrio furnissi</i>	0	0	0	0	0	0	0
<i>Vibrio harveyi</i>	0	0	0	0	0	0	0
<i>Vibrio inusitatus</i>	0.8	0.8	0.8	0.8	0.8	0.9	0.9
<i>Vibrio litoralis</i>	0.8	0.8	0.8	0.8	0.8	0.8	0.9
<i>Vibrio mediterranei</i>	0	0	0.7	0.7	0.7	0.8	0.8
<i>Vibrio natriegens</i>	0	0	0	0	0	0	0
<i>Vibrio ordalii</i>	0	0	0	0	0	0	0
<i>Vibrio salmonicida</i>	0	0	0	0	0	0	0
<i>Vibrio splendidus</i> II	0	0	0	0	0	0	0
<i>Vibrio vulnificus</i>	0.8	0.8	0.8	0.8	0.8	0.8	0.8

3. Discussion

In the present study, the lipidic extract of the Mediterranean invasive alga *C. cylindracea* was analyzed by multinuclear and multidimensional NMR spectroscopy and its antioxidant, antibacterial, and antifungal activities have been evaluated.

Some important issues can be inferred from the obtained results:

By the NMR analysis the ^1H NMR spectrum in CDCl_3 of *C. cylindracea* algal lipid fraction interestingly, for the first time, showed the presence of polyhydroxybutyrate (PHB), a natural, linear biodegradable, and biocompatible polymer belonging to the polyesters group of bioplastics. PHB is synthesized by microorganisms as a form of energy-storage granules and utilized when the other energy sources are not available [38]. PHB has been already evidenced in the red algae *Plocamium cartilagineum* (Linnaeus) Dixon [39], *Gracilariopsis longissima* [17], and *Cladophora rupestris* [40]. Due to the rapid degradation and the non-toxicity of the final products, PHB represents an important ingredient for the production of polymers used in various biotechnological applications such as in the production of several medical devices and compostable plastic [41,42]. As crude oil reserves decrease throughout the world, petroleum based plastics are becoming less economical. Moreover, petroleum based plastics cannot be considered environmentally friendly due to their resistance to

natural or biological decomposition. In this framework, *Caulerpa cylindracea* could represent one of the alternatives for the production of bioplastics because it is an excellent renewable resource due to its high bioinvasion potential and high growth rate. The technology development for seaweeds-based bioplastics are still in the research phase. The challenge is to have significant advancements in the bioplastics industries to make seaweed bioplastics a concrete alternative in the future as already underlined in a recent review by Noreen et al. [43].

The NMR analysis besides other metabolites also showed the characteristic signals of SAFAs, UFAs, FFAs, and EFAs. NMR has indeed the desirable property of providing a global profiling tool for monitoring rapidly the molecular components of marine organisms [44]. Fatty acids are compounds usually bound to other molecules including glycerol, sugars, or phosphate groups thus constituting the lipids. As already reported for other algal lipid extracts [39,45], the presence of monogalactosyl diacylglycerol (MGDG) was recognized in *C. cylindracea* by NMR spectroscopy. Our results are in accordance with other studies on *C. racemosa* and other seaweeds confirming their nutritional value [21]. In particular, seaweeds are low in calories, have high content of dietary fiber, are a good source of polyunsaturated fatty acids DHA and EPA, and may contain proteins with an amino acid profile of interest [46]. Apart from the importance of fatty acids for diet, their ability to interfere with bacterial growth and survival has been established in several organisms including seaweeds [40,47]. This is noteworthy since in this study we observed for the first time an antibacterial activity of *C. cylindracea* against *Vibrio aestuarianus*, *V. fischeri*, *V. inusitatus*, *V. littoralis*, *V. mediterranei*, and *V. vulnificus*. This is a critical result considering that aquaculture is emerging as the fastest growing food-producing industry in the world because of the increasing demand for food fish consumption. However, the intensive culture of food fish has led to outbreaks of various microbial diseases, resulting in annual economic losses to the aquaculture industry estimated at billions of dollars worldwide [48]. Bacteria, mainly from the genus *Vibrio*, have been identified as the etiological agents responsible for the most common disease outbreaks in fish and shellfish causing mortality in several countries [49]. Disease outbreaks caused by several *Vibrio* species including *V. aestuarianus*, *V. mediterranei*, and *V. vulnificus* are called vibriosis [50–52]. Treatments of infected fish with antibiotic-mediated food are a common practice but have led to the development of bacterial antibiotic resistance, resulting in a higher dose requirement for effective control and a consequent increase of chemical residues released into the environment posing serious risks to animal and human health. Therefore, there is currently an increasing request for more environment-friendly alternatives to conventional antibiotics with similar or enhanced properties for aquatic animals. In the last 20 years, there has been an increasing interest in using various seaweed extracts as prophylactic and/or therapeutic agents in aquaculture [53]. Thus, the ability of the *C. cylindracea* lipidic extract to act against aquatic pathogens at a concentration of 5 mg/mL, evidenced in the present study, highlights its potential exploitation as a source of antibacterial compounds, of great importance in the control of disease in the mariculture industry, which is a largely unexplored field of research.

The seaweed extract did not show activity against the yeasts *C. albicans* and *C. glabrata*, and against the bacterial strains *Enterococcus* sp., *E. coli*, *Staphylococcus* sp., *Streptococcus* sp. It is well known that the antibacterial activity can be affected by many factors and that the method of extraction is one of the most important ones since different compounds, capable of acting on different bacterial strains, can be obtained by different extraction procedures [53,54]. The substances isolated from green, brown, and red algae showing powerful antimicrobial activity belong to different classes and include polysaccharides, fatty acids, phlorotannins, pigments, lectins, alkaloids, terpenoids, and halogenated compound. As an explanation, *C. racemosa* ethyl acetate extract exhibits antibacterial activity against *Enterococcus faecalis*, *Staphylococcus aureus*, *Bacillus cereus*, and *Escherichia coli* [54,55]. By contrast, methanolic extracts of *C. racemosa* shows antibacterial activity against *Klebsiella pneumoniae*, *Enterobacter aerogenes*, *Pseudomonas aeruginosa*, *Micrococcus luteus*, *Enterobacter faecalis*, *Streptococcus faecalis*, *Staphylococcus aureus*, and *Bacillus subtilis* [26,38]. In the present study, we evaluated the antibacterial

activity of the *C. cylindracea* lipidic extract but further studies are needed to evaluate whether the crude extract or aqueous extract lead to evidence other relevant biological activities.

The seaweed lipidic extract exerts an important antioxidant activity. Even though a comparison of antioxidant activity across seaweeds is difficult due to seasonal, environmental, and genetic variations, the obtained ORAC mean value of 552 μm Trolox equivalent/g extract for *C. cylindracea* is similar to those of lipid-soluble extracts of *Macrocystis pyrifera* (462 μm Trolox equivalent/g extract) and *Ecklonia radiata* (363 μm Trolox equivalent/g extract). This is noteworthy since these seaweeds, with powerful activity, have been recently proposed as potential sources of natural antioxidants instead of chemical antioxidant [56].

The TEAC and ORAC methods were simultaneously utilized in the present work since many studies stress the need of adopting at least two approaches to take into account the different mechanisms of antioxidant action [57,58], as single assay may disregard some radical sources or all antioxidants [59]. The methods utilized here are among the most popular assays and differ for their reaction mechanisms: TEAC is an electron transfer (ET) based method, while ORAC is based on hydrogen atom transfer (HAT) method. Other studies have already been conducted on the antioxidant activity of *C. racemosa* [21,60]. The utilized methodology is generally different across studies so that a comparison of seaweed antioxidant activity is challenging. The results obtained in the present study indicate that the ORAC values were higher in the measurement of antioxidant capacity of the lipidic extract of *C. cylindracea* than the TEAC. These results are in accordance with literature data indicating that carotenoids contain a chain of isoprene residues bearing numerous conjugated double bonds and are mostly involved in the scavenging of two of the reactive oxygen species, singlet molecular oxygen (O_2), and peroxy radicals ($\text{ROO}\bullet$). The HAT-based antioxidant capacity assays, like ORAC, utilize a radical initiator to generate peroxy radicals and measure the antioxidant activities of antioxidant molecules against peroxy radicals [61]. Then the antioxidant activity of carotenoids is higher when measured with ORAC assay on comparison to TEAC assay. By contrast, the antioxidant activity of phenolic acids is similar using both methods [62].

Interestingly, the characterization of *C. cylindracea* lipidic extract by 1D and 2D NMR spectroscopy assessed the presence of several types of molecules known for their antioxidant activity. Carotenoids, which produce signals in the 6.00–6.70 ppm range, were identified as well as the signals of chlorophylls and its thermal by-product, metal-free pheophytins. Both chlorophylls and pheophytins have demonstrated protective activity against auto-oxidation of vegetable oils in the dark [63,64]. Endo et al. [63] suggested that chlorophyll derivatives may act as electron donors as evidenced by their ability to reduce free radicals such as 1,1-diphenyl-2-picrylhydrazyl. Moreover, chlorophyll *a* was shown to act synergistically with vitamin E by quenching tocopherol radicals, thereby enhancing the observed antioxidant effects of vitamin E [65,66]. The ORAC specificity, the medium polarity of the extraction solvent, together with the content of carotenoids, chlorophylls, pheophytins, and sterols could explain the higher ORAC value of the *Caulerpa* lipidic extract compared to those obtained by TEAC according with other studies on *Caulerpa* [67,68]. Previous studies highlighted greater antioxidant activities of the green seaweed *C. racemosa* from Malaysian North Borneo compared to other brown and red seaweeds, observing that the phenolic compounds were involved in the antioxidant activity [69]. Further studies conducted on *C. racemosa* highlighted that the major contributors to the antioxidant activities are medium polarity phenolic compounds [70,71]. The further identification, characterization, and isolation of the compounds involved in *C. cylindracea* antioxidant activity might contribute to the employment of algal extracts in disease treatments related to oxidative stress, taking into account that, recently, an interest in natural antioxidants has increased because they are safer than synthetic antioxidants.

In conclusion, our findings open a new research area on the possible employment of the examined seaweed in the biotechnological field as source of bioactive natural products including antibiotics, antioxidants, fatty acids, and PHB potentially carrying benefits to human and marine organisms' health. At present, eradication of recent alien introduction [72] has been proposed as a promising management

action to assist the recovery of highly invaded areas under protection regime such as Marine Protected Areas. In this framework, the overproduced biomass of the invasive seaweed *C. cylindracea* could transform into a gain offering a potential tool with recycling options in eradication programs.

4. Materials and Methods

4.1. Study Site and Species Collection

Caulerpa cylindracea was collected in the Marine Protected Area of Torre Guaceto located in the Mediterranean Sea (Southern Adriatic Sea, Brindisi, Italy) at the depth of 8–10 m on the rocky bottom, during the season of maximum growth of the species when the alga dominates the benthic assemblages forming continuous dense meadows across the areas. *Caulerpa cylindracea* shows a seasonal cycle with a period of vegetative growth approximately between June and November alternated with a period of vegetative rest (a quasi-complete withdrawal) approximately from December to May [73,74]. Seasonal variations in the growth rate were highly significant: during the maximum development the biomass is $82 \pm 3 \text{ g} \cdot \text{DW} \cdot \text{m}^{-2}$ and length of stolons $1162 \pm 86 \text{ m} \cdot \text{m}^{-2}$. By contrast, during the minimum development the biomass is reduced to $0.3 \pm 0.1 \text{ g} \cdot \text{DW} \cdot \text{m}^{-2}$ and length of stolons to $3 \pm 1 \text{ m} \cdot \text{m}^{-2}$ [74]. Three replicates of about 500 g of fresh material were collected by SCUBA diving. All the harvested material was transferred into aseptic containers to the laboratory under controlled temperature (4 °C). The species was identified on the basis of morphological features. In detail, the following morphological features of the thalli were analyzed: height, width and attachment to stolons of the fronds; height, diameter, shape, and arrangement of the ramuli; diameter of the stolons; length, width, spacing, and morphology of the rhizoids. Algal morphological features fit the description given by several authors for the invasive species *C. cylindracea* [29,36,75–77].

4.2. Preparation of Lipidic Extracts from the Macroalga

Algae samples were cleaned of epiphytes and other marine organisms with a mixture of ethanol and (40%) and sodium hypochlorite (1%) for 10 s [78] and necrotic parts were removed. The samples were further rinsed with sterile water to remove any associated debris. The freshly cleaned material was air-dried and powdered, then 3 g of each sample was extracted in 150 mL of chloroform/methanol (2:1 at 55–60 °C for 24 h) using a soxhlet apparatus. Extraction solvents were evaporated under vacuum at controlled temperature. 5 mg of extract was then dissolved in 1 mL of absolute ethanol (95%; by J.T. Baker), and assayed for antimicrobial and antioxidant activity.

4.3. NMR Spectroscopy

The lipid fraction of *C. cylindracea* was characterized by ^1H and ^{13}C 1D and 2D NMR spectroscopy with the same methodology already reported in Stabili et al. [47]. 1D ^1H and ^{13}C , 2D ^1H Jres, ^1H COSY, ^1H – ^{13}C HSQC, and ^1H – ^{13}C HMBC spectra were recorded at 298.15 K on a Bruker Avance III 400 MHz spectrometer using CDCl_3 as solvent and chemical shift was referenced to TMS by the residual protic solvent peaks as internal references ($^1\text{H} = 7.24 \text{ ppm}$; $^{13}\text{C} = 77.0 \text{ ppm}$). High resolution ^{13}C -NMR spectra were acquired semi-quantitatively [79], with short relaxation times, and high number scans, to achieve sufficient S/N ratio to calculate the integrals. The following parameters were used: 64 K data points, spectral width of 20,161.291 Hz, 16 K scans with a 0.5 s repetition delay, and 60° at ^{13}C excitation pulse. The acquisition and processing of spectra were performed using the software Topspin 2.1 (Bruker Biospin). Resonances of fatty acids and metabolites were assigned on the basis of literature data [40,47,80,81].

4.4. Antioxidant Activity

4.4.1. Oxygen Radical Absorbance Capacity Assay (ORAC)

For ORAC the method of Davalos et al. [82] was used. Extracts were diluted with 75 mM phosphate buffer (pH 7.4). The assay was carried out in black-walled 96-well plates (Greiner-Bio One) and each well contained a final volume of 200 μ L. To each well 20 μ L of extract and 120 μ L of fluorescein (FL; 70 nM final concentration) were added and the plate was incubated at 37 °C for 15 min. The AAPH (60 μ L; 12 mM final concentration) was added to each well and fluorescence intensity was estimated using an Infinite200 Pro plate reader (Tecan, Männedorf, Switzerland), every minute for a total of 80 min using an excitation wavelength of 485/9 nm and an emission wavelength of 535/20 nm. A standard curve was constructed using 6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid (Trolox, Sigma-Aldrich, Oakville, ON, Canada, 1.5–10.5 μ M). A blank (fluorescein + AAPH) using phosphate buffer instead of the antioxidant solution was carried out in each assay. Results were determined by using Magellan v 7.2 software (Tecan, Männedorf, Switzerland), on the basis of the difference in area under the curve between the control and the sample and expressed as μ moles of Trolox equivalents (TE) per g of lipidic extract. All the reaction mixtures were prepared in triplicate and at least three independent assays were performed for each sample.

4.4.2. Trolox Equivalent Antioxidant Capacity Assay (TEAC)

The TEAC assay was performed as previously described by Re et al. [83] with minor modifications to adapt the assay to a microplate reader. Briefly 2,2'-azinobis (3-ethylbenzothiazoline-6-sulfonic acid) diammonium salt (ABTS, Sigma-Aldrich) radical cations were prepared by mixing an aqueous solution of potassium persulfate 2.45 mM (final concentration) and an aqueous solution of ABTS 7 mM (final concentration) and allowing it to stand in the dark at room temperature for 12–16 h, before use. The ABTS radical cation solution was diluted in PBS (pH 7.4) to an absorbance of 0.40 at 734 nm \pm 0.02; this value was adopted to obtain about 80% of maximum inhibition of the blank absorbance using the highest concentration of the Trolox standard curve. Trolox was used as antioxidant standard and a standard calibration curve was constructed for Trolox (0–16 μ M). After addition of 200 μ L of diluted ABTS to 10 μ L of Trolox standard or extracts diluted in PBS, in each well of a 96 well-plate (Costar), the absorbance reading at 734 nm was taken 6 min after initial mixing using an Infinite200 Pro plate reader (Tecan, Männedorf, Switzerland). Appropriate solvent blanks were run in each plate. The lipidic extract was assayed in at least three separate dilutions and in triplicate. The percentage inhibition of absorbance at 734 nm is calculated and plotted as a function of concentration of Trolox and the TEAC value expressed as Trolox equivalent (in μ molar) per g of lipidic extract, using Magellan v 7.2 software.

4.5. Test Microorganisms

The antibacterial activity was tested on six human pathogenic microbial strains: *Candida albicans*, *Candida glabrata*, *Enterococcus* sp., *Escherichia coli*, *Staphylococcus* sp., *Streptococcus* sp., and several *Vibrio* strains isolated from marine environment: *Vibrio aestuarinus*, *Vibrio campbellii*, *Vibrio carchariae*, *Vibrio diazotrophicus*, *Vibrio fischeri*, *Vibrio fluvialis*, *Vibrio furnissii*, *Vibrio harveyi*, *Vibrio inusitatus*, *Vibrio littoralis*, *Vibrio mediterranei*, *Vibrio natriegens*, *Vibrio ordalii*, *Vibrio salmonicida*, *Vibrio splendidus* II, *Vibrio vulnificus*. The tested strains were isolated and identified from seawater samples of the Mar Piccolo of Taranto as previously reported by Cavallo et al. [84] and Stabili et al. [85].

4.6. Antimicrobial Activity

Antimicrobial activity was evaluated using the Kirby Bauer method [86]. Sterile paper discs, 6 mm in diameter (AA; Whatman International Ltd., Maidstone, Kent, UK), were impregnated with 10, 20, 30, 40, 60, 80, 100 μ L of each extract and left to air-dry at room temperature for 4 h, as already described by Cavallo et al. [87]. By contrast, discs impregnated with an equivalent volume of carrier solvent were used as controls. Moreover, an 'extraction blank' as a negative control (MeOH/CHCl₃ extraction

with no algae, dry, then suspended in ethanol) was also used. For each assay, sterile medium-agar plates opportune for each selected bacterial and fungal strain tested were seeded with 100 µL of microbial suspension (about 10^8 CFU mL⁻¹) [88,89], using a sterile swab to give a uniform covering. Impregnated discs and controls were laid onto the agar surface, the *Vibrio* plates were then incubated for 24 h at 30 °C, the other tests were conducted at 37 °C. The clear zone around the discs was evidence of antibacterial activity. The diameter of the microbial growth inhibition was taken as the diameter of the clear zone (measured in centimeters).

Acknowledgments: The European Union's Horizon 2020 research for the project MERCES (Grant agreement No. 689518, <http://www.merces-project.eu>) and RITMARE Flagship Project funded by the Italian Ministry of University and Research are acknowledged. We thank the three anonymous reviewers for the revision process as well as for their constructive comments which proved to be particularly useful to improve the manuscript.

Author Contributions: L.S., S.F. and L.R. conceived and designed the experiments; L.S., M.I.A., R.A.C., S.A.D.P., C.G., M.N. and L.R. performed the experiments; L.S., S.F., M.I.A., R.A.C., S.A.D.P., F.P.F., C.G., M.N. and L.R. analyzed the data; L.S., S.F. and F.P.F. contributed reagents/materials/analysis tools; L.S., S.F. and L.R. wrote the paper.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Raniello, R.; Mollo, E.; Lorenti, M.; Gavagnin, M.; Buia, M.C. Phytotoxic activity of caulerpenyne from the Mediterranean invasive variety of *Caulerpa racemosa*: A potential allelochemical. *Biol. Invasions* **2007**, *9*, 361–368. [[CrossRef](#)]
2. Mollo, E.; Gavagnin, M.; Carbone, M.; Castelluccio, F.; Pozzone, F.; Roussis, V.; Templado, J.; Ghiselin, M.T.; Cimino, G. Factors promoting marine invasions: A chemoecological approach. *Proc. Natl. Acad. Sci. USA* **2008**, *105*, 4582–4586. [[CrossRef](#)] [[PubMed](#)]
3. Terlizzi, A.; Fellingine, S.; Lionetto, M.G.; Caricato, R.; Perfetti, V.; Cutignano, A.; Mollo, E. Detrimental physiological effects of the invasive alga *Caulerpa racemosa* on the Mediterranean white seabream *Diplodus sargus*. *Aquat. Biol.* **2011**, *12*, 109–117. [[CrossRef](#)]
4. Pusceddu, A.; Frascchetti, S.; Scopa, M.; Rizzo, L.; Danovaro, R. Meiofauna communities, nematode diversity and C degradation rates in seagrass (*Posidonia oceanica* L.) and unvegetated sediments invaded by the algae *Caulerpa cylindracea* (Sonder). *Mar. Environ. Res.* **2016**, *119*, 88–99. [[CrossRef](#)] [[PubMed](#)]
5. Rizzo, L.; Frascchetti, S.; Alifano, P.; Pizzolante, G.; Stabili, L. The alien species *Caulerpa cylindracea* and its associated bacteria in the Mediterranean Sea. *Mar. Biol.* **2016**, *163*, 1–12. [[CrossRef](#)]
6. Rizzo, L.; Frascchetti, S.; Alifano, P.; Tredici, M.S.; Stabili, L. Association of *Vibrio* community with the Atlantic Mediterranean invasive alga *Caulerpa cylindracea*. *J. Exp. Mar. Biol. Ecol.* **2016**, *475*, 129–136. [[CrossRef](#)]
7. Ciminiello, P.; Dell'Aversano, C.; Forino, M.; Tartaglione, L. Marine toxins in Italy: The more you look, the more you find. *Eur. J. Org. Chem.* **2014**, *2014*, 1357–1369. [[CrossRef](#)]
8. Gorbi, S.; Giuliani, M.E.; Pittura, L.; d'Errico, G.; Terlizzi, A.; Fellingine, S.; Grauso, L.; Mollo, E.; Cutignano, A.; Regoli, F. Could molecular effects of *Caulerpa racemosa* metabolites modulate the impact on fish populations of *Diplodus sargus*? *Mar. Environ. Res.* **2014**, *96*, 2–11. [[CrossRef](#)] [[PubMed](#)]
9. Fellingine, S.; Mollo, E.; Ferramosca, A.; Zara, V.; Regoli, F.; Gorbi, S.; Terlizzi, A. Can a marine pest reduce the nutritional value of Mediterranean fish flesh? *Mar. Biol.* **2014**, *161*, 1275–1283. [[CrossRef](#)]
10. Mollo, E.; Cimino, G.; Ghiselin, M.T. Alien biomolecules: A new challenge for natural product chemists. *Biol. Invasions* **2015**, *17*, 941–950. [[CrossRef](#)]
11. Villarreal-Gómez, L.J.; Soria-Mercado, I.E.; Guerra-Rivas, G.; Ayala-Sánchez, N.E. Antibacterial and anticancer activity of seaweeds and bacteria associated with their surface. *Rev. Biol. Mar. Oceanogr.* **2010**, *45*, 267–275. [[CrossRef](#)]
12. Pandian, P.; Selvamuthukumar, S.; Manavalan, R.; Parthasarthy, V. Screening of antibacterial and antifungal activities of red marine algae *Acanthaphora spicifera* (Rhodophyceae). *J. Biomed. Sci. Res.* **2011**, *3*, 444–448.
13. Senthilkumar, K.; Kim, S.K. Anticancer effects of fucoidan. *Adv. Food Nutr. Res.* **2014**, *72*, 195–213. [[PubMed](#)]
14. Chakraborty, K.; Joseph, D.; Joy, M.; Raola, V.K. Characterization of substituted aryl meroterpenoids from red seaweed *Hypnea musciformis* as potential antioxidants. *Food Chem.* **2016**, *212*, 778–788. [[CrossRef](#)] [[PubMed](#)]

15. Hardouin, K.; Bedoux, G.; Burlot, A.S.; Donnay-Moreno, C.; Bergé, J.P.; Nyvall-Collén, P.; Bourgoignon, N. Enzyme-assisted extraction (EAE) for the production of antiviral and antioxidant extracts from the green seaweed *Ulva armoricana* (Ulvales, Ulvophyceae). *Algal Res.* **2016**, *16*, 233–239. [[CrossRef](#)]
16. Jiménez-Escrig, A.; Sánchez-Muniz, F.J. Dietary fibre from edible seaweeds: Chemical structure, physicochemical properties and effects on cholesterol metabolism. *Nutr. Res.* **2000**, *20*, 585–598. [[CrossRef](#)]
17. Yang, Y.J.; Nam, S.J.; Kong, G.; Kim, M.K. A case-control study on seaweed consumption and the risk of breast cancer. *Br. J. Nutr.* **2010**, *103*, 1345–1353. [[CrossRef](#)] [[PubMed](#)]
18. Matsumura, Y. Nutrition trends in Japan. *Asia Pac. J. Clin. Nutr.* **2001**, *10*, S40–S47. [[CrossRef](#)] [[PubMed](#)]
19. Terrados, J.; Lopez-Jimenez, J.A. Fatty acid composition and chilling resistance in the green alga *Caulerpa prolifera* (Forsskal) Lamouroux (Chlorophyta, Caulerpales). *Biochem. Mol. Biol. Int.* **1996**, *39*, 863–869. [[CrossRef](#)] [[PubMed](#)]
20. Blažina, M.; Iveša, L.; Najdek, M. *Caulerpa racemosa*: Adaptive varieties studied by fatty acid composition (Northern Adriatic Sea, Vrsar, Croatia). *Eur. J. Phycol.* **2009**, *44*, 183–189. [[CrossRef](#)]
21. Kumar, M.; Gupta, V.; Kumari, P.; Reddy, C.R.K.; Jha, B. Assessment of nutrient composition and antioxidant potential of Caulerpaceae Seaweeds. *J. Food Compos. Anal.* **2010**, *24*, 270–278. [[CrossRef](#)]
22. Paul, N.A.; Neveux, N.; Magnusson, M.; de Nys, R. Comparative production and nutritional value of “sea grapes”—The tropical green seaweeds *Caulerpa lentillifera* and *C. racemosa*. *J. Appl. Phycol.* **2014**, *26*, 1833. [[CrossRef](#)]
23. Kumar, C.S.; Ganesan, P.; Suresh, P.V.; Bhaskar, N. Seaweeds as a Source of Nutritionally Beneficial Compounds—A Review. *J. Food Sci. Technol.* **2008**, *45*, 1–13.
24. Gupta, S.; Rajauria, G.; Abu-Ghannam, N. Study of the microbial diversity and antimicrobial properties of Irish edible brown seaweeds. *Int. J. Food Sci. Technol.* **2010**, *45*, 482–489. [[CrossRef](#)]
25. Ballesteros, E.; Martin, D.; Uriz, M.J. Biological activity of extracts from some Mediterranean macrophytes. *Bot. Mar.* **1992**, *35*, 481–486. [[CrossRef](#)]
26. Kandhasamy, M.; Arunachalam, K.D. Evaluation of in vitro antibacterial property of seaweeds of southeast coast of India. *Afr. J. Biotechnol.* **2008**, *7*, 1958–1961.
27. Jebasingh, S.E.J.; Rosemary, S.; Elaiyaraja, S.; Sivaraman, K.; Lakshmikandan, M.; Murugan, A.; Raja, P. Potential antibacterial activity of selected green and red seaweeds. *J. Pharmac. Biomed. Sci.* **2011**, *5*, 1–7.
28. Etcherla, M.; Narasimha Rao, G.M. In vitro study of antimicrobial activity in marine algae *Caulerpa taxifolia* and *Caulerpa racemosa* (C. Agardh). *Int. J. Appl. Biol. Pharm. Technol.* **2014**, *5*, 57–72.
29. Belton, G.S.; Prud’homme van Reine, W.F.; Huisman, J.M.; Draisma, S.G.A.; Gurgel, C.F.D. Resolving phenotypic plasticity and species designation in the morphology challenging *Caulerpa racemosa-peltata* complex (Caulerpaceae, Chlorophyta). *J. Phycol.* **2014**, *50*, 32–54. [[CrossRef](#)] [[PubMed](#)]
30. Verlaque, M.; Durand, C.; Huisman, J.M.; Boudouresque, C.F.; Le Parco, Y. On the identity and origin of the Mediterranean invasive *Caulerpa racemosa* (Caulerpales, Chlorophyta). *Eur. J. Phycol.* **2003**, *38*, 325–339. [[CrossRef](#)]
31. Durand, C.; Manuel, M.; Boudouresque, C.F.; Meinesz, A.; Verlaque, M.; Le Parco, Y. Molecular data suggest a hybrid origin for the invasive *Caulerpa racemosa* (Caulerpales, Chlorophyta) in the Mediterranean Sea. *J. Evol. Biol.* **2002**, *15*, 122–133. [[CrossRef](#)]
32. Sauvage, T.; Payri, C.; Draisma, S.G.A.; Prud’homme van Reine, W.F.; Verbruggen, H.; Belton, G.S.; Gabriel, D.; Sherwood, A.R.; Fredericq, S.; Gurgel, C.F.D. Molecular diversity of the *Caulerpa racemosa*–*Caulerpa peltata* complex (Caulerpaceae, Bryopsidales) in New Caledonia, with new Australasian records for *C. racemosa* var. *cylindracea*. *Phycologia* **2013**, *52*, 6–13. [[CrossRef](#)]
33. Wright, J.T.; Davis, A.R. Demographic feedback between clonal growth and fragmentation in an invasive seaweed. *Ecology* **2006**, *87*, 1744–1754. [[CrossRef](#)]
34. Smith, C.M.; Walters, L.J. Fragmentation as a strategy for *Caulerpa* species: Fates of fragments and implications for management of an invasive weed. *Mar. Ecol.* **1999**, *20*, 307–319. [[CrossRef](#)]
35. Holmer, M.; Marbà, N.; Lamote, M.; Duarte, C.M. Deterioration of Sediment Quality in Seagrass Meadows (*Posidonia oceanica*) Invaded by Macroalgae (*Caulerpa* sp.). *Estuaries Coasts* **2009**, *32*, 456–466. [[CrossRef](#)]
36. Cebrian, E.; Linares, C.; Marchal, C.; Garrabou, J. Exploring the effects of invasive algae on the persistence of gorgonian populations. *Biol. Invasions* **2012**, *14*, 2647–2656. [[CrossRef](#)]

37. Box, A.; Deudero, S.; Sureda, A.; Blanco, A.; Alòs, J.; Terrados, J.; Grau, A.M.; Riera, F. Diet and physiological responses of *Spondyliosoma cantharus* (Linnaeus, 1758) to the *Caulerpa racemosa* var. *cylindracea* invasion. *J. Exp. Mar. Biol. Ecol.* **2009**, *380*, 11–19. [[CrossRef](#)]
38. Gupta, M.P.; Gómez, N.E.; Santana, A.I.; Solis, P.N.; Palacios, G. Antimicrobial activity of various algae of the Panamanian Atlantic coast. *Rev. Med. Panama* **1991**, *16*, 64–68. [[PubMed](#)]
39. Dawczynski, C. Amino acids, fatty acids, and dietary fibre in edible seaweed products. *Food Chem.* **2007**, *103*, 891–899. [[CrossRef](#)]
40. Stabili, L.; Acquaviva, M.I.; Biandolino, F.; Cavallo, R.A.; De Pascali, S.A.; Fanizzi, F.P.; Narracci, M.; Cecere, E.; Petrocelli, A. Biotechnological potential of the seaweed *Cladophora rupestris* (Chlorophyta, Cladophorales) lipidic extract. *New Biotechnol.* **2014**, *31*, 436–444. [[CrossRef](#)] [[PubMed](#)]
41. Luengo, J.M.; Garcia, B.; Sandoval, A.; Naharro, G.; Olivera, E.R. Bioplastics from microorganisms. *Curr. Opin. Microbiol.* **2003**, *6*, 251–260. [[CrossRef](#)]
42. Pachekoski, W.M.; Marcondes Agnelli, J.A.; Belem, L.P. Thermal, mechanical and morphological properties of poly (hydroxybutyrate) and polypropylene blends after processing. *Mater. Res.* **2009**, *12*, 159–164. [[CrossRef](#)]
43. Noreen, A.; Zia, K.M.; Zuber, M.; Ali, M.; Mujahid, M. A critical review of algal biomass: A versatile platform of bio-based polyesters from renewable resources. *Int. J. Biol. Macromol.* **2016**, *86*, 937–949. [[CrossRef](#)] [[PubMed](#)]
44. De Pascali, S.A.; Del Coco, L.; Feline, S.; Mollo, E.; Terlizzi, A.; Fanizzi, F.P. ¹H NMR spectroscopy and MVA analysis of *Diplodus sargus* eating the exotic pest *Caulerpa cylindracea*. *Mar. Drugs* **2015**, *13*, 3550–3566. [[CrossRef](#)] [[PubMed](#)]
45. Schmid, M.; Guihéneuf, F.; Stengel, D. Fatty acid contents and profiles of 16 macroalgae collected from the Irish Coast at two seasons. *J. Appl. Phycol.* **2014**, *26*, 451–463. [[CrossRef](#)]
46. Holdt, S.L.; Kraan, S. Bioactive compounds in seaweed: Functional food applications and legislation. *J. Appl. Phycol.* **2011**, *23*, 543–597. [[CrossRef](#)]
47. Stabili, L.; Acquaviva, M.I.; Biandolino, F.; Cavallo, R.A.; De Pascali, S.A.; Fanizzi, F.P.; Narracci, M.; Petrocelli, A.; Cecere, E. The lipidic extract of the seaweed *Gracilariopsis longissima* (Rhodophyta, Gracilariales): A potential resource for biotechnological purposes? *New Biotechnol.* **2012**, *29*, 443–450. [[CrossRef](#)] [[PubMed](#)]
48. Pridgeon, J.W.; Klesius, P.H. Major bacterial diseases in aquaculture and their vaccine development. *Anim. Sci. Rev.* **2012**, *7*, 1–16. [[CrossRef](#)]
49. Sudheesh, P.S.; Al-Ghabshi, A.; Al-Mazrooei, N.; Al-Habsi, S. Comparative pathogenomics of bacteria causing infectious diseases in fish. *Int. J. Evol. Biol.* **2012**, *2012*, 1–16. [[CrossRef](#)] [[PubMed](#)]
50. Labreuche, Y.; Soudant, P.; Gonçalves, M.; Lambert, C.; Nicolas, J.L. Effects of extracellular products from the pathogenic *Vibrio aestuarianus* strain 01/32 on lethality and cellular immune responses of the oyster *Crassostrea gigas*. *Dev. Comp. Immunol.* **2006**, *30*, 367–379. [[CrossRef](#)] [[PubMed](#)]
51. Kushmaro, A.; Banin, E.; Loya, Y.; Stackebrandt, E.; Rosenberg, E. *Vibrio shiloi* sp. nov., the causative agent of bleaching of the coral *Oculina patagonica*. *Int. J. Syst. Evol. Microbiol.* **2001**, *51*, 1383–1388. [[CrossRef](#)] [[PubMed](#)]
52. Strom, M.S.; Paranjpye, R.N. Epidemiology and pathogenesis of *Vibrio vulnificus*. *Microbes Infect.* **2000**, *2*, 177–188. [[CrossRef](#)]
53. Vatsos, I.N.; Rebours, C. Seaweed extracts as antimicrobial agents in aquaculture. *J. Appl. Phycol.* **2015**, *27*, 2017–2035. [[CrossRef](#)]
54. Chandrasekaran, M.; Venkatesalu, V.; Raj, G.A. Antibacterial activity of selected marine macro algae against vancomycin resistant *Enterococcus faecalis*. *J. Coast. Life Med.* **2014**, *2*, 940–946.
55. Salem, W.M.; Galal, H.; Nasr El-deen, F. Screening for antibacterial activities in some marine algae from the red sea (Hurghada, Egypt). *Afr. J. Microbiol. Res.* **2011**, *5*, 2160–2167. [[CrossRef](#)]
56. Kindleysides, S.; Quek, S.Y.; Miller, M.R. Inhibition of fish oil oxidation and the radical scavenging activity of New Zealand sea weed extracts. *Food Chem.* **2012**, *133*, 1624–1631. [[CrossRef](#)]
57. Frankel, E.N.; Meyer, A.S. The problems of using one-dimensional methods to evaluate multifunctional food and biological antioxidants. *J. Sci. Food Agric.* **2000**, *80*, 1925–1941. [[CrossRef](#)]
58. Prior, R.L.; Cao, G. In vivo total antioxidant capacity: Comparison of different analytical methods. *Free Radic. Biol. Med.* **1999**, *27*, 1173–1181. [[CrossRef](#)]

59. Prior, R.L.; Wu, X.; Schaich, K. Standardized methods for the determination of antioxidant capacity and phenolics in foods and dietary supplements. *J. Agric. Food Chem.* **2005**, *53*, 4290–4302. [[CrossRef](#)] [[PubMed](#)]
60. Lakmal, H.C.; Samarakoon, K.W.; Lee, W.; Lee, J.H.; Abeytunga, D.T.U.; Lee, H.S.; Jeon, Y.J. Anticancer and antioxidant effects of selected Sri Lankan marine algae. *J. Natl. Sci. Found. Sri Lanka* **2014**, *42*, 315–323. [[CrossRef](#)]
61. Zulueta, A.; Esteve, M.A.; Frigola, A. ORAC and TEAC assay comparison to measure the antioxidant capacity of food product. *Food Chem.* **2009**, *114*, 310–316. [[CrossRef](#)]
62. Ou, B.; Huang, D.; Hampsch-Woodill, M.; Flanagan, J.A.; Deemer, E.K. Analysis of antioxidant activities of common vegetables employing oxygen radical absorbance capacity (ORAC) and ferric reducing antioxidant power (FRAP) assays: A comparative study. *J. Agric. Food Chem.* **2002**, *50*, 3122–3128. [[CrossRef](#)] [[PubMed](#)]
63. Endo, Y.; Usuki, R.; Kaneda, T. Antioxidant effects of chlorophyll and pheophytin on the autooxidation of oils in the dark. II. The mechanism of antioxidative action of chlorophyll. *J. Am. Oil Chem. Soc.* **1985**, *62*, 1387–1390. [[CrossRef](#)]
64. Gutierrez-Rosales, F.; Garrido-Fernandez, J.; Gallardo-Guerrero, L.; Gandul-Rojas, B.; Minguéz-Mosquera, M.I. Action of chlorophylls on the stability of the virgin olive oil. *J. Am. Oil Chem. Soc.* **1992**, *69*, 866–871. [[CrossRef](#)]
65. Le Tutour, B.; Brunel, C.; Quemeneur, F. Synergistic effect of chlorophyll a on the oxidation properties of vitamin E. *New J. Chem.* **1996**, *20*, 707–721.
66. Le Tutour, B.; Benslimane, F.; Gouleau, M.P.; Gouygou, J.P.; Saadan, B.; Quemeneur, F. Antioxidant and pro-oxidant activities of the brown algae, *Laminaria digitata*, *Himantalia elongata*, *Fucus vesiculosus*, *Fucus serratus* and *Ascophyllum nodosum*. *J. Appl. Phycol.* **1998**, *19*, 121–129. [[CrossRef](#)]
67. Pires-Cavalcante, K.M.D.S.; de Alencar, D.B.; de Sousa, M.B.; Sampaio, A.H.; Saker-Sampaio, S. Seasonal changes of α -tocopherol in green marine algae (*Caulerpa* genus). *J. Food Sci.* **2011**, *76*, C775–C781. [[CrossRef](#)] [[PubMed](#)]
68. Ornano, L.; Donno, Y.; Sanna, C.; Ballero, M.; Serafini, M.; Bianco, A. Phytochemical study of *Caulerpa racemosa* (Forsk.) J. Agarth, an invading alga in the habitat of La Maddalena Archipelago. *Nat. Prod. Res.* **2014**, *28*, 1795–1799. [[CrossRef](#)] [[PubMed](#)]
69. Matanjun, P.; Mohamed, S.; Mustapha, N.M.; Muhammad, K.; Ming, C.H. Antioxidant activities and phenolics content of eight species of seaweeds from north Borneo. *J. Appl. Phycol.* **2008**, *20*, 367–373. [[CrossRef](#)]
70. Vinayak, R.C.; Sudha, S.A.; Chatterji, A. Bio-screening of a few green seaweeds from India for their cytotoxic and antioxidant potential. *J. Sci. Food Agric.* **2011**, *91*, 2471–2476. [[CrossRef](#)] [[PubMed](#)]
71. Li, Z.; Wang, B.; Zhang, Q.; Qu, Y.; Xu, H.; Li, G. Preparation and antioxidant property of extract and semipurified fractions of *Caulerpa racemosa*. *J. Appl. Phycol.* **2012**, *24*, 1527–1536. [[CrossRef](#)]
72. Anderson, L.W. California's reaction to *Caulerpa taxifolia*: A model for invasive species rapid response. *Biol. Invasions* **2005**, *7*, 1003–1016. [[CrossRef](#)]
73. Piazzini, L.; Ceccherelli, G.; Cinelli, F. Threat to macroalgal diversity: Effects of the introduced green alga *Caulerpa racemosa* in the Mediterranean. *Mar. Ecol. Prog. Ser.* **2001**, *210*, 161–165. [[CrossRef](#)]
74. Ruitton, S.; Verlaque, M.; Boudouresque, C.F. Seasonal changes of the introduced *Caulerpa racemosa* var. *cylindracea* (Caulerpales, Chlorophyta) at the northwest limit of its Mediterranean distribution. *Aquat. Bot.* **2005**, *82*, 55–70. [[CrossRef](#)]
75. Klein, J.; Verlaque, M. The *Caulerpa racemosa* invasion: A critical review. *Mar. Pollut. Bull.* **2008**, *56*, 205–225. [[CrossRef](#)] [[PubMed](#)]
76. Guillén, J.; Jiménez, S.; Martínez, J.; Triviño, A.; Múgica, Y.; Argilés, J.; Bueno, M. Expansion of the invasive algae *Caulerpa racemosa* var. *cylindracea* (Sonder) Verlaque, Huisman & Boudouresque, 2003 on the region of Valencia seabed. *Thalassas* **2010**, *26*, 135–149.
77. Verlaque, M.; Boudouresque, C.F.; Meinesz, A.; Gravez, V. The *Caulerpa racemosa* complex (Caulerpales, Ulvophyceae) in the Mediterranean Sea. *Bot. Mar.* **2000**, *43*, 49–68. [[CrossRef](#)]
78. Kientz, B.; Thabard, M.; Cragg, S.M.; Pope, J.; Hellio, C. A new method for removing microflora from macroalgal surfaces: An important step for natural product discovery. *Bot. Mar.* **2011**, *54*, 457–469. [[CrossRef](#)]
79. Standal, I.B.; Axelson, A.E.; Aursand, M. Differentiation of fish oils according to species by ^{13}C -NMR regiospecific analyses of triacylglycerols. *J. Am. Oil Chem. Soc.* **2009**, *86*, 401–407. [[CrossRef](#)]

80. Del Coco, L.; Papadia, P.; De Pascali, S.A.; Bressani, G.; Storelli, C.; Zonno, V.; Fanizzi, F.P. Comparison among different gilthead sea bream (*Sparus aurata*) farming systems: Activity of intestinal and hepatic enzymes and ^{13}C -NMR analysis of lipids. *Nutrients* **2009**, *1*, 291–301. [[CrossRef](#)] [[PubMed](#)]
81. Kamal, C.; Sethuraman, M.G. Caulerpin—A bis-Indole Alkaloid as a Green Inhibitor for the Corrosion of Mild Steel in 1 M HCl Solution from the Marine Alga *Caulerpa racemosa*. *Ind. Eng. Chem. Res.* **2012**, *51*, 10399–10407. [[CrossRef](#)]
82. Dávalos, A.; Gómez-Cordovés, C.; Bartolomé, B. Extending applicability of the oxygen radical absorbance capacity (ORAC-fluorescein) assay. *J. Agric. Food Chem.* **2004**, *52*, 48–54. [[CrossRef](#)] [[PubMed](#)]
83. Re, R.; Pellegrini, N.; Proteggente, A.; Pannala, A.; Yang, M.; Rice-Evans, C. Antioxidant activity applying an improvement ABTS radical cation decolorization assay. *Free Radic. Biol. Med.* **1999**, *26*, 1231–1237. [[CrossRef](#)]
84. Cavallo, R.A.; Stabili, L. Presence of vibrios in seawater and *Mytilus galloprovincialis* (Lam.) from the Mar Piccolo of Taranto (Ionian Sea). *Water Res.* **2002**, *36*, 3719–3726. [[CrossRef](#)]
85. Stabili, L.; Acquaviva, M.I.; Cavallo, R.A. *Mytilus galloprovincialis* filter feeding on the bacterial community in a Mediterranean coastal area (Northern Ionian Sea, Italy). *Water Res.* **2005**, *39*, 469–477. [[CrossRef](#)] [[PubMed](#)]
86. Bauer, A.W.; Kirby, W.M.; Sherris, J.C.; Turcck, M. Antibiotic susceptibility testing by a standardized single disk method. *Am. J. Clin. Pathol.* **1966**, *45*, 493–496. [[PubMed](#)]
87. Cavallo, R.; Acquaviva, M.; Stabili, L.; Cecere, E.; Petrocelli, A.; Narracci, M. Antibacterial activity of marine macroalgae against fish pathogenic *Vibrio* species. *Open Life Sci.* **2013**, *8*, 646–653. [[CrossRef](#)]
88. Ely, R.; Supriya, T.; Naik, C.G. Antimicrobial activity of marine organisms collected off the coast of South East India. *J. Exp. Biol. Ecol.* **2004**, *309*, 121–127. [[CrossRef](#)]
89. Kim, I.H.; Lee, D.G.; Lee, S.H.; Ha, J.M.; Ha, B.J.; Kim, S.K.; Lee, J.H. Antibacterial activity of *Ulva lactuca* against methicillin resistant *Staphylococcus aureus* (MRSA). *Biotechnol. Bioprocess Eng.* **2007**, *112*, 579–582. [[CrossRef](#)]



© 2016 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC-BY) license (<http://creativecommons.org/licenses/by/4.0/>).