

This article was downloaded by: [Cornell University Library]

On: 12 November 2014, At: 14:16

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number:

1072954 Registered office: Mortimer House, 37-41 Mortimer Street,
London W1T 3JH, UK



Annals and Magazine of Natural History: Series 4

Publication details, including instructions
for authors and subscription information:
<http://www.tandfonline.com/loi/tnah10>

XXXVIII.—Descriptions of five birds and a hare from Abyssinia

William T. Blanford F.G.S. C.M.Z.S.

Published online: 16 Oct 2009.

To cite this article: William T. Blanford F.G.S. C.M.Z.S. (1869)
XXXVIII.—Descriptions of five birds and a hare from Abyssinia ,
Annals and Magazine of Natural History: Series 4, 4:23, 329-330, DOI:
[10.1080/00222936908696065](https://doi.org/10.1080/00222936908696065)

To link to this article: <http://dx.doi.org/10.1080/00222936908696065>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly

or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

EXPLANATION OF PLATE XII.

- Fig. 1.* Two rows of teeth of *Janassa (C.) linguaeformis*, a little over the natural size, arranged in order, the anterior row merely indicated : *a*, central primary tooth ; *b*, root ; *c*, first lateral primary tooth ; *d*, second oblique ditto ; *e*, secondary or Petalodontoid form ; *f*, root of ditto.
- Fig. 2.* Primary tooth of *J. linguaeformis*, smooth variety, slightly enlarged : *a*, scoop-like cutting-margin ; *b*, grinding- or crushing-surface.
- Fig. 3.* Primary tooth of the same, a little enlarged ; worn variety, intermediate between the smooth variety and those much undulated.
- Fig. 4.* Diagram of profile of primary tooth : *a*, scoop-like cutting-margin ; *b*, crushing- or grinding-surface ; *c*, root.

XXXVIII.—*Descriptions of five Birds and a Hare from Abyssinia.* By WILLIAM T. BLANFORD, F.G.S., C.M.Z.S.

Hirundo aethiopica, sp. nov.

H. similis *H. albiculari*, Strickl. (Contrib. to Ornith. 1849, pl. 17), sed conspicue minor et torque pectorali interrupta, gutture pectoreque rufescenti-lavatis.
Long. tota 5·25, al. 4·3, rect. med. 1·55, ext. 2·3, tarsi 0·45, rostr. a fr. 0·3, a rict. 0·5 poll. Angl.

Syn. *Cecropis rufifrons*, auct., ex Abyssinia.

Hirundo albicularis, Strickl. apud Heuglin, Ornithologie N. O. Africa's, p. 113 (nec Strickland, l. c.).

Hab. in Abyssinia septentrionali, et in Nubia (teste Heuglin).

Ruticilla (?) fuscicaudata, sp. nov.

R. supra brunnescenti-fusca, uropygio magis rufescente ; remigibus reetricibusque fuscis, vix pallidiore marginatis ; macula anteroculari nigrescente, albido circumdata ; mento, gula, abdomineque medio sordide albis, pectore et hypochondriis cinerascentibus. Rostro pedibusque fuscis.

Long. tota circa 5·5 poll. Angl., al. 2·95, caud. 2·2, tars. 0·9, rostr. a fr. 0·45, a rict. 0·7.

Hab. in Abyssinia septentrionali.

The form of this bird resembles *Ruticilla*, the bill being similar and the tarsi smooth in front ; but the sombre plumage rather resembles that of a *Sylvia*. Its nearest allies are *R.* (*Saxicola*) *familiaris*, Stephens, and *R.* (*Erythacus*) *sinuata*, Schlegel. The tail is somewhat rounded, and the wing is less pointed than in *Ruticilla*.

Phylloscopus habessinicus, sp. nov.

Ph. *P. trochili* similis, sed supra magis viridescens, subtus isabellinus vix flavescens, cauda longiore.

Long. al. 2·5, caud. 2·05, tars. 0·78, rostr. a fr. 0·36, a rict. 0·5, tota circa 4·5 poll. Angl.

Hab. in provincia Habessinica "Tigré" dicta.

Alauda prætermissa, sp. nov.

A. supra fusco-umbrina, capitis, colli postice interscapuliique plumis late et pallide rufesceni-marginatis; dorso posteriore magis cinerascente, vix striolato; superciliis et gastræo toto isabellinis; genis, colli lateribus pectoreque saturationibus et fusco guttatis, regione parotica fuscescente, remigibus et tectricibus alarum umbrinis, primariis extus isabellino, secundariis cum tectricibus alarum rufo marginatis; remigibus omnibus intus versus basin rufofulvis; uropygio et rectricibus medianis rufescenti-umbrinis, exteris (pogonis internis basin versus exceptis) et pogonis externis secundarum rufo-isabellinis, cæteris cum partibus reliquis 4 externalium fumoso-nigricantibus. Caput suberistatum; rostrum supra fuscum, subtus pallidum; pedes carnei.

Long. al. 3·9, caud. 2·15, tars. 1, ung. post. 0·45, rostr. a fr. 0·5, a rictu 0·75, tota circa 6·25 poll. Angl. Fœmina vix minor.

Hab. in provincia Habessinica "Tigré" dicta. Circa pagum Senafé frequentissimam inveni.

Criithagra flavivertex, sp. nov.

C. supra olivacea, obsolete fusco maculata, fronte late aurea, pileo summo paullatim olivascente; superciliis flavis, postice productis; loris fuscis; colli lateribus olivaceis, immaculatis; uropygio flavo; remigibus cum tectricibus alarum rectricibusque fuscis flavo marginatis, remigum marginibus internis pallidis; gastræo sordide flavo, olivascenti-lavato, crasso albescente; rostro brunneo, pedibus fuscis. Fœmina vix dilutius colorata.

Long. al. 3·15, caud. 2·15, tars. 0·6, rostr. a fr. 0·35, tota circa 5·25 poll. Angl.

Hab. in provincia Habessinica "Tigré" dicta.

Lepus tigrensis, sp. nov.

L. persimilis *L. saxatili*, sed minor, cauda breviore, plantarum pilis ferrugineis, haud umbrinis.

Long. capitis 4, corporis circa 13, caudæ cum pilis ad extremitatem 4, sine pilis 3, auris 5, lat. ejusdem 2·8; long. cranii 3·5, lat. 1·6, alt. 2·2; long. tibiæ 5·3, tarsi 4·5, radii 4·5, carpi 2·3 poll. Angl.

Syn. *Lepus abyssinicus*, Lefèbvre, Voyage in Abyssinie, Atlas, pl. 5. fig. 1 (nec Hempr. et Ehrenb.).

More complete descriptions, where necessary, and figures will be given in my forthcoming account of natural-history observations made during the Abyssinian expedition.