

made. An operation found the colon in this region tightly bound by adhesions, and left them broken up; now there are no more psychical manifestations. The author believes that in psychic states as well as in neurasthenia a cure results not from direct treatment of the nervous system, but from medical or surgical treatment of one or other of the viscera.

W. A. BASTEDO.

TREATMENT OF INSANITY BY CONTINUOUS BATHS. P. Keraval (*Le Progrès Médical*, May 3, 1902).

The baths at Heidelberg used by Kraepelin and W. Alter are continued day and night for weeks and months, the patients eating from tables placed over the bath, and being allowed to smoke, read, crochet, etc. At night a rubber air-pillow is supported beneath the head. The water is kept at 34 deg. C. (93 deg. F.)

Such baths are specific in mania, agitation and the acute delirium of general paralysis. By suspension in water in a hammock the feeble and paralytic escape injuries and bed-sores. Almost without exception the patients accept the treatment, though in some cases it must be begun in short sances. The ordinary functions of urination and defecation make the most trouble, but not more than they do in a cell. The patient is quickly calmed and rapidly gains an appetite. It is not of much use in epilepsy, but for delirium tremens it is valuable, combined with caffeine in large doses. Alzheimer reports success with these baths at Frankfort, twelve bathers being constantly employed. Bieberbach has used them for a year at Heppenheim with very favorable results. In periodical insanity the subsequent attacks are especially mild. Insomnia is never persistent.

W. A. BASTEDO (New York).

INFANTILE CONVULSIONS. Prof. Déspine (*La Méd. Moderne*, April 9, 1902).

Infantile convulsions may be divided into two main groups, symptomatic convulsions, due to disease of the cerebral nervous system (meningo-encephalitis, tumors, hemorrhages, etc.), and idiopathic or essential. The latter group includes three main forms, external convulsions (eclampsia), internal convulsions (spasm of the glottis), and contractions of the extremities (tetany). The idiopathic forms are here studied as to etiology and symptomatology. Heredity plays an important part in causation. The influence of rickets is that of an auto-intoxication. Dentition is of no importance. In the newborn convulsions are usually due to organic lesion, and the prognosis is grave. Often asphyxia is a cause, and is due to congenital cyanosis, whooping-cough, bronchopneumonia. Many cases are attributed to poisoning by lead, alcohol, opium, santonin, etc., The most frequent cause is auto-intoxication (uremia, athyroidism, hypertrophy of the thymus, gastrointestinal disease). Reflex eclampsia may be due to irritation of the skin, ears or digestive tract. Clinically, eclampsia in infants is preceded by a stage of hyperexcitability of the nervous centers, manifested by hyperesthesia, exaggerated reflexes, electrical hyperexcitability. As to form, the convulsion is neither purely tonic nor clonic; exceptionally a purely tonic convulsion may occur. Commonly the spasm begins at the eyeball, extending to the face, neck, extremities and trunk; if localized, the face is usually the part involved. Bilateral symmetry is the rule; exceptionally the spasm is exaggerated on one side. Spasm of the glottis is characterized by a series of inspiratory or expiratory sounds; the paroxysm may end fatally by syncope, but typical cases are described which were saved by tracheotomy or by intubation. Spasm of the glottis is characteristically a disease of the first year of life;