

parts in case of falls. 7. The orbital foramen was implicated in 21.4 per cent. of the cases of orbital fossa fracture. 8. Inequality and immobility of pupils, or both, furnish the most frequent and unfavorable sign of fracture of the base. In the forty-four cases in which the pupils were recorded, they were normal in only thirteen. 9. Injury to the ciliospinal tract in its intracranial course is a more probable cause of the Hutchinson pupil and the other pupillary changes than injury to the third nerve or to the cortex, though no single lesion explains all cases. 10. The reflexes may be lessened or lost in fracture of the base, as in any case of violent jarring of the brain. On the other hand, they may be increased even to spasticity, probably through direct pressure on the pyramidal tract as by hemorrhage. It is probable that the initial result of the impact in all cases is a tendency towards lessening or loss of the reflexes. 11. Profuse and persistent bleeding from the ear does not suggest middle meningeal hemorrhage. No middle meningeal hemorrhage was found in the cases of profuse and persistent bleeding, and, conversely, hemorrhage from this artery occurred eight times without, and once with only slight, bleeding from the ear.

NOYES (New York.)

POLYNEURITIS. L. Harrison Mettler (Medicine, July, 1904).

Author states, as a result of the increasing number of observations revealing central as well as peripheral lesions in multiple neuritis, the opinion is fast gaining ground that in the toxic and infectious case, at least, the peripheral degeneration is dependent upon the central damage, organic or functional, of the nutritive cell bodies of the respective neurones. In other words, the peripheral changes are secondary in very many of the cases, at least, and are poliomyelitic in origin. Many conservative pathologists still believe, however, that there is a direct effect of a destructive character exerted upon the peripheral nerves by the toxin simultaneously with the effect of the same poison upon the central elements.

J. E. CLARK (New York.)

THE REFLEXES IN ALCOHOLISM (Deut. Med. Woch. No. 2, 1904).

Gudden has described a slow reaction of the pupil in acute alcoholism. Kutner has noted the condition of loss of muscular tone and tendon reflexes in the same condition. In cases under his observation transitory disturbances of consciousness and fixed ideas were noted. Typical epileptic attacks and hemiparesis were seen. In all these cases the reaction appeared after a relatively small amount of alcohol had been taken, and sometimes when they had been abstemious for some time. Sometimes the knee-jerks were increased, indicating central irritation. In all these cases of intoxication a marked increase of the passive mobility of the limbs, hypotonia and feebleness of tendon reflexes were noted. Skin, plantar and abdominal reflexes were variable. The corneal and conjunctival reflex were constant.

NOYES (New York.)

OVERWORK AND MYASTHENIA GRAVE. PEL (Berlin klin. Woch. No. 25).

The author observes that myasthenia, with or without bulbar paralysis, is found to be related to the morbid phenomena of hysteria, with the cardinal symptoms of grave neurasthenia, or with the cerebral or spinal symptoms of multiple sclerosis, and with syphilitis of the nervous system. He publishes a case where neurasthenic phenomena predominated. Concerning atrophy of the tongue, he states that muscular atrophy does not of necessity belong to grave myasthenia, and the lack of reaction of degeneration is characteristic. Atrophy of the muscles of the neck may occur. Oppenheim suggests the term of myasthenia pseudo-paralytica. Overwork of certain muscles is the chief cause of the disease.

NOYES (New York)