

presented no traces of inflammation; there was, at a few points, a hyaline exudation into the walls of the capillaries. The examination of the motor regions of the brain, which were about as hard as an olive, demonstrated that the neo-formation had begun in the neuroglia fibres.

The fifth brain presented no appreciable lesion to the naked eye. Under the microscope, however, in the region of the paracentral lobule, the same neuroglia changes were observed that were found in the other brains.

The author reaches the following conclusions:

1. Certain lesions, described under the name of cerebral sclerosis, are due, in many instances, to proliferation of the connective-tissue, particularly of the fibres of the neuroglia. For these cases the name of *neuroglia sclerosis* or *glioma* was proposed.

2. The induration of some portions of the brain, particularly of the cornua Ammonis and the olives, has been known for a long time in connection with epilepsy. This induration is the external evidence of the internal proliferation of the neuroglia. *When there are no macroscopical lesions, the same neuro-pathological fundamental process can nevertheless be recognized.*

G. M. H.

THERAPEUTICS OF THE NERVOUS SYSTEM.

HYSTERICAL CHOREA: CRISES CHARACTERIZED BY HYPOCHONDRIA, FOLLOWED BY AN IMPERIOUS DESIRE TO WALK, TO CRY, AND TO SING; CURED BY COPPER USED EXTERNALLY AND INTERNALLY.

Dr. Moricourt, at the Société Médicale (Journ. des Conn. Méd., April 18, 1889), reports the following case: The patient, a woman fifty years of age, had had these nervous crises for five years, ever since the death of her child. She had been subject for a long time to attacks of migraine, accompanied by nausea, which had diminished in frequency with the advent of the nervous crises. One of her sisters had genuine hysterical attacks. The crises usually came on between two and five o'clock in the morning. They

generally began with melancholy thoughts and with a desire to cry, frequently with headache, anorexia, nausea, and diarrhœa. This was soon followed by trembling in all the limbs, then by an imperious desire to walk, to cry, to sing, and to embrace people. These crises lasted about five hours, and occurred from three times in a month to several times in a week. Anæsthesia and analgesia were present on the outer surface of both forearms. An armlet, composed of four disks of steel, was applied to each forearm. At the end of ten minutes the patient experienced slight warmth in the right arm, which soon passed to the left arm, oscillated from one arm to the other, and then disappeared. The anæsthesia and analgesia were diminished, and the muscular strength also to a slight extent. The next day two armlets of copper were applied in the same manner. The patient experienced sensations of pain and heat much more acutely than with the steel; analgesia was diminished, while anæsthesia was increased. Zinc, tin, gold, silver, and aluminium were successively tried without result.

Dr. Moricourt therefore prescribed *eau d'Orezza* and external applications of copper. Under this treatment the intervals between the crises became longer, the appetite improved, and the menstruation, which had been absent for three months, returned. Shortly afterward the patient, having neglected the treatment for two days, was seized with another crisis, but much less severe than formerly. The water was then discontinued, and, in its place, pilules of sulphate of copper were given. Each pill contained one milligramme of sulphate of copper. The dose was two pills four times a day.

This treatment was continued from the 28th of January till the 23d of March. In this time the sensibility became normal on both sides, menstruation returned, and the crises disappeared.

G. M. H.

TREPHINING FOR CEREBRAL ABSCESS.

M. Terrillon, at a meeting of the Société de Chirurgie, July 3, 1889, reported the case of a child, thirteen years of age, who, following an attack of typhoid fever, presented a

generally began with melancholy thoughts and with a desire to cry, frequently with headache, anorexia, nausea, and diarrhœa. This was soon followed by trembling in all the limbs, then by an imperious desire to walk, to cry, to sing, and to embrace people. These crises lasted about five hours, and occurred from three times in a month to several times in a week. Anæsthesia and analgesia were present on the outer surface of both forearms. An armlet, composed of four disks of steel, was applied to each forearm. At the end of ten minutes the patient experienced slight warmth in the right arm, which soon passed to the left arm, oscillated from one arm to the other, and then disappeared. The anæsthesia and analgesia were diminished, and the muscular strength also to a slight extent. The next day two armlets of copper were applied in the same manner. The patient experienced sensations of pain and heat much more acutely than with the steel; analgesia was diminished, while anæsthesia was increased. Zinc, tin, gold, silver, and aluminium were successively tried without result.

Dr. Moricourt therefore prescribed *eau d'Orezza* and external applications of copper. Under this treatment the intervals between the crises became longer, the appetite improved, and the menstruation, which had been absent for three months, returned. Shortly afterward the patient, having neglected the treatment for two days, was seized with another crisis, but much less severe than formerly. The water was then discontinued, and, in its place, pilules of sulphate of copper were given. Each pill contained one milligramme of sulphate of copper. The dose was two pills four times a day.

This treatment was continued from the 28th of January till the 23d of March. In this time the sensibility became normal on both sides, menstruation returned, and the crises disappeared.

G. M. H.

TREPHINING FOR CEREBRAL ABSCESS.

M. Terrillon, at a meeting of the Société de Chirurgie, July 3, 1889, reported the case of a child, thirteen years of age, who, following an attack of typhoid fever, presented a

phlegmonous swelling over the left temporal region which was followed by aphasia and by paralysis of the right side of the face and the right arm. An incision into this subperiosteal abscess in the temporal fossa was not followed by any amelioration of the cerebral symptoms. The next day trephining was performed, and a button of bone was removed from over the area of the centre for the upper extremity, and a lesion was sought for which M. Terrillon believed would be found between the bone and the dura; but nothing was discovered.

Suspecting a cerebral abscess, three exploratory punctures were made, instead of an incision, as the operator did not believe that the rules laid down for cranio-cerebral topography in the adult applied equally as well to the child's brain. On the third puncture an abscess was found, which was freely opened. Following the evacuation of the contents of the abscess, the paralysis almost completely disappeared. Three days after the operation, meningo-encephalitis developed, and the patient died.

G. M. H.

OBSERVATIONS AND THEORIES RELATING TO HYPNOTISM.

In the *Gazette Degli Ospitale* are reported several interesting cases in which good results have been obtained by means of hypnotism.

The following case of nervous vomiting is reported by Dr. A. Colombi, in the *Journal* of March 31, 1889. The patient was a girl of twelve, who for a month had been affected with nervous vomiting occurring suddenly after each meal. She was irritable and showed a great lack of strength. She was hypnotized twice by means of fixing her eyes on a clinical thermometer for a few minutes. The vomiting and general weakness disappeared after the second treatment, and the irritability likewise, the child becoming happy and contented.

Dr. G. Ficano, in the same journal, August 11, 1889, publishes a case of hysterical cough cured by what he calls "moral" influence. The patient, a young woman, had suffered with a severe cough for several months. Her par-