

rare disease, though somewhat more common in young children than in adults. He regards it as similar in etiology to the other spasmodic disorders, and finds it, in children, quite amenable to treatment.

C. A. ALLEN.

TETANY FOLLOWING SCARLATINA.

Bradford McConnell (Canada Med. Record, Sept., 1896) reports a typical case of tetany following scarlet fever. The inception of the trouble was noticed fourteen days after the beginning of the fever and during desquamation. The specific action of the infectious disease in causing the spasmodic affection is more than doubtful, as the patient, aged 5½ years, had had a similar attack at 18 months. There was a vicious heredity, and the child himself had had numerous convulsions.

PATRICK (Chicago).

EIN FALL VON TETANY MIT EIGENTÜMLICHEM SECTIONSBEFUND. (A Case of Tetany with Peculiar Findings.) Deutsche Zeitschrift für Nervenheilkunde, Band ix., Heft 3 und 4. By H. KÖSTER.

The diagnosis of tetany in this case was founded on tonic, intermitting frequently painful cramps of individual muscles or cramps of muscles, on the presence of Trousseau's and Erb's signs, on intermitting spasm of the ocular muscles, retention of urine, dysphagia, tinnitus aurium, absence of mental symptoms, vertigo and the development of the disease in the first months of the year.

The simultaneous appearance of the cramps in all four extremities, affecting the extensors as well as the flexors, the flexion of the leg, the intensity of the spasm in the neuchal muscles, the ability of the patient by an effort of the will to control the cramps, the position of the hand as a fist, the clonic convulsions, at times, in the muscles of the extremities and eyes, were peculiarities noticed in this case. A slight rigidity of the muscles prevented the knee jerk. The autopsy revealed contracted kidneys, circumscribed hemorrhages on the anterior part of the cauda equina and about the roots of the third to the fifth cervical nerves, as well as hemorrhages and hyperæmia in the sheath of the right sciatic nerve.

Köster is unable to explain the symptoms from the post-mortem findings.

SPILLER.

PATHOLOGICAL ANATOMY.

ASCENDING DEGENERATION IN NERVES AND THE CONSEQUENT CHANGES IN NERVE CELLS.

Fleming has made an unusually extensive and careful study of the changes that occur in the central stump, and in the cells of the spinal cord after division or ligation of the nerve trunk.

In the nerve he found: "First, that in process of time a slow atrophy of 'motor' fibres occurs. Secondly, that certain 'sensory' fibres degenerate centrally, possibly because severed from their peripheral trophic centres. Thirdly, that the minute fibres found in a normal nerve undergo very marked change. Fourthly, that distinctive connective tissue increase occurs."

The changes found in the cells are summarized as follows:

"1. The cells of the ganglia on the posterior nerve roots undergo definite changes as the result of nerve section or ligature, and do so at a much earlier period than the multipolar cells in the cord—beginning probably as early as the fourth day and certainly by the seventh day.

"2. That one of the very first changes observed in the cells of ganglia and anterior cornu is a diminution in the size of the nucleus—in proportion to the size of the cell and that, sometimes, but not in all cases—nucleoli also becomes smaller and very frequently the nuclei take up an eccentric position, sometimes even bulging the cell wall.