



Annals and Magazine of Natural History: Series 6

Publication details, including instructions
for authors and subscription information:
<http://www.tandfonline.com/loi/tnah12>

XXXVIII.—Description of a new snake from Borneo

G.A. Boulenger F.R.S.

Published online: 06 Oct 2009.

To cite this article: G.A. Boulenger F.R.S. (1895) XXXVIII.—Description of a new snake from Borneo , Annals and Magazine of Natural History: Series 6, 15:88, 329-329, DOI: [10.1080/00222939508677890](https://doi.org/10.1080/00222939508677890)

To link to this article: <http://dx.doi.org/10.1080/00222939508677890>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs,

expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

skull of this *Geotrypetes*, I come to the conclusion that the genus is valid, for the squamosals are not in contact with the parietals, as in *Hypogeophis rostratus*. It may be thus defined :—

GEOTRYPETES.

(Peters, Sitz. Ges. naturf. Fr. Berl. 1880, p. 55.)

Squamosals separated from parietals. Teeth large, numerous in inner row of mandible. Eyes visible. Tentacle flap-shaped; tentacular groove horseshoe-shaped, situated below and behind the nostril. Cycloid scales imbedded in the skin.

A single species, from West Africa :—

Geotrypetes Petersii.

Hypogeophis seraphini, Peters, Mon. Berl. Ac. 1879, p. 937; Boulenger, Cat. Batr. Caud. Ap. p. 97 (1882).

XXXVIII.—*Description of a new Snake from Borneo.*

By G. A. BOULENGER, F.R.S.

Calamaria Brookii.

Rostral as deep as broad, the portion visible from above half as long as its distance from the frontal; frontal once and one fourth as long as broad, much shorter than the parietals, thrice as broad as the supraocular; a præocular and a post-ocular; diameter of eye equal to its distance from the mouth; five upper labials, third and fourth entering the eye; first pair of lower labials forming a suture behind the symphysial; two pairs of chin-shields, in contact with each other. Scales in 13 rows. Ventrals 147; anal entire; subcaudals 23. Tail obtusely pointed. Yellowish brown above, with five black stripes, the median the broadest and occupying one scale and two halves; head marbled with black; a black nuchal collar; two similar black bars on the tail, one at the base, the other near the end; outer row of scales, ventrals, and subcaudals yellowish white, the upper third of the outer scale black, otherwise unspotted.

Total length 220 millim.; tail 23.

A single male specimen, from Matang. Presented to the British Museum by H.H. Rajah Brooke.