

light on the question whether a microörganism is the sole cause of multiple sclerosis or if there are other infectious factors, as would seem to be the case from clinical indications. Malaria, for example, may run a course either as multiple or pseudosclerosis, and H. Duerk has even proved the presence of granuloma in the central nervous system in malaria, which he is inclined to regard as multiple sclerotic foci in an initial stage. The author thinks that the picture which Duerk describes as multiple sclerosis bears more resemblance to acute encephalomyelitis, though this latter disease is not a form wholly contradictory of multiple sclerosis of the type usually encountered. The author inoculated rabbits with the spinal fluid of two fresh cases of multiple sclerosis. The fluid was sterile, excluding lues. Nine days after one animal manifested slight paralysis in the hind leg together with other paralytic signs. The other animals remained healthy. In the next ten days the forelegs of the affected rabbit were paralyzed; the animal was helpless and refused food. In nineteen days after the injection it died without spasms. On section, the place of injection, the spinal cord, the skin, the brain and the internal organs were found to reveal not the slightest pathological changes, either generally or upon transverse section. A rabbit inoculated with fresh sterile fluid from this animal showed signs of severe paralysis of the hind leg. The animal was killed after fourteen days but there were no pathological findings. The author agrees with Siemerling and Raecke, whose patient attempts to transmit multiple sclerosis were unsuccessful, that failure to find the exciting cause should not discourage, when it is remembered how many years of fruitless search preceded the discovery of the cause of lues.

4. MIDBRAIN: CEREBELLUM.

Maas, Otto. DYSTONIA MUSCULORUM DEFORMANS. [Neurol. Centralbl., March 16, 1918, No. 6, Vol. 37.]

In 1908 Schwalbe described three cases in one family where during childhood muscle contractions of varying intensity made their appearance followed in later years in one case with total inability to walk, and in the two others by extreme disturbances of gait. There were no symptoms of the pyramidal tract and no disturbance of sensibility. Ziehen described a similar case which he called "tonic torsion neurosis." Oppenheim, who had opportunity to observe a large number of cases of this disease extended the knowledge of the clinical picture, showing that tonic cramps as well as clonic are present and that in isolated muscle groups there is hypotonia. He was of the opinion, in contradiction to other writers, that the symptoms were due to a disease of the central nervous system, and proposed the name *dysbasia lordica progressiva* or *dystonia musculorum deformans*. The author gives a case in which a disturbance in gait set in following typhus. A tendon operation was performed which only increased the difficulty. Soon a trembling of the

head and arms began and a curvature of the body. Some years afterward a second operation on the tendon of the leg was performed. It is to the course of the disease after the second operation to which the author would call attention. The patient claims that it resulted in marked improvement and her assertion was confirmed by two physicians who had observed her during the earlier period of the disease. Other authors who have described similar cases claim that this disease is progressive. Ziehen asserts that no cases of improvement have been observed, that the disease progresses until it reaches a certain chronic standstill. Oppenheim says that the spasms may improve in certain muscle regions, especially in the arm, but that the general course is progressive. In his case the author found the whole condition improved and during a period of years there was no regression. Again, Flatau and Sterling state that in this new sort of spasm the face is not affected, as is nearly always the case in athetosis; a second difference in the author's case from the former experiences is that there was a tremor of the face. Schwalbe and Ziehen mention psychic symptoms, while Oppenheim says that he does not believe they belong to the picture of this disease; in the author's case they were present. Further, there was close resemblance of this disease to Wilson's disease and to pseudosclerosis. He thinks that none of these diseases are hereditary but arise gradually, from unknown causes in youth and the abnormal muscle tone is their principal feature, but, in his case of dystonia, the trembling did not seem to be as great as it is in pseudosclerosis. Just how close the relation is cannot be definitely decided, because there have not been any anatomical examinations of dystonia. The two points emphasized are that his observations lead him to believe, contrary to his predecessors, that psychic symptoms belong to the clinical picture of the disease, and that, in the course of the disease, considerable improvements may take place.

Paulian, D. E. STUNTED PYRAMIDAL-CEREBELLAR SYSTEM. [Rev. Neur., Nov., 1919.]

Paulian had two cases, a woman 31 and her brother 26. They both had arrested development of the pyramidal system and afferent cerebellar tracts. They were both about 20 when first ill. An uncle had a history of spinal cord disease, and the mother had unmistakable signs of syphilis. Paulian labels them familial pyramido-cerebellar dysgenesia, emphasizing also the inherited taint.

Brouwer, B. CHRONIC DIFFUSE DISEASE OF THE CEREBELLUM. [Neurol. Centralbl., Nov. 1, 1919, Vol. 38, p. 674.]

The author describes two cases, the first being important from both clinical and pathological points of view; it shows that the same symptom complex may occur in "*atrophie lamelleuse des cellules de Purkinje*" as