

# THE LOCATION WITHIN THE SPINAL CORD OF THE FIBERS FOR TEMPERATURE AND PAIN SENSATIONS.<sup>1</sup>

BY WILLIAM G. SPILLER, M.D.,  
OF PHILADELPHIA,

PROFESSOR OF NEUROPATHOLOGY AND ASSOCIATE PROFESSOR OF NEUROLOGY IN  
THE UNIVERSITY OF PENNSYLVANIA.

Many believe that the tracts of Gowers may contain the fibers for the sensations of pain and temperature, but evidence supporting this opinion has been sadly deficient. There has been little to show that the fibers for these sensations are confined to Gowers' tracts and are not present elsewhere in the antero-lateral columns. J. Piltz<sup>2</sup> is the most recent author to support the view referred to above.

A case has been in my wards at the Philadelphia General Hospital within the last few months which seems to me to afford the best evidence that has as yet been offered for the location of the fibers for temperature and pain within the tracts of Gowers.

A man, twenty-three years of age, began to have pain in the lumbar region and abdomen four months before he came under my observation, Sept. 1, 1904. Numbness was soon felt in the feet, and gradually involved the lower limbs below the knees. When standing he had extreme genu valgum, which had existed about five years. He had scars on the abdomen which he said were caused by painless burns in childhood. Resistance to passive movement was slightly diminished in the lower limbs. Sensation for touch was normal or nearly normal in the lower limbs, but sensation for pain was abolished or almost abolished in these parts. Sensation for temperature was almost lost in the lower limbs below the knees, and much impaired in the thighs and lower part of the trunk. Walking was difficult because of the deformity of the knees.

---

<sup>1</sup> Part of a paper read by invitation before the Section on Neurology and Psychiatry of the Medical and Chirurgical Faculty of Maryland, March 8, 1905. The paper will be published in full in the University of Pennsylvania Medical Bulletin.

<sup>2</sup>Piltz. *Neurologisches Centralblatt*, March 16, 1905.

He had a slight kyphosis at about the tenth thoracic vertebra. The upper limbs and face were not affected.

After a fall backwards down a flight of steps he became completely paralyzed in the lower limbs. Sensation for touch was then lost in the right lower limb for a time, but was preserved in the left lower limb. Sensations for temperature and pain were lost in the right leg and foot and in the left foot, but were preserved elsewhere, although it is not stated by Dr. McConnell, who made this note, that they were normal else-




Fig. 1. Small tubercle in the right lateral column extending forward into Gowers' tract.

where. The patellar reflexes were much exaggerated and Babinski's sign was present.

Still later sensation for touch was found preserved everywhere in the lower limbs. Sensation for pain was irregularly present in the lower limbs, the response to pain stimulation being very uncertain, and sensation for temperature also was much impaired in the lower limbs.

Tuberculous meningitis and caries of the vertebræ were found, and a small tubercle was present in the right lateral column at the extreme lower end of the thoracic cord, in-

volving the area of Gowers' tract. About one-half to one inch higher another small tubercle was found involving the left tract of Gowers. Although there was some myelitis, it was very evident from the symptoms that it had chiefly devel-




Fig. 2. Small tubercle in the left antero-lateral column implicating Gowers' tract.

oped after the fall down the flight of steps, as at the time the patient first came under my observation weakness in the lower limbs was slight. He lived four months after the accident.