

This article was downloaded by: [University of Glasgow]
On: 20 December 2014, At: 06:52
Publisher: Taylor & Francis
Informa Ltd Registered in England and Wales Registered Number:
1072954 Registered office: Mortimer House, 37-41 Mortimer
Street, London W1T 3JH, UK



Annals and Magazine of Natural History: Series 6

Publication details, including instructions for authors and subscription information:
<http://www.tandfonline.com/loi/tnah12>

LIVIII.—Characters of a new genus and species of Cicindelidæ

Charles O. Waterhouse
Published online: 09 Oct 2009.

To cite this article: Charles O. Waterhouse (1889) LIVIII.—Characters of a new genus and species of Cicindelidæ, *Annals and Magazine of Natural History: Series 6*, 3:18, 486-486, DOI: [10.1080/00222938909460373](https://doi.org/10.1080/00222938909460373)

To link to this article: <http://dx.doi.org/10.1080/00222938909460373>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever

caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

LVIII.—*Characters of a new Genus and Species of Cicindelidæ.* By CHARLES O. WATERHOUSE.

AMONG some insects from South Brazil which recently came under my notice were two male examples of a Cicindelid allied to *Oxycheila* and *Raminagrobis*, but which differ from both those genera in the form of the labrum and mandibles; these latter are quite unlike those of any Cicindelid with which I am acquainted and remind one of some species of Lucanidæ.

COPHOGNATHUS.

General characters of *Oxycheila*. Central tooth of the mentum equilaterally triangular, very acute at its apex. Labial palpi very slender, the basal joint extending a little beyond the central tooth, the second joint only a little shorter than the first, the fourth joint a little widened at its apex, rather shorter than the third. Labrum scarcely longer than broad, triangular, denticulate near the apex. Maxillary palpi with the fourth joint a little shorter than the third. Mandibles strong and thick, flattened, curved, with two obtuse approximate teeth near the apex. Thorax, as compared with that of *Oxycheila tristis*, broad, constricted at its base. The anterior tarsi in the male with the three basal joints dilated. Wings present.

Cophognathus opacipennis.

Niger; capite thoraceque nitidis; antennis articulis 5^o–11^m testaceis; elytris opacis, basi punctatis, ad apicem emarginatis. Long. 12 lin.

Hab. São Paulo, South Brazil.

This interesting species has somewhat the appearance of *Oxycheila tristis*, but the small labrum and blunt mandibles are quite different. The eyes are relatively smaller. The thorax is shorter and broader, shining, strongly constricted at the base, with a deep lunate impression on the disk a little in front of the middle, and with the basal transverse impression strongly marked. The elytra are a little shorter, dull, with some not very strong punctures at the base, a few of which are asperate; the apex narrowly truncate (or slightly emarginate), the outer angle of the truncature slightly dentiform. Labrum, mandibles, and tarsi obscurely pitchy.