

This article was downloaded by: [University of Cambridge]
On: 08 October 2014, At: 06:08
Publisher: Taylor & Francis
Informa Ltd Registered in England and Wales Registered Number:
1072954 Registered office: Mortimer House, 37-41 Mortimer
Street, London W1T 3JH, UK



Annals and Magazine of Natural History: Series 6

Publication details, including instructions
for authors and subscription information:
<http://www.tandfonline.com/loi/tnah12>

Aspidiotus bicarinatus a Lepidopterous Larva

E.E. Green

Published online: 12 Oct 2009.

To cite this article: E.E. Green (1890) *Aspidiotus bicarinatus* a
Lepidopterous Larva , Annals and Magazine of Natural History: Series 6,
6:36, 503-503, DOI: [10.1080/00222939008694078](https://doi.org/10.1080/00222939008694078)

To link to this article: <http://dx.doi.org/10.1080/00222939008694078>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of
all the information (the "Content") contained in the publications
on our platform. However, Taylor & Francis, our agents, and our
licensors make no representations or warranties whatsoever as
to the accuracy, completeness, or suitability for any purpose of
the Content. Any opinions and views expressed in this publication
are the opinions and views of the authors, and are not the views
of or endorsed by Taylor & Francis. The accuracy of the Content
should not be relied upon and should be independently verified
with primary sources of information. Taylor and Francis shall not be
liable for any losses, actions, claims, proceedings, demands, costs,
expenses, damages, and other liabilities whatsoever or howsoever

caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

north in the Atlantic, namely Les Passages, near San Sebastian; Paul Fischer, however, has not met with it on the closely adjoining south-west coast of France.

It may be added that Lamarek, in his original description of the species, speaks of its being confounded with *A. glacialis*, and points out the differences between the species. A. M. NORMAN.

Burnmoor,
Nov. 3, 1890.

Since writing the above, on turning to Dujardin and Hupé, I find that they, as I have done, refer Linck's and Pennant's form to *A. glacialis*.

Aspidiotus bicarinatus a *Lepidopterous Larva*.

To the Editors of the Annals and Magazine of Natural History.

10 Observatory Gardens,
Campden Hill, Kensington, W.,
November 6, 1890.

GENTLEMEN,—On looking through the collection of Coccidæ in the Students' Insect-Room at the Natural-History branch of the British Museum my attention was drawn to two insects labelled *Aspidiotus bicarinatus*. The tickets attached to these insects showed them to be the veritable types described by Walker in the supplement of the British Museum 'Catalogue of Homoptera,' p. 306, as *Aspidiotus bicarinatus*. But upon examination I found them to be the dried larvæ of a Limacodid moth closely resembling those of the Indian species *Narosa conspersa*. The caterpillar of this moth has a coriaceous integument with two well-developed dorsal ridges. As the specimens in question were received from North China, they are probably the larvæ of some allied species.

It is difficult to understand how the mistake could have originated. In its dried state the caterpillar has certainly some superficial resemblance to a Coccid; but its head and mouth-parts at once proclaim its true character. E. E. GREEN.

Note on Irrisor Jacksoni, sp. n. By R. BOWDLER SHARPE.

In a letter just received from Mr. F. J. Jackson he has given some very interesting notes on birds, some of which he believes to be new to science. The *Irrisor* is certainly undescribed, and I herewith name it after the explorer.

Irrisor Jacksoni, sp. n.

Ad. Similis *I. Bollei*, sed minor rostro brevior, tectricibus alarum intimis chalybeis nec cuprescentibus distinguendus.

Long. tot. 13·0, culmin. 1·35, alæ 4·85, caudæ 7·4, tarsi 0·8.

Hab. Kikuyu Country, Eastern Africa.