


This article was downloaded by: [University of Glasgow]
On: 30 December 2014, At: 12:47
Publisher: Taylor & Francis
Informa Ltd Registered in England and Wales Registered Number:
1072954 Registered office: Mortimer House, 37-41 Mortimer Street,
London W1T 3JH, UK



Annals and Magazine of Natural History: Series 2

Publication details, including instructions
for authors and subscription information:
<http://www.tandfonline.com/loi/tnah08>

Note on Helix Cantiana, Mont

Wm. Lonsdale Esq. F.G.S.
Published online: 26 Oct 2009.

To cite this article: Wm. Lonsdale Esq. F.G.S. (1856) Note on Helix Cantiana, Mont , Annals and Magazine of Natural History: Series 2, 18:104, 187-187, DOI: [10.1080/00222935608697614](https://doi.org/10.1080/00222935608697614)

To link to this article: <http://dx.doi.org/10.1080/00222935608697614>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly

or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

fore propose to change it into *Heterocnemis*. A second species of the genus seems to be the bird figured in Buffon's Pl. Enl. 73, fig. 2, under the name of 'Le Bambla de Cayenne.'—(*Turdus bambla*, Bodd.—*Heterocnemis bambla*, mihi.)—The present bird is very closely allied to the latter. In the upper plumage it is very similar, being only of a more cinnameous tinge; but it may be distinguished by the want of the white markings on the wings, and the throat, breast and upper belly being white, each feather narrowly margined with brown. In *H. bambla* these parts are ash-brown, with obsolete transverse markings. Lesson's *Myrmothera troglodytes* (Desc. d. Mamm. et Ois. p. 301, no. 118) seems the same as *H. bambla*. This form is indeed very closely connected with some of the Wrens, and hardly to be separated from certain birds that are usually placed in the genus *Scytalopus*.

TODIROSTRUM GRACILIPES, Sclater.

T. supra olivaceum; alis caudaque nigricantibus, olivaceo anguste limbatis; pileo fuscescente; loris mentoque albidis; subtus flavum, lateribus olivascens; gutture et pectore longitudinaliter nigro striatis; tectricibus subalaribus sulphureis; rostro nigro; pedibus carneis; tarsis gracillimis.

Long. tota 3·8; alæ 2·0; caudæ 1·7.

This apparently new species is most nearly allied to *T. maculatum* (Desm.) and *T. striaticolle*, Lafr. (in both of which the neck is also striated), but has the whole throat yellow. The bill agrees in form with that of the former bird, but is rather shorter and narrower. The type specimen is in the British Museum.—*Proc. Zool. Soc.* July 24, 1855.

ON THE BRITISH DIASTYLIDÆ.

To the Editors of the Annals of Natural History.

8 Mulgrave Place, Plymouth, July 10, 1856.

GENTLEMEN,—It has recently been pointed out to me by Mr. Alder that I have unfortunately made use of two generic names in the paper on "the British *Diastylidæ*," lately published in the 'Annals,' that have been previously employed to designate other genera of animals. It is therefore desirable that the names *Halía* and *Venilia* should be changed, and I propose instead to use *Iphinoë* and *Cyrianassa* respectively. The names as applied will therefore be *Iphinoë trispinosa* and *Cyrianassa gracilis*.

Believe me, yours obediently,

C. SPENCE BATE.

Note on Helix Cantiana, Mont. By WM. LONSDALE, Esq., F.G.S.

"*Helix Cantiana* is abundant around Keynsham, Somerset. It was first found in the spring of 1825, alive and near the entrance to Dr. Fox's establishment, between Keynsham and Brislington; and more recently in lanes close to the town."