

Several subcommittees were appointed.

Subcommittee on Infant Feeding: *Chairman*, Dr. Harvey W. Wiley, Chief of Bureau of Chemistry, U. S. Department of Agriculture; Gen. Geo. M. Sternberg, Dr. Stitt, U. S. Navy; Dr. J. W. Kerr, Asst. Surgeon General, Bureau of Public Health and Marine-Hospital Service; Drs. D. P. Hickling, Wm. M. Sprigg, and W. C. Woodward.

Subcommittee on the Sanitary Condition of the Milk Supply: *Chairman*: Dr. Geo. M. Kober, Professor of Hygiene, School of Medicine, Georgetown University. His associates were Drs. A. D. Melvin, Chief of Bureau of Animal Industry, U. S. Dept. of Agriculture; M. J. Rosenau, Director of Hygienic Laboratory, U. S. P. and M.-H. S.; John R. Mohler, Chief Pathologist, Bureau of Animal Industry, Department of Agriculture; Chas. T. Mason, Asst. Surgeon General, U. S. Army; G. Lloyd Magruder, Emeritus Professor of Materia Medica and Therapeutics, School of Hygiene, Georgetown University, together with Messrs. E. H. Webster, Chief of Dairy Division, Bureau of Animal Industry, Dept. of Agriculture, Col. Robert G. Smith, milk producer, and E. Berliner.

The matter assigned to these committees was divided into several subheads and was so carefully and exhaustively considered by men well versed on the subject that copious excerpts from the reports which were submitted to the full commission on April 17 and May 1 and unanimously adopted, are presented in abstract with the feeling that much of value will be given:

Extracts from the recommendations unanimously adopted by the committee to consider milk in relation to the food of infants:

1. The milk which is supplied to infants under the age of 3 years in the District of Columbia should be certified by the health officer, and should contain not more than 10,000 bacteria per cubic centimeter, should be not more than twelve hours old, and should be delivered in artificially cooled packages.
2. The health officer of the district should be authorized to advertise for dairies which will be willing to so modify their barns, stables, etc., if necessary, as to secure a license from him for the production of certified milk under the most modern improved sanitary conditions.
3. Each cow furnishing the milk in the dairy should be tested, under the supervision of the health officer, for tuberculosis or other contagious or infectious diseases, and that any animal so suffering be excluded from the herd.
4. A daily sample, or samples, drawn from the supply of each certified milk furnished to the city should be secured for the purpose of making the bacterial count and determining the chemical composition, etc.
5. Parents and guardians should be urged to use only certified milk, at least for infants' food in the District of Columbia for all infants under the age of 3 years.
6. If private dairies can not be induced to furnish certified milk, the commissioners of the District of Columbia, as a health measure affecting in the highest degree the welfare of the District of Columbia, should be asked, if it can legally be done, to establish a municipal dairy for the purpose of furnishing certified milk for the use of infants under 3 years of age in the District of Columbia.
7. For the purpose of securing modified milk for infants in ill health, according to the prescription of physicians the Milk Commission of the District Medical Society should be requested to secure the establishment of a laboratory under its supervision in which only certified milk shall be used, and in which, by the use of a separator, or otherwise, milk of a definite chemical composition may be prepared from certified milk, in harmony with the physicians' prescriptions relating thereto.
10. The term "certified milk" as used herein is to be applied to milk secured at dairies subject to a periodic inspection, and the products of which are subjected to constant analysis. The cows providing the milk are to be properly fed, free of tuberculosis or other contagious diseases, and housed in clean stables, properly ventilated, and they are to be supplied with

wholesome water and food, and kept clean. The milk is to be drawn under precautions to avoid infection, is to be immediately strained and cooled, and packed in sterilized bottles which are to be kept at a temperature of about 40 degrees F. until delivered to the consumer.

#### VARIOUS DISEASES AND CONDITIONS OF CATTLE THAT MAY AFFECT THE QUALITY OF THE MILK SUPPLY.

BY JOHN R. MOHLER.

The importance of obtaining a hygienic and wholesome milk supply for the residents of this city is recognized by all intelligent people and should require no argument. Public health demands the purity of all milk and milk products. Next to bread, milk is more extensively used as an article of diet than any other foodstuff. It forms a portion of the food of almost every person practically every day of the year. Moreover, unlike many other articles of diet, milk is consumed in most cases in an uncooked state, making it a very dangerous food should it perchance contain any deleterious organisms. The reasons for securing a supply of pure and wholesome milk are so numerous and so important that the consumer should become acquainted with some of the more essential of them in order that he may render assistance in bringing about a satisfactory improvement. Not only is milk a very suitable medium for almost every description of germ life which may gain access to it in its journey from the cow to the consumer, but it may also become contaminated while still in the udder through infectious or poisonous material present in the cow herself. Consideration in this report will be given only to the latter aspect of the question of a wholesome milk supply. In this connection it will be necessary to keep in mind the requirements of an awakened public for a clean and wholesome milk as well as the effect of any unreasonable or irrational demand on the producer which may cause him heavy losses, or even to discontinue his business.

It will also be apparent from the report of the Committee on Sanitation that in order to produce milk in compliance with the requirements hereinafter to be described, certain precautions must be taken which will necessarily entail additional expense on the producer of this higher grade of milk. The customer must therefore expect to pay his portion of any legitimate advance in the cost of production, and such increase in the price of milk due to its improved quality should be considered as money well expended. Furthermore, we can not have good milk of safe quality without a realization on the part of the farmer, transportation agent, dairyman and housewife of the danger in utilizing old, warm or dirty milk. Education is an important factor, therefore, in the improvement of the milk supply which can not be accomplished through laws and regulations alone. In view of these facts it is recommended that the subject be taught in the schools, that popular articles be frequently prepared for the press, that lectures and demonstrations be given in towns and townships, that pamphlets in plain language be prepared by the Health Officers for general distribution, and especially that rules and suggestions, with reasons therefor, be placed in the homes of dairy-men and dairy attendants.

#### MILK FROM UNHEALTHY COWS AS A FACTOR IN THE SPREAD OF DISEASE.

*Tuberculosis.*—Probably the most important disease of cows from the standpoint of public health is tuberculosis, and it is also the most prevalent. Notwithstanding the announcement of Koch that, in his opinion, tuberculosis in man and cattle is a different disease, it is now the generally accepted opinion among scientists that people, especially children, may become infected with tuberculosis from cattle. It is not known to what extent such infection occurs, nor is it possible to obtain any definite percentage by the method formerly adopted of looking for the primary lesions in the intestinal canal, although much statistical evidence is recorded, showing that even by these figures primary intestinal tuberculosis of chil-

dren has been observed in as high as 45.5 per cent. of the tuberculous cases examined. (Heller.)

Evidence which must be considered conclusive has been obtained by the Bureau of Animal Industry as well as by Ravenel and a number of French investigators, showing that tuberculous infection may take place through the intestinal tract without leaving any lesion in the abdominal cavity, the first alteration being found in the lungs or thoracic glands. Therefore the presence of pulmonary tuberculosis in infants without intestinal lesions is no indication that the disease was not transmitted by the food, and the statistics above referred to are thus shown to be below the true percentage of cases of tuberculosis of intestinal origin. These figures, however, do not give any satisfactory idea as to whether the bacilli entering the intestines originated from human or bovine sources. Owing to this fact, it follows that the only way of determining the infection of people by bacilli of the bovine type will be to study the lesions in the body of as many cases of human tuberculosis as is possible.

The finding of the bovine type of tubercle bacillus in human lesions is the most direct and positive proof that tuberculosis of cattle is responsible for a certain amount of tuberculosis in the human family. Numerous experiments with this object in view have already proven this fact. Thus the German commission on tuberculosis examined 16 different cultures obtained from tubercular lesions in children and found four which were more virulent than is usual for human tubercle bacilli, causing tuberculosis in the cattle inoculated with them, and making 25 per cent. of the cases tested, affected with a form of tuberculosis which, by Koch's own method, must be classified as of bovine origin. In a similar series of tests conducted by the British royal commission on tuberculosis, 60 cases of the disease in the human were tested, with the result that 14 cases were claimed by this commission to have been infected from bovine sources. Ravenel reports that of 5 cases of tuberculosis in children, 2 received their infection from cattle.

Theobald Smith has also reported on one culture of the bovine tubercle bacillus obtained from the mesenteric glands of a child, out of 5 cases examined, and according to a recent paper by Goodale, Smith has recently been at work on 7 other cultures from different children, 4 of which conformed to his idea of tubercle bacilli emanating from cattle. Of 4 cases of generalized tuberculosis in children examined in the Biochemic Division of the Bureau of Animal Industry, 2 were found to be affected with very virulent organisms, which warranted the conclusion that such children had been infected from bovine source. The Pathological Division of the same bureau has likewise, out of the 9 cases of infantile tuberculosis examined, obtained two cultures of tubercle bacilli that could not be differentiated from bovine culture. In Europe so many similar instances of bovine tubercle bacilli having been recovered from human tissues are on record that it appears to your committee entirely proven that man is susceptible to tuberculosis caused by animal infections, and while the proportion of such cases can not be decided with even approximate accuracy, it is nevertheless incumbent on us to recommend such measures as will guard against these sources of danger, when enforced.

The two principal sources of infection from cattle and the only ones necessary to be considered are the meat and milk of tuberculous animals. The fact that most of the cases of bovine tuberculosis above enumerated which occurred in the human were cases of infantile tuberculosis points with grave suspicion to the milk rather than the meat supply. This naturally leads to the question of how and under what condition does the milk become infectious. That milk coming from a tuberculous udder is capable of transmitting the infectious principle is conceded by all who have given the subject any consideration. It has been equally established that in advanced generalized tuberculosis the udder may secrete tubercle bacilli without showing any indication of being affected.

Careful experiments performed by trained and eminently responsible investigators have also demonstrated beyond reasonable doubt that tubercle bacilli at certain times may be present in the milk of cows affected with tuberculosis to a

degree that can be detected only by the tuberculin test, so that in a herd of cows in the various stages of tuberculosis it is to be expected that some of them will secrete tuberculous milk, which, when mixed with other cows' milk, makes the entire product dangerous. Even when the tubercle bacilli are not being excreted by the udder, it has been shown by Eber in Germany and Schroeder in this country, that the dust and manure of the stable where the diseased animals are kept are in many cases contaminated with tubercle bacilli.

In a recent examination of the manure passed by twelve cows just purchased from dairy farms in this city and affected with tuberculosis to an extent only demonstrable by the tuberculin test, tubercle bacilli were found in over 41 per cent. of the cases both by microscopic examination and animal inoculations. The danger from this method of infecting milk is impressed on us as consumers when we realize that on an average probably 25 per cent. of all the cows which supply milk to the district are tuberculous, and this is considered to be a conservative estimate, judging from the results of recent tuberculin tests. Vital statistics show that 14 out of every 100 people that die succumb to tuberculosis, while of the remaining 86 more than one-half show lesions of tuberculosis on post-mortem examination, although dying from some other cause.

The statement of von Behring above mentioned is particularly pertinent in regard to the relationship of human tuberculosis to the milk supply, and especially in connection with the results of those investigators who have studied market milk and found from 5.2 to 55 per cent. of the samples examined to contain tubercle bacilli. LeBlanc considers the milk of tuberculous cows dangerous even when bacilli are not present, on account of the toxin it contains. Michellazzi has injected such milk into tuberculous animals and obtained a reaction. To eliminate all tuberculous cattle from the herd, or to pasteurize all milk coming from untested cattle, should therefore be the object of all producers of milk and the sanitarians will be remiss in their whole duty should they neglect to guard against the products of tuberculous animals in their attempts to eradicate tuberculosis from man.

*Other Diseases.*—Foot-and-mouth disease, anthrax, cow-pox, and rabies may be transmitted by the milk, and while no case of actinomycosis in man has been traced to the use of milk it is desirable that the use of milk of cows so affected should be prohibited, especially when the disease affects the udder. Such conditions as gastroenteritis and septic and febrile conditions may render the milk injurious to the consumer. Local diseases of the udder like Botryomycosis, mastitis, mastitis, etc., may render the milk unwholesome, especially when pus organisms are present. The color, taste and odor of milk may be altered so as to make it unpalatable if not unwholesome; these changes may be due to the food of the cow or to bacterial changes in the milk after it is drawn from the udder. Milk may acquire poisonous properties from the food eaten by the cow. The milk produced shortly before or during the first five days after parturition should not be used.

*Recommendations.*—In view of the facts above enumerated, your committee desires to make the following recommendations:

1. All cows on dairy farms producing milk for the District of Columbia shall be tagged, tattooed or otherwise marked, for purpose of identification.
2. All milk produced on such dairy farms shall come either from tuberculin tested cattle, which shall be retested at least once a year, or be subjected to pasteurization under the supervision of the health department in case the herd is not tuberculin tested.
3. No additions to any herd, whether tested or not tested, shall be made in the future without subjecting such cattle to the tuberculin test.
4. No license shall in future be granted except to applicants having herds free from tuberculosis.
5. The milk of cattle showing any of the udder affections above mentioned, anthrax, rabies, gastroenteritis, septic conditions or clinical symptoms of tuberculosis, shall not be utilized as human food, even though the milk be pasteurized. Milk from cows 15 days before and 5 days after parturition, and

that from animals receiving any of the deleterious medications of foodstuffs previously mentioned shall likewise be excluded.

6. That the veterinary inspectors of the health department make frequent visits to dairies having untested herds, in order that they may discover all advanced cases of tuberculosis, or udder tuberculosis, as early as possible.

#### MILK SEDIMENTS OR DIRTY MILK IN RELATION TO DISEASE.

BY GEORGE M. KOBER, M.D.

Professor of Hygiene, School of Medicine, Georgetown University.

Every consumer of milk has doubtless observed the presence of more or less foreign matter at the bottom of the bottle in which it is kept; indeed, it is a matter of such common occurrence that it hardly excites our attention, and many are disposed to look on it as a matter of course. Professor Soxhlet of Munich was perhaps the first to point out that these deposits are largely made up of the excrementitious matter from the cow, which adhering to the udder of the animal, gained access to the bucket during the act of milking.

If these sediments are subjected to microscopical examination we will find that they are composed of epithelial debris, hairs of the cow, organic and inorganic dust particles, excrementitious matter, vegetable fibers, bacteria, fungi and spores of every description: fully 90 per cent. of the germs are fecal bacilli—all of which is not only disgusting but extremely suggestive of danger. The number of micro-organisms is largely increased and we know that under a suitable temperature bacterial development and consequent decomposition are materially hastened in such a medium. The greatest danger from milk of this class is the possible presence of ptomaines or toxins produced by certain saprophytic germs from the albuminoids of the milk.

The views of Vaughan, Booker, Escherich, Harrington, Kober and others on the relation of saprophytic germs and toxins to cholera infantum and the summer diarrheas in bottle-fed children are gaining ground and will doubtless lead to great reform in the management of dairies. It may be urged that all such modern innovations involve unnecessary hardship, but it should be remembered that by attacking all the various factors concerned in the causation of the disease we may hope for the best results. The reduction in the general mortality in the registration area of the United States from 19.6 in 1890 to 16.2 per 1,000 shows what may be accomplished by preventive medicine and sanitation.

*Milk in Relation to Infectious Diseases.*—Milk may acquire infective properties after it leaves the udder of the animal. Numerous instances have been observed in which outbreaks of typhoid fever, scarlet fever and diphtheria, by their sudden and explosive character, affecting families living in streets and localities supplied by the same milkman, naturally pointed to the milk supply as a common cause. Dr. Michael Taylor, however, was the first physician (in 1857) to point out definitely that cow's milk might serve as the medium of spreading typhoid fever from a dairy where the disease prevailed. In 1867 he also showed that scarlet fever might be distributed in the same way. In 1877 Mr. Jacob traced a diphtheria epidemic at Sutton to the milk supply, and in 1872 MacNamara traced an outbreak of cholera at Calcutta to an infected dairy. These facts could not fail to sharpen the powers of observation in others, so that in consequence similar outbreaks were more frequently reported. I was enabled to present to the International Medical Congress held in Paris in 1900 the history of 195 outbreaks of typhoid fever, ninety-nine of scarlet fever and six of diphtheria, all traceable to milk supply.

It has been demonstrated that disease germs may not only survive, but in many instances actually proliferate, in the milk, and it is not a difficult matter to point out the many ways by which these germs gain access, especially when some of the employees are also engaged in nursing the sick, or are suffering themselves from some mild infection while continuing their duties, or are convalescent from the disease, and thus infect the milk in handling it.

After the recital of numerous epidemics and milk-borne diseases, I need hardly insist on the necessity of compulsory notification of all infectious diseases, and that milk should not be permitted to leave a farm, dairy or milk-shop during the existence of any of these diseases among the inmates or employees, nor should the latter be permitted to reside in or visit infected houses while engaged in the milk traffic without permission from the health department. The farmer or retailer should, in fact, be prepared by previous instructions to guard the milk supply from these sources of danger and call on the authorities for an immediate inspection. To prevent great loss incident to those restrictions, they may be modified so as to use the milk after proper sterilization under the direction of the health department. There is nothing strained in the requirements of the proposed legislation and by their enforcement we may hope to obtain such a standard of milk as will not only effect a decided reduction in infantile mortality, but will render the dissemination of infectious diseases through the milk supply a matter of history only. Until this is accomplished, we should patronize only such dealers as sell "certified milk," or subject the milk to pasteurization, or simply bring the milk to the boiling point in the household, and, after cooling, keep it on ice: this will not make bad milk good, but it will at least destroy its infectiousness.

#### SANITARY INSPECTION OF DAIRIES AND DISTRIBUTING DEPOTS.

BY E. H. WEBSTER.

Chief of Dairy Division, Bureau of Animal Industry.

A systematic sanitary inspection of dairy farms and milk-distributing depots are anything but ideal; they are, in fact, about as bad as it is possible to conceive. Stables are poorly lighted, many having no windows whatever, and ventilation is left to care for itself. Little attention is paid to floors, ceilings, walks, or stable yards. Swine, horses and poultry are often found in the same barn with the cows. Manure is not removed, or, when removed, is thrown through an opening in the wall or just outside the door, frequently near the milk room.

The necessary appliances for sterilizing and cooling in milk room are often lacking, making it impossible properly to wash and sterilize pails, cans, bottles and other appliances, or to properly cool and hold at a low temperature the milk before delivery. Milk dealers, as a rule, have more respect for sanitation and have better appliances than the average farm, but some common practices are deplorable. Very few have appliances for sterilizing bottles.

The situation in Washington is not different from that confronting the health departments of most of the cities of any size throughout the country. The public is gradually awakening to the fact that these conditions must be changed. In order to bring about these changes within the district, it is recommended that:

1. A sufficient number of inspectors shall be employed so that each inspector shall have not over 100 farm dairies.
2. These men shall have technical training in the production and handling of milk.
3. The health officers shall have full authority to make rules and regulations, and to enforce the same, so as to safeguard the milk supply of the district from contamination through carelessness, ignorance or malicious intent.
4. The health officer, or any authorized inspector, shall have authority to revoke instantly the license or right to sell milk in the city if the provisions in these regulations are not complied with, where, in his judgment, such violation endangers the health of the consumer.

#### WATER SUPPLY OF DAIRY FARMS.

BY G. LLOYD MAGRUDER, M.D.

Emeritus Professor, Materia Medica and Therapeutics, School of Medicine, Georgetown University.

The influence of a polluted water supply has long been recognized as a most potent cause of disease. Many virulent epidemics have been directly traced to this origin. The literature