

ART. XIX.—*Correspondence of JOACHIM BARRANDE, SIR WILLIAM LOGAN and JAMES HALL, on the Taconic System and the age of the Fossils found in the Rock of Northern New England, and the Quebec Group of Rocks.*

I. INTRODUCTORY REMARKS.

As some of our foreign readers may not be acquainted with the question to which the following important correspondence relates, we think it advisable to make a few explanatory observations by way of introduction. A complete history of the whole subject would require a greater amount of space than can be afforded and we shall therefore touch only upon a few of the more salient points.

The rocks under discussion occupy a belt of country east and west from twenty to sixty miles wide, stretching from the vicinity of the city of New York in a northerly direction to Lake Champlain and thence through Vermont and Lower Canada to Cape Gaspé at the mouth of the St. Lawrence. The strata consisting of slates, limestones, sandstones and conglomerates are greatly disturbed, plicated and dislocated, and are often, especially along the eastern side of the belt, in a highly metamorphic condition. On this side they are overlaid unconformably by upper Silurian and Devonian rocks, but on the western and northern margin they are in contact with and in general seem to be a continuation of the Lower Silurian. Some of the slates of the formation closely resemble in lithological characters those of the Hudson River group and thus along the western side of the region, where the junction of the two formations occurs, it is often almost impossible to draw the line between them. The dip and strike of both are in the same direction and throughout extensive areas the newer rocks appear to plunge beneath the older. The whole district affords an excellent example of those cases, so well known to field geologists, where the true relations of the different masses cannot be clearly worked out without the aid of fossils and where the best of observers may arrive at diametrically opposite opinions.

Dr. Emmons, one of the geologists of the New York Survey, early convinced himself by a careful examination of these rocks, that they constituted a distinct physical group more ancient than the Potsdam sandstone, the latter being regarded by him as the base of the Lower Silurian System in North America. His views were given in detail in 1842 in his final report on that part of the State confided to his charge, and in a more special manner in another work entitled "THE TACONIC SYSTEM," published in 1844. In this latter work he figured several species of fossils which had been collected in different parts of the formation. Two of these were trilobites and were described under

the names of *Atops trilineatus* and *Elliptocephala asaphoides*. The others were graptolites, fucoides and apparently trails of annelides; he considered all the species to be distinct from any that had been found in American rocks of undoubted Silurian age. The pre-silurian age of the formation has also been maintained by him in several more recent publications such as his "American Geology"—the several reports on the geological survey of North Carolina and in his "Manual of Geology."

On the other hand, Professor Hall placed the whole region in the Hudson River group. In the first volume of the Palæontology of New York he identifies *Atops trilineatus* with *Triarthrus Beckii* the characteristic trilobite of the Utica slate;—*Elliptocephala asaphoides* he refers to the genus *Olenus* and describes as congeneric therewith another trilobite (*O. undulostriatus*) said to be from the true Hudson River shales. It is scarcely necessary to state that these identifications have always afforded an extremely powerful objection against the correctness of the position assumed by Emmons, because no species of trilobite is known to range from the Primordial Zone up to the top of the Lower Silurian. Hall's first volume was published in 1847 and as it is unquestionably the most important work on the Lower Silurian Fossils of North America it has been very generally accepted by our physical geologists as a guide. It is not surprising therefore that, in all the discussions that have taken place during the last fourteen years upon the age of these rocks, the majority of those who did not profess to be naturalists should have arranged themselves on the side of the leading Palæontologist of the country.

The formation was traced from New York through Vermont and there identified, by Prof. Adams the State Geologist, with the Hudson River group. The Canadian Surveyors continued it with great labor through a mountainous and partially uninhabited country for nearly five hundred miles further, from the northern extremity of Vermont to the neighborhood of Quebec and thence along the south side of the St. Lawrence to the mouth of that river at Cape Gaspé. In Canada the nomenclature of the New York Survey was adopted for all the formations and it appears from his several reports that Sir W. E. Logan could find nothing in the physical structure of the country to authorize him to make an exception in favor of this particular series of rocks. It has therefore always been called the Hudson River group in the publications of the Canadian Survey.

It will be seen by the following correspondence that the new light thrown upon the question of the age of these rocks by the fortunate discovery of a large number of fossils near Quebec, now leads him to place them at the base of the Lower Silurian, and as he states that the shales in Vermont in which the trilobites no-

ticed in Mr. Barrande's letter to Prof. Bronn have been found may be subordinate to the Potsdam it seems probable that the sequence contended for by Emmons will turn out to be at least for the greater part the true one.

II.

ON THE PRIMORDIAL FAUNA AND THE TACONIC SYSTEM OF EMMONS, IN A LETTER TO PROF. BRONN OF HEIDELBERG.*

"Paris, July 16, 1860.

"..... I have recently received, thanks to the kindness of Mr. E. Billings, the learned palæontologist of the Geological Survey of Canada, a very interesting pamphlet entitled 'Twelfth Annual Report of the Regents of the University of the State of New York, 1859.' If you possess this publication, you will find there, at page 59, a memoir of Prof. J. Hall, entitled 'Trilobites of the Shales of the Hudson River Group.' This savant there describes three species under the names *Olenus Thompsoni*, *Olenus Vermontana*, and *Peltura (Olenus) holopyga*. The well-defined characters of these trilobites are described with the clearness and precision to be expected from so skillful and experienced a palæontologist as James Hall.

"Although the specimens are incomplete, their primordial nature cannot admit of the least doubt, when the descriptions are read, accompanied with wood engravings which the large dimensions of these three species render sufficiently exact. The first is 105 millim. long by 80 broad, the other two are somewhat smaller.

"The heads of the two *Oleni* being deteriorated, the furrows of the glabella cannot be recognized. The thorax has a common and remarkable character, which consists in the greater development of the third segment, the point of which is stronger and longer than in all the other pleura. This is a striking resemblance to the *Paradoxides*, the second segment of which has the same peculiarity. Besides, there is an intimate relation between these two primordial types, and we should not be surprised if America furnished us with forms uniting most of their characteristics. The pygidium of *O. Thompsoni*, the only one that is known, shows no segmentation, and attests by its exiguity its relation to a primordial trilobite. *P. holopyga*, by its whole appearance, resembles the species of Sweden so well known by the name of *P. Scarabæoides*.

"Thus all the characters of these three trilobites, as they are recognized and described by J. Hall, are those of the trilobites of the primordial fauna of Europe. This is so true, that I think I may say without fear, if M. Angelin, or any other palæontologist practised in distinguishing the trilobites of Scandinavia, had met with these three American forms in Sweden or Norway, he would not have hesitated to class them among the species of the Primordial fauna, and to place the schists enclosing them in one of the formations containing this fauna. Such is my profound conviction, and I think any one who has made a serious study of the trilobitic forms and of their vertical distribution in the oldest formations will be of the same opinion.

* Proceed. Boston S. N. Hist., vol. vii, Dec., 1860, p. 371.

"Besides, all who have seriously studied palæontology know well that each geological epoch, or each fauna, has its proper and characteristic forms, which once extinct reappear no more. This is one of the great and beautiful results of your immense researches, which have generalized this law, recognized by each one of us within the limits of the strata he describes.

"The great American palæontologist arrived long since at the same conclusion, for in 1847 he wrote the following passage in the *Introduction* to the first volume of the Monumental Work consecrated to the Palæontology of New York.

"Every step in this research tends to convince us that the succession of strata, when clearly shown, furnishes conclusive proofs of the existence of a regular sequence among the earlier organisms. We are more and more able, as we advance, to observe that the Author of nature, though always working upon the same plan and producing an infinite variety of forms almost incomprehensible to us, has never repeated the same forms in successive creations. The various organisms called into existence have performed their parts in the economy of creation, have lived their period and perished. This we find to be as true among the simple and less conspicuous forms of the palæozoic series, as in the more remarkable fauna of later periods.'—*J. Hall, 'Pal. of New York,' i, p. xxiii.*"

"When an eminent man expresses such ideas so eloquently, it is because they rise from his deepest convictions. It must then be conceived that J. Hall, restrained by the artificial combinations of stratigraphy previously adopted by him, has done violence to his palæontological doctrines, when, seeing before him the most characteristic forms of the *Primordial fauna*, and giving them names the most significant of this first creation, he thinks it his duty to teach us that these three trilobites belong to a horizon *superior* to that on which the second fauna is extinguished.

"In effect, according to the text of J. Hall, the three trilobites in question were found near the town of Georgia, Vermont, in schists which are superior to the *true Hudson River group*. In his works J. Hall does not go beyond indicating the horizon of certain fossils, and no one would think of asking a guaranty for such indications. But on this occasion the great American palæontologist thinks it necessary to support his stratigraphical determination by another authority, chosen from the most respectable names in geology. The following is the note which terminates his Memoir.

"NOTE.—In addition to the evidence heretofore possessed regarding the position of the shales containing the Trilobites, I have the testimony of Sir W. E. Logan, that the shales of this locality are in the upper part of the Hudson River group, or forming a part of a series of strata which he is inclined to rank as a distinct group, above the Hudson River proper. It would be quite superfluous for me to add one word in support of the opinion of the most able stratigraphical geologist of the American continent."

"Now, when a savant like J. Hall thinks himself obliged to invoke testimony to guarantee the exactness of the position of a few fossils, it is clear that the determination of this position is difficult.

"In order to understand these difficulties I have consulted the maps and documents relating to the State of Vermont and the country in which the town of Georgia is situated, and, although the library of our Geological Society does not contain all that one could wish on this subject, I recognized easily that Georgia is placed in the region where the order of succession of the deposits is the most obscured by foldings and dislocations; so that the position of the schists in question could not have been determined by the incontestable evidence of direct superposition. Besides, the physical appearance of these schists is not that of the rocks constituting the typical group of Hudson River. This is verified by the Note of J. Hall, for it tells us that Sir W. E. Logan is inclined to make a distinct group of these schists *superior* to that of the Hudson, and which consequently *would crown the whole Lower Silurian division* of the continent.

"For the above reasons, the geological horizon on which the three *Oleni* of Georgia were found appears to me, at first view, to have been but doubtfully determined, and in complete opposition to palæontological documents.

"I do not think, then, that I weaken in the least degree the respect and confidence justly inspired by the labors of the American savants whose names have just been mentioned, when I ask them in the name of science to make new researches and new studies, that may lead to a final and certain solution of this important question.

"Doubtless, thanks to the progress of our knowledge, we are now no longer bound by the ancient conception of the simultaneous extinction and the total renovation of the faunæ. As for myself, in particular, it would not be possible to accuse me of similar views at the moment when I am publishing the explanation of my doctrine of colonies. But you will perceive that the facts which I invoke in support of this doctrine are far from sustaining the reappearance of a fauna after the extinction of the following fauna, which the three trilobites of Georgia would do, if they had really lived after the deposit of the Hudson River group.

"This reappearance would be still more astonishing, as among the three great Silurian faunæ the second fauna occupies the greatest vertical space and is probably the one which enjoyed the longest existence. Thus, to verify such a reappearance, the most incontestable proofs are required, for such a decision would compel the entire re-formation of one of our most important scientific creeds.

"Yours very truly,

J. BARRANDE."

In another letter, dated Paris, 14th August, 1860, Mr. Barrande says:—

"You will easily perceive the interest and importance of the question, even if it were only raised on account of the three *Oleni* of Georgia; but it takes in now a much wider field, owing to a letter I have just received from Mr. Billings, official Palæontologist of the Geological Survey of Canada, who informs me that he has found lately, in the schists and limestones near Quebec, considered as being the prolongation of those in question in Vermont, nearly one hundred species, almost all new. Twenty-six of these come from a white limestone, and seem to him to be the true representatives of the Primordial fauna, and he cites among them *Conocephalites*,

Arionellus, *Dikellocephalus*, etc., that is, very characteristic forms of this fauna.

"In another limestone, which is gray, he finds thirty-nine species, all different from the first, and representing, on the contrary, the most distinct types of the second fauna. Finally, the black schists furnish him with *Graptolites*, *Lingulæ*, etc., etc., fossils which at first sight cannot determine a horizon, because they are found upon several Silurian horizons.

"While waiting for these very obscure stratigraphical relations to be disentangled, and without committing in any manner Mr. Billings, who should preserve the independence of his opinion, I may yet express to you my view wholly personal, and of which at this moment I take the entire responsibility. I think, then, that this region of schists and limestones of Vermont, in other words the *Taconic system*, will reproduce in America what took place in England as to the Malvern Hills, and in Spain for the Cantabrian chain,—that is to say, the Primordial fauna, after having been disregarded, will regain its rights and its place, usurped for a time by the second fauna.

"You see it is a great and noble question, whose final solution will complete the imposing harmonies existing already between the series of palæozoic faunæ of America and that of the contemporaneous faunæ of Europe, leaving to each the imprint peculiar to its continent.

"I can well imagine, from the position previously taken by our learned American brethren on the subject of the Taconic system, that the final solution of which I speak will not be obtained without debate, and perhaps some wounding of self-love, for some opinions that appear to be dominant must be abandoned.

"But experience has taught me that in such cases the most elevated minds turn always first to the light, and put themselves at the head of the movement of reform. Thus, when in 1850 I recognized the Primordial fauna in the Malvern Hills, where the second fauna only had been found, Sir Henry de la Beche and Sir Roderick Murchison were the first to adopt my views, to which little by little the other official geologists agreed; Edward Forbes ranged himself publicly on my side in 1853 in 'The Geological Survey,' while others still hesitated, until now there is no longer any opponent.

"I think there will be the same experience in America, and that in a few years from this time the opinions of your savans will have undergone a great change as regards this question.

"It is a fine opportunity for Dr. Emmons to reproduce his former observations and ideas with more success than in 1844.

"Yours very truly,

J. BARRANDE."